

**MOTION SICKNESS RELIEF- meclizine hcl tablet, chewable**  
**Rite Aid Corporation**

-----  
**Rite Aid 44-404A**

***Active ingredient (in each chewable tablet)***

Meclizine HCl 25 mg

***Purpose***

Antiemetic

***Uses***

for prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness

***Warnings***

**Do not use**

for children under 12 years of age unless directed by a doctor.

**Ask a doctor before use if you have**

- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis

**Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers.

**When using this product**

- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### ***Directions***

- **do not take more than directed**
- dosage should be taken one hour before travel starts
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 1 to 2 chewable tablets once daily or as directed by a doctor

### ***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- protect from heat and humidity
- see end flap for expiration date and lot number

### ***Inactive ingredients***

colloidal silicon dioxide, dextrose, FD&C red #40 aluminum lake, flavor, lactose monohydrate, magnesium stearate, pregelatinized starch, propylene glycol, saccharin sodium, silicon dioxide

### ***Questions or comments?***

**1-800-426-9391**

### ***Principal Display Panel***

NDC 11822-4040-1

Compare to the active ingredient  
in **Bonine® Chewable Tablets\***

**MOTION SICKNESS RELIEF**  
MECLIZINE HCl 25 mg CHEWABLE TABLETS  
ANTIEMETIC  
PREVENTS MOTION SICKNESS  
Ages 12 Years and Over

RASPBERRY  
FLAVORED

**16** CHEWABLE TABLETS

ACTUAL SIZE

Chew or crush tablets completely  
before swallowing.

**TAMPER EVIDENT: DO NOT USE IF PACKAGE  
IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN  
OR SHOWS ANY SIGNS OF TAMPERING**

\*This product is not manufactured or distributed by WellSpring Pharmaceutical Corporation, owner of the registered trademark Bonine® Chewable Tablets.  
50844      ORG082340421

**DISTRIBUTED BY:** RITE AID,  
200 NEWBERRY COMMONS  
ETTERS, PA 17319  
[www.riteaid.com](http://www.riteaid.com)

**SATISFACTION  
GUARANTEE**

If you're not satisfied, we'll  
happily refund your money.

NDC 11822-4040-1

Compare to the active ingredient  
in Bonine® Chewable Tablets\*

# MOTION SICKNESS RELIEF

MECLIZINE HCl 25 mg CHEWABLE TABLETS

NDC 11822-4040-1

Compare to the active ingredient  
in Bonine® Chewable Tablets\*

# MOTION SICKNESS RELIEF

MECLIZINE HCl 25 mg CHEWABLE TABLETS

ANTIEMETIC  
PREVENTS MOTION SICKNESS

Ages 12 Years and Over



RASPBERRY  
FLAVORED

ACTUAL SIZE

# 16

CHEWABLE  
TABLETS

Chew or crush tablets completely  
before swallowing.



E1 042  
FLATFOLD

B-1702-404A-21-R3F  
08032340421

\*This product is not manufactured or distributed by  
WellSpring Pharmaceutical Corporation, owner of the  
registered trademark Bonine® Chewable Tablets.  
50844 ORG082340421

**Drug Facts (continued)**  
*Questions or comments? 1-800-426-9391*

**SATISFACTION  
GUARANTEE**  
If you're not satisfied, we'll  
happily refund your money.

no print / no varnish area  
lot no & exp date

**Drug Facts**  
KEEP OUTER PACKAGE FOR  
COMPLETE PRODUCT INFORMATION

**Active ingredient**  
(in each chewable tablet)  
Meclizine HCl 25 mg.....Antiemetic

**Warnings**  
Do not use for children under 12 years of age unless  
directed by a doctor.  
Ask a doctor before use if you have  
■ difficulty in urination due to enlargement of the prostate  
gland ■ glaucoma  
■ a breathing problem such as emphysema or chronic  
bronchitis  
Ask a doctor or pharmacist before use if you are  
taking sedatives or tranquilizers.

**When using this product** ■ drowsiness may occur  
■ alcohol, sedatives, and tranquilizers may increase  
drowsiness ■ avoid alcoholic beverages  
■ use caution when driving a motor vehicle or operating  
machinery

**Drug Facts (continued)**

**Purpose**  
If pregnant or breast-feeding, ask a health professional  
before use.  
Keep out of reach of children. In case of overdose, get  
medical help or contact a Poison Control Center right  
away.

**Directions** ■ do not take more than directed  
■ dosage should be taken one hour before travel starts  
■ chew or crush tablets completely before swallowing;  
do not swallow tablets whole  
■ adults and children 12 years and over: take 1 to 2  
chewable tablets once daily or as directed by a doctor

**Other information**  
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE  
IS OPENED OR BLISTER IS TORN OR BROKEN  
■ store at 25°C (77°F); excursions permitted between  
15°-30°C (59°-86°F) ■ protect from heat and humidity  
■ see end flap for expiration date and lot number

**Inactive ingredients** colloidal silicon dioxide,  
dextrose, FD&C red #40 aluminum lake, flavor, lactose,  
monohydrate, magnesium stearate, pregelatinized starch,  
propylene glycol, saccharin sodium, silicon dioxide



DISTRIBUTED BY: RITE AID,  
200 NEWBERRY COMMONS  
ETTERS, PA 17319  
www.riteaid.com

TAMPER EVIDENT: DO NOT USE IF PACKAGE  
IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN  
OR SHOWS ANY SIGNS OF TAMPERING

## Rite Aid 44-404A

### MOTION SICKNESS RELIEF

meclizine hcl tablet, chewable

#### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-4040
Route of Administration	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

#### Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DEXTROSE, UNSPECIFIED FORM (UNII: IY9XDZ35W2)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

#### Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	9mm
Flavor	RASPBERRY	Imprint Code	44;404
Contains			

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-4040-1	2 in 1 CARTON	11/10/2023	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	11/10/2023	

**Labeler** - Rite Aid Corporation (014578892)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-4040) , pack(11822-4040)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-4040)

Revised: 11/2024

Rite Aid Corporation