

MOTION SICKNESS RELIEF- meclizine hcl tablet, chewable
Rite Aid Corporation

Rite Aid 44-404A

Active ingredient (in each chewable tablet)

Meclizine HCl 25 mg

Purpose

Antiemetic

Uses

for prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness

Warnings

Do not use

for children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- dosage should be taken one hour before travel starts
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 1 to 2 chewable tablets once daily or as directed by a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- protect from heat and humidity
- see end flap for expiration date and lot number

Inactive ingredients

colloidal silicon dioxide, dextrose, FD&C red #40 aluminum lake, flavor, lactose monohydrate, magnesium stearate, pregelatinized starch, propylene glycol, saccharin sodium, silicon dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

NDC 11822-4040-1

Compare to the active ingredient
in **Bonine® Chewable Tablets***

MOTION SICKNESS RELIEF
MECLIZINE HCl 25 mg CHEWABLE TABLETS
ANTIEMETIC
PREVENTS MOTION SICKNESS
Ages 12 Years and Over

RASPBERRY
FLAVORED

16 CHEWABLE TABLETS

ACTUAL SIZE

Chew or crush tablets completely
before swallowing.

**TAMPER EVIDENT: DO NOT USE IF PACKAGE
IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

*This product is not manufactured or distributed by WellSpring Pharmaceutical Corporation, owner of the registered trademark Bonine® Chewable Tablets.
50844 ORG082340421

DISTRIBUTED BY: RITE AID,
200 NEWBERRY COMMONS
ETTERS, PA 17319
www.riteaid.com

**SATISFACTION
GUARANTEE**

If you're not satisfied, we'll
happily refund your money.

NDC 11822-4040-1

Compare to the active ingredient
in Bonine® Chewable Tablets*

MOTION SICKNESS RELIEF

MECLIZINE HCl 25 mg CHEWABLE TABLETS

NDC 11822-4040-1

Compare to the active ingredient
in Bonine® Chewable Tablets*

MOTION SICKNESS RELIEF

MECLIZINE HCl 25 mg CHEWABLE TABLETS

ANTIEMETIC
PREVENTS MOTION SICKNESS

Ages 12 Years and Over



RASPBERRY
FLAVORED

ACTUAL SIZE

16

CHEWABLE
TABLETS

Chew or crush tablets completely
before swallowing.



E1042
FLATFOLD

B-1702-404A-21-R3F
08032340421

*This product is not manufactured or distributed by
WellSpring Pharmaceutical Corporation, owner of the
registered trademark Bonine® Chewable Tablets.
50844 ORG082340421

Drug Facts (continued)
Questions or comments? 1-800-426-9391

**SATISFACTION
GUARANTEE**
If you're not satisfied, we'll
happily refund your money.

no print / no varnish area
lot no & exp date

Drug Facts
KEEP OUTER PACKAGE FOR
COMPLETE PRODUCT INFORMATION

Active ingredient
(in each chewable tablet)
Meclizine HCl 25 mg.....Antiemetic

Uses for prevention and treatment of nausea, vomiting,
or dizziness associated with motion sickness

Warnings
Do not use for children under 12 years of age unless
directed by a doctor.
Ask a doctor before use if you have
■ difficulty in urination due to enlargement of the prostate
gland ■ glaucoma
■ a breathing problem such as emphysema or chronic
bronchitis

Ask a doctor or pharmacist before use if you are
taking sedatives or tranquilizers.

When using this product ■ drowsiness may occur
■ alcohol, sedatives, and tranquilizers may increase
drowsiness ■ avoid alcoholic beverages

■ use caution when driving a motor vehicle or operating
machinery

Drug Facts (continued)

Purpose
If pregnant or breast-feeding, ask a health professional
before use.
Keep out of reach of children. In case of overdose, get
medical help or contact a Poison Control Center right
away.

Directions ■ do not take more than directed
■ dosage should be taken one hour before travel starts
■ chew or crush tablets completely before swallowing;
do not swallow tablets whole

■ adults and children 12 years and over: take 1 to 2
chewable tablets once daily or as directed by a doctor

Other information
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE
IS OPENED OR BLISTER IS TORN OR BROKEN
■ store at 25°C (77°F); excursions permitted between
15°-30°C (59°-86°F) ■ protect from heat and humidity
■ see end flap for expiration date and lot number

Inactive ingredients colloidal silicon dioxide,
dextrose, FD&C red #40 aluminum lake, flavor, lactose,
monohydrate, magnesium stearate, pregelatinized starch,
propylene glycol, saccharin sodium, silicon dioxide



DISTRIBUTED BY: RITE AID,
200 NEWBERRY COMMONS
ETTERS, PA 17319
www.riteaid.com

TAMPER EVIDENT: DO NOT USE IF PACKAGE
IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING

Rite Aid 44-404A

MOTION SICKNESS RELIEF

meclizine hcl tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-4040
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: 008PUV3986) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DEXTROSE, UNSPECIFIED FORM (UNII: IY9XDZ35W2)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	9mm
Flavor	RASPBERRY	Imprint Code	44;404
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-4040-1	2 in 1 CARTON	11/10/2023	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	11/10/2023	

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-4040) , pack(11822-4040)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-4040)

Revised: 11/2025

Rite Aid Corporation