

**IBUPROFEN- ibuprofen capsule, liquid filled**  
**Cardinal Health (Leader) 70000**

-----

**Drug Facts**

**Active ingredient (in each capsule)**

**Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\***

**(present as the free acid and potassium salt)**

**\*nonsteroidal anti-inflammatory drug**

**Purpose**

**Pain reliever/fever reducer**

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - slurred speech
  - trouble breathing
  - leg swelling
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- **do not take more than directed**

- **the smallest effective dose should be used**
- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

#### **Other information**

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F)
- swallow whole; do not crush, chew, or dissolve

#### **Inactive ingredients**

FD&C green #3, gelatin, medium-chain triglycerides, polyethylene glycol, potassium hydroxide, purified water, sorbitan solution, sorbitol, white ink

#### **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

#### **Principal Display Panel**

**Ibuprofen** capsules, 200 mg

Pain Reliever/ Fever Reducer (**NSAID**)

COMPARE TO ADVIL® LIQUI-GELS® active ingredient\*\*

SOFTGELS<sup>†</sup>

(<sup>†</sup>Liquid Filled Capsules)

**TAMPER-EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

\*\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liquid Gels®.

DISTRIBUTED BY CARDINAL HEALTH, DUBLIN, OHIO 43017

www.myleader.com 1-800-200-6313

#### **Product Label**

Exp. Date: Lot No.: Made in China  
0 96295-13405 6  
CN 5425764 REV. 3/18  
Return to place of purchase if not satisfied.

ALL LEADER® Brand Products Have A 100% Money Back Guarantee

CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health, DISTRIBUTED BY CARDINAL HEALTH, DUBLIN, OHIO 43007  
www.mylender.com 1-800-200-5555

Cardinal Health, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health, DISTRIBUTED BY CARDINAL HEALTH, DUBLIN, OHIO 43007

**Drug Facts (continued)**  
■ slurred speech ■ trouble breathing ■ leg swelling  
■ weakness in one part or side of body ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear  
**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** ■ do not take more than directed  
■ the smallest effective dose should be used  
■ adults and children 12 years and over: ■ take 1 capsule every 4 to 6 hours while symptoms persist  
■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

**Other information** ■ each capsule contains: potassium 20 mg ■ read all warnings and directions before use. Keep carton. ■ store at 20° to 25°C (68° to 77°F) ■ avoid excessive heat above 40°C (104°F) ■ swallow whole; do not crush, chew, or dissolve

**Drug Facts (continued)**  
product ■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery  
**Ask a doctor before use if** ■ the stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic  
**Ask a doctor or pharmacist before use if you are** ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug  
**When using this product**, take with food or milk if stomach upset occurs.  
**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain

**Drug Facts (continued)**  
**Inactive ingredients** FD&C green #3, gelatin, medium-chain triglycerides, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution, white ink

**Questions or comments?**  
Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

"This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®."

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.** PLD-A524A FC004785

**Drug Facts**  
**Active ingredient (in each capsule)**  
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*  
(present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**Purpose**  
Pain reliever/fever reducer

**Uses**  
■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever

**Warnings**  
**Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this

NDC 70000-0360-1

**LEADER**

**Ibuprofen**  
Capsules, 200 mg  
Pain Reliever / Fever Reducer (NSAID)

**20 SOFTGELS\***  
(Liquid Filled Capsules)

**COMPARE TO ADVIL® LIQUI-GELS®**  
active ingredient\*\*

**100% Money Back Guarantee**

LEADER Ibuprofen 200 mg Capsules

IBUPROFEN			
ibuprofen capsule, liquid filled			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0360
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)			IBUPROFEN	200 mg
Inactive Ingredients				
Ingredient Name				Strength
GELATIN (UNII: 2G86QN327L)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)				
WATER (UNII: 059QF0KO0R)				
SORBITOL (UNII: 506T60A25R)				
SORBITAN (UNII: 6O92ICV9RU)				
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)				
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)				
Product Characteristics				
Color	GREEN	Score	no score	
Shape	CAPSULE	Size	19mm	
Flavor		Imprint Code	PC16	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0360-1	1 in 1 BOX	02/28/2018	
1		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:70000-0360-2	1 in 1 BOX	02/28/2018	
2		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:70000-0360-3	80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/28/2018	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA206568		02/28/2018	

**Labeler** - Cardinal Health (Leader) 70000 (097537435)