

**SEVERE NIGHTTIME COLD AND FLU RELIEF- acetaminophen dextromethorphan hbr doxylamine succinate phenylephrine hci liquid
Family Dollar (FAMILY WELLNESS)**

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal Decongestant

Uses

- temporarily relieves common cold/flu symptoms:
 - nasal congestion
 - sore throat
 - headache
 - sinus congestion and pressure
 - minor ache and pain
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation
- temporarily reduces
 - fever
 - cough to help you sleep
 - swelling of nasal passages
 - temporarily restores freer breathing through the nose
 - promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or last more than 3 days
- redness or swelling is present
- cough comes back, or occurs with rash or headache that lasts.

These could be a signs of a serious conditions.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **take only as directed - see Overdose warning**
- do not exceed 4 doses per 24 hours
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over: 30 mL every 4 hours
- children under 12 years of age: do not use
- **When using Daytime or Nighttime products, carefully read each label to ensure correct dosing.**

Other information

- **each 30 mL contains;** sodium 64 mg
- store between 20-25°C (68-77°F). Do not refrigerate

Inactive ingredients

anhydrous citric acid, FD&C blue1, FD&C red 40, Flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum

Questions or comments?

Call

Principal Display Panel

*Compare to the active ingredients of VICKS® NYQUIL® Severe Cold & Flu

Severe Nighttime

Cold & Flu Relief

Acetaminophen 650 mg

Pain Reliever/Fever Reducer

Dextromethorphan HBr 20 mg

Cough suppressant

Doxylamine succinate 12.5 mg

Antihistamine

Phenylephrine HCl 10 mg

Nasal decongestant

Relieves

- Headache, fever, sore throat, minor aches & pains
- Nasal/sinus congestion & sinus pressure
- Sneezing
- Runny nose
- Cough

Max strength

Alcohol free

FL OZ (mL)

Berry Flavor

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY

Product Label

NDC XXXXX-XXX-XX
*Compare to the active ingredients in Vicks® NyQuil® Severe Cold & Flu

Severe Nighttime Cold & Flu Relief

Acetaminophen 650 mg
Pain reliever/Fever reducer

Dextromethorphan HBr 20 mg
Cough suppressant

Doxylamine succinate 12.5 mg
Antihistamine

Phenylephrine HCl 10 mg
Nasal decongestant

Relieves
✓ Headache, fever, sore throat, minor aches & pains
✓ Nasal/sinus congestion & sinus pressure
✓ Sneezing
✓ Runny nose
✓ Cough

Max strength
Alcohol free

BERRY FLAVOR

PL-ANIC LB009767

8 fl oz (237 mL)

Drug Facts

Active ingredients (in each 30 mL)
Acetaminophen 650 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Doxylamine succinate 12.5 mg.....Antihistamine
Phenylephrine HCl 10 mg.....Nasal decongestant

Purposes

Uses
■ temporarily relieves common cold/flu symptoms
■ nasal congestion
■ sore throat
■ headache
■ sinus congestion and pressure

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

PLD-A341C LB009766

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

- minor aches and pains
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- temporarily reduces
 - fever
 - cough to help you sleep
 - swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening
■ blisters
■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ liver disease ■ heart disease
■ high blood pressure ■ thyroid disease
■ diabetes ■ glaucoma
■ trouble urinating due to an enlarged prostate gland
■ cough that occurs with too much phlegm (mucus)
■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ a sodium-restricted diet

Ask a doctor or pharmacist before use if you are
■ taking sedatives or tranquilizers
■ taking the blood thinning drug warfarin

Drug Facts (continued)

When using this product
■ do not take more than directed
■ excitability may occur, especially in children
■ marked drowsiness may occur
■ avoid alcoholic drinks
■ be careful when driving a motor vehicle or operating machinery
■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ new symptoms occur
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical

Drug Facts (continued)

help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions
■ take only as directed (see Overdose warning)
■ do not exceed 4 doses per 24 hours
■ measure only with dosing cup provided. Do not use any other dosing device.
■ mL = milliliter
■ keep dosing cup with product
■ adults and children 12 years and over: 30 mL every 4 hours
■ children under 12 years of age: do not use
■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information
■ each 30 mL contains: sodium 64 mg
■ store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

FAMILY WELLNESS Severe Nighttime Cold & Flu Relief Berry Flavor

SEVERE NIGHTTIME COLD AND FLU RELIEF

acetaminophen dextromethorphan hbr doxylamine succinate phenylephrine hci liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-641
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-641-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/31/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/31/2024	

Revised: 5/2024

Family Dollar (FAMILY WELLNESS)