

ASPIRIN- aspirin tablet, coated
Aarkish Pharmaceuticals NJ Inc.

ACTIVE INGREDIENT (IN EACH TABLET)

Aspirin 81 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

PURPOSE

Pain reliever

USES

temporarily relieves minor aches and pains

Because of its delayed action, this product will not produce fast relief of headaches or other symptoms needing immediate relief.

ask your doctor about other uses for enteric-coated 81 mg Aspirin

WARNINGS

Reye's syndrome:Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be early sign of Reye's syndrome, a rare but serious illness.

OTHER SAFETY INFORMATION

Allergy Alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naprozen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for longer time than directed

DO NOT USE

Do not use: if you are allergic to aspirin or any other pain reliever/fever reducer

ASK DOCTOR

Ask a doctor before use if

stomach bleeding warning applies to you
you have a history of stomach problems, such as heartburn
you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
you are taking a diuretic
you have asthma
you have not been drinking fluids
you have lost a lot of fluid due to vomiting or diarrhea

ASK A DOCTOR OR PHARMACIST

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug
- under a doctor's care for any serious condition

STOP USE

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
 - you are experience any of the following signs of stomach bleeding: feel faint, have bloody or black stools, vomit blood, have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - redness or swelling is present
 - new symptoms occur
 - ringing in the ears or a loss of hearing occurs.
- these could be signs of a serious condition.

PREGNANCY OR BREASTFEEDING

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

KEEP OUT OF REACH OF CHILDREN

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

DIRECTIONS

- Drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

STORAGE


- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- Use by expiration date on package

INACTIVE INGREDIENTS

corn starch, pregelatinized starch, povidone, microcrystalline cellulose, colloidal silicon dioxide, stearic acid, methacrylic acid and ethyl acrylate copolymer, talc, titanium dioxide, triethyl citrate, sodium bicarbonate, sodium lauryl sulfate, D&C yellow #10, hypromellose, triacetin.

PRINCIPAL DISPLAY PANEL

Aspirin Tablets, 81 mg, 300 counts



NDC 81005-103-30

ASPIRIN

ADULT LOW DOSE


Pain Reliever (NSAID)
Enteric Coated Aspirin Regimen
Delayed Release
Safe Pain Relief

81
MG


300
TABLETS

+Compare to the Active Ingredient in BAYER® Low Dose Aspirin®

MADE IN USA



†EagleHealth is not affiliated with the owner of the trademark, BAYER®. Manufactured by: AARKISH PHARMACEUTICALS NJ Inc. Rev. 01 Fairfield, NJ 07004 www.aarkishpharma.com



3 81005 10330 0

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts	<p>Active ingredient (in each tablet) Purpose Aspirin 81 mg (NSAID*) Pain reliever *nonsteroidal anti-inflammatory drug</p> <p>Uses ■ temporarily relieves minor aches and pains Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin</p> <p>Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed</p> <p>Do not use if you are allergic to aspirin or any other pain reliever/fever reducer</p>
	<p>UNVARNISHED AREA</p> <p>PEEL HERE FOR MORE DRUG FACTS</p> <p>LOT #:</p> <p>EXP. DATE:</p>


Drug Facts continued on back of label

Glue Area

Drug Facts (continued)	<p>Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma</p> <p>Ask a doctor or pharmacist before use if you are ■ taking a prescription drug for diabetes, gout, or arthritis</p> <p>Stop use and ask a doctor if ■ an allergic reaction occurs. Speak medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ vomit blood ■ have bloody or black stools ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present ■ new symptoms occur ■ ringing in the ears or a loss of hearing occurs</p> <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor</p> <p>Other information store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)</p> <p>Inactive ingredients colloidal silicon dioxide, corn starch, D&C yellow #10, hypromellose, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, povidone, pregelatinized starch, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate.</p>
-------------------------------	--

STOP PEELING

Aspirin Tablets, 81 mg, 1000 counts



NDC 81005-103-10

ASPIRIN

ADULT LOW DOSE

Pain Reliever (NSAID)
Enteric Coated Aspirin Regimen
Delayed Release
Safe Pain Relief

81
MG

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN


1000
TABLETS

NOT ACTUAL SIZE

Compare to the Active Ingredient in BAYER® Low Dose Aspirin®

MADE IN USA

†EagleHealth is not affiliated with the owner of the trademark BAYER®. Manufactured by: AARKISH PHARMACEUTICALS NJ Inc. Rev. 01 Fairfield, NJ 07004 www.aarkishpharma.com



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin 81 mg (NSAID)* Pain reliever
*nonsteroidal anti-inflammatory drug

Uses ■ Temporarily relieves minor aches and pains
Because of its delayed action, this product will not provide last relief of headaches or other symptoms needing immediate relief.
■ ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings
Reye's syndrome. Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Drug Facts continued on back of label

UNVARNISHED AREA

LOT #:
EXP. DATE:

PEEL HERE FOR MORE DRUG FACTS

Glue Area

Drug Facts (continued)

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are
■ taking a prescription drug for diabetes, gout, or arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ have bloody or black stools
■ vomit blood ■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ redness or swelling is present
■ new symptoms occur
■ ringing in the ears or a loss of hearing occurs
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
■ children under 12 years: consult a doctor

Other information store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients colloidal silicon dioxide, corn starch, D&C yellow #10, hypromellose, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, povidone, pregelatinized starch, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate.

STOP PEEING

ASPIRIN			
aspirin tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:81005-103
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg	
Inactive Ingredients			
Ingredient Name	Strength		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
STARCH, CORN (UNII: O8232NY3SJ)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			

METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	5mm
Flavor		Imprint Code	S;17
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81005-103-30	300 in 1 BOTTLE; Type 0: Not a Combination Product	03/18/2026	
2	NDC:81005-103-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	03/18/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/18/2026	

Labeler - Aarkish Pharmaceuticals NJ Inc. (117729250)

Revised: 3/2026

Aarkish Pharmaceuticals NJ Inc.