

**FOSTER AND THRIVE READY TO USE SALINE LAXATIVE- sodium phosphate, dibasic, unspecified form and sodium phosphate, monobasic, unspecified form enema
McKesson**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

**Foster & Thrive™ Ready to Use Saline Enema
Laxative**

Drug Facts

Active ingredients (in each 118mL delivered dose)	Purpose
Dibasic Sodium Phosphate 7 g	Saline laxative
Monobasic Sodium Phosphate 19 g	Saline laxative

Use

- relieves occasional constipation
- this product generally produces bowel movement in 1 to 5 minutes

Warnings

Dosage warning

Using more than one enema in 24 hours can be harmful.

For rectal use only.

Do not use

- on children under 2 years of age

Ask a doctor before use if you

- have already used a laxative for more than 3 days
- have kidney disease, have heart problems, or are dehydrated
- are 55 years of age or older
- are on a sodium-restricted diet
- have abdominal pain, nausea, or vomiting
- have a sudden change in bowel habits lasting more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

When using this product

- **do not use more than directed. Serious side effects may occur from excess dosage**
- do not use for more than 3 days, without asking a doctor

Stop use and ask a doctor if

- you have rectal bleeding
- you have no bowel movement within 30 minutes of enema use
- you have symptoms of dehydration (thirstiness, dizziness, vomiting, urinating less often than normal)

These symptoms may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away: 800-222-1222.

Directions (or as directed by a doctor)

Single daily dosage (per 24 hours) Do not use if taking another sodium phosphate product. Do not use more unless directed by a doctor. See Warnings.

Adults and children 12 years of age and older	1 bottle once daily
Children 2 to under 12 years of age	Use Pediatric Enema
Children under 2 years of age	DO NOT USE

Other information

- **each 118 ml contains** sodium 4.3 g
- store at 15 to 30°C (59 to 86°F)
- additional liquids by mouth are recommended while using this product
- **Tamper Evident:** DO NOT USE IF TOP OR BOTTOM FLAP OF CARTON IS TORN OR MISSING.

Inactive ingredients

benzalkonium chloride, disodium EDTA, purified water

Questions or comments?

833-358-6431

Distributed by: McKesson Corp., via Strategic

Sourcing Services LLC. Memphis, TN 38141

PRINCIPAL DISPLAY PANEL - 133 mL Bottle Carton

NDC 70677-1089-1

COMPARE TO FLEET®
SALINE ENEMA
ACTIVE INGREDIENTS*

Foster & Thrive™

READY TO USE
Enema
SALINE LAXATIVE

Lubricated Tip

Works In Minutes

Not Made With
Natural Rubber Latex

For Relief of:

- Occasional
Constipation

NOT ACTUAL SIZE
1 BOTTLE
4.5 FL OZ (133 mL)

Foster & Thrive™

READY TO USE

Enema

SALINE LAXATIVE

Lubricated Tip

Works In Minutes

Not Made With Natural Rubber Latex

NDC 70677-1089-1

COMPARE TO FLEET®
SALINE ENEMA
ACTIVE INGREDIENTS*

Foster & Thrive™

READY TO USE

Enema

SALINE LAXATIVE

Lubricated Tip

Works In Minutes

Not Made With

Natural Rubber Latex

For Relief of:

- Occasional Constipation



NOT ACTUAL SIZE
1 BOTTLE
4.5 FL OZ (133 mL)

Drug Facts

Active ingredients Purpose (in each 118mL delivered dose)

Dibasic Sodium Phosphate 7 g Saline laxative
Monobasic Sodium Phosphate 19 g Saline laxative

Use

- relieves occasional constipation
- this product generally produces bowel movement in 1 to 5 minutes

Warnings

Dosage warning: Using more than one enema in 24 hours can be harmful.

For rectal use only.

Do not use

- on children under 2 years of age
- if you have already used a laxative for more than 3 days
- if you have kidney disease, have heart problems, or are dehydrated
- if you are 55 years of age or older
- if you are on a sodium-restricted diet
- if you have abdominal pain, nausea, or vomiting
- if you have a sudden change in bowel habits lasting more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

When using this product

- do not use more than directed. Serious side effects may occur from excess dosage
- do not use for more than 3 days, without asking a doctor

Stop use and ask a doctor if

- you have rectal bleeding
 - you have no bowel movement within 30 minutes of enema use
 - you have symptoms of dehydration (thirstiness, dizziness, vomiting, urinating less often than normal)
- These symptoms may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away: 800-222-1222.

TAMPER RESISTANT: DO NOT USE IF TOP OR BOTTOM FLAP IS TORN OR MISSING.

CAUTION: REMOVE PROTECTIVE SHIELD BEFORE INSERTING.

*This product is not manufactured or distributed by C.B. Fleet Company Inc., owner of the registered trademark Fleet®.

©2023 McKesson Corporation
Distributed by: McKesson Corp., via Strategic Sourcing Service LLC, Memphis, TN 38141
Money Back Guarantee
www.fosterandthrive.com

Drug Facts (continued)

Directions (or as directed by a doctor)
Single daily dosage (per 24 hours)
Do not use if taking another sodium phosphate product.
Do not use more unless directed by a doctor. See Warnings.

Adults and children 12 years of age and older	1 bottle once daily
Children 2 to under 12 years of age	Use Pediatric Enema
Children under 2 years of age	DO NOT USE

Other information

- each 118 mL contains sodium 4.3 g
- store at 15 to 30°C (59 to 86°F)
- additional liquids by mouth are recommended while using this product
- Tamper Evident: DO NOT USE IF TOP OR BOTTOM FLAP OF CARTON IS TORN OR MISSING.**

Inactive ingredients

benzalkonium chloride, disodium EDTA, purified water

Questions or comments?

833-358-6431

Complete Enema In A Disposable Squeeze Bottle With Soft, Pre-Lubricated Tip.



Protective shield prevents contamination
Bottle sealed for safety

Comfortable, lubricated tip for ease of insertion

One-way safety valve controls flow and prevents reflux

Easy, soft squeeze bottle

Positions for using this enema

Left-side position: Lie on left side with knees bent and arms resting comfortably.



Knee-chest position: Kneel, then lower head and chest forward until left side of face is resting on surface with left arm folded comfortably.



How to use this enema

- Remove the peel-away safety seal and protective shield

Hold bottle upright, grasping bottle cap with fingers. Grasp protective shield with other hand and pull gently to remove.



Administering enema:

- With steady pressure, gently insert enema tip into rectum with a slight side-to-side movement, with tip pointing toward navel. Insertion may be easier if person receiving enema bears down, as if having a bowel movement. Doing so helps relax muscles around anus.
- Do not force enema tip into rectum. Stop using if tip is hard to insert. Forcing the tip into the rectum can cause injury (especially if you have hemorrhoids). If enema tip causes rectal bleeding or pain, get immediate medical care.
- Squeeze bottle until nearly all liquid is gone. It is not necessary to empty bottle completely, as it contains more liquid than need ed.
- Remove tip from rectum and stay in position until urge to evacuate is strong (usually 1 to 5 minutes). If no urge is felt after 5 minutes of using, try to empty bowel. Call a doctor promptly if no liquid comes out of the rectum after 30 minutes because dehydration could occur.

Bottle, cap, and shield are NOT MADE WITH NATURAL RUBBER LATEX



PL313568
V01

FOSTER AND THRIVE READY TO USE SALINE LAXATIVE

sodium phosphate, dibasic, unspecified form and sodium phosphate, monobasic, unspecified form enema

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1089
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM (UNII: GR686LBA74) (PHOSPHATE ION - UNII:NK08V8K8HR, SODIUM CATION - UNII:LYR4MONH37)	SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM	7 g in 118 mL
SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW) (PHOSPHATE ION - UNII:NK08V8K8HR, SODIUM CATION - UNII:LYR4MONH37)	SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM	19 g in 118 mL

Inactive Ingredients

Ingredient Name	Strength
Benzalkonium Chloride (UNII: F5UM2KM3W7)	
EDETATE DISODIUM ANHYDROUS (UNII: 8NLQ36F6MM)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1089-1	1 in 1 CARTON	05/02/2023	
1		133 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:70677-1089-2	2 in 1 CARTON	05/02/2023	
2		133 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	05/02/2023	

Labeler - McKesson (116956644)

Establishment

Name	Address	ID/FEI	Business Operations
Natureplex LLC		062808196	MANUFACTURE(70677-1089)

Revised: 5/2023

McKesson