

TAGAMET- cimetidine tablet
Medtech Products Inc.

Tagamet

Drug Facts

Active ingredient (in each tablet)

Cimetidine 200mg

Purpose

Acid reducer

Uses

- **relieves** heartburn associated with acid indigestion and sour stomach
- **prevents** heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers.

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

Ask a doctor or pharmacist before use if you are taking a prescription drug.

Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- you need to take this product for more than 14 days
- stomach pain continues
- your heartburn continues or worsens

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **right before or any time up to 30 minutes before** eating food or drinking beverages that cause heartburn
 - do not take more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

store at 15-30°C (59-86°F)

Inactive ingredients

cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions?

Call toll-free **1-800-482-4394** weekdays

tagamet.com

PRINCIPAL DISPLAY PANEL

Tagamet® HB 200
Cimetidine Tablets 200 mg
Acid Reducer
70 tablets (70 doses)

TAKE ANYTIME YOU NEED IT:
before meal · during meal · after meal

Tagamet[®] HB 200

Cimetidine Tablets 200 mg
Acid Reducer

JUST ONE TABLET
RELIEVES and PREVENTS
Heartburn and Acid Indigestion

Actual size

70 TABLETS
(70 DOSES)

Tagamet[®] HB 200
Cimetidine Tablets 200 mg
Acid Reducer

Just ONE TABLET
RELIEVES
and PREVENTS
Heartburn
and
Acid Indigestion

Keep cards for future reference.

Tagamet[®] HB 200 Tablets (70 doses) are not for use if blister or foil is open or torn.

Drug Facts	
Active ingredient (in each tablet) Cimetidine 200 mg	Purpose Acid reducer
Uses <ul style="list-style-type: none"> relieves heartburn associated with acid indigestion and sour stomach prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages 	
Warnings Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers. Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. <ul style="list-style-type: none"> with other acid reducers 	
Ask a doctor before use if you have <ul style="list-style-type: none"> frequent chest pain frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain had heartburn over 3 months. This may be a sign of a more serious condition. heartburn with lightheadedness, sweating or dizziness chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness kidney disease liver disease 	
Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.	
Stop use and ask a doctor if <ul style="list-style-type: none"> you need to take this product for more than 14 days stomach pain continues your heartburn continues or worsens 	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) immediately.	
Directions Adults and children 12 years and over: <ul style="list-style-type: none"> to relieve symptoms, swallow 1 tablet with a glass of water to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn do not take more than 2 tablets in 24 hours. Children under 12 years: ask a doctor	
Other information store at 15°-30°C (59°-86°F)	
Inactive ingredients cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polyorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide	
Questions? Call toll-free 1-800-482-4394 weekdays.	

PRINCIPAL DISPLAY PANEL

Tagamet[®] HB 200
Cimetidine Tablets 200 mg
Acid Reducer
30 tablets (30 doses)

TAKE ANYTIME YOU NEED IT:
before meal · during meal · after meal

Tagamet[®] HB 200

Cimetidine Tablets 200 mg
Acid Reducer

JUST ONE TABLET
RELIEVES and PREVENTS
Heartburn and Acid Indigestion

Actual size

30 TABLETS
(30 DOSES)

Tips for Managing Heartburn

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcoholic, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Drug Facts	
Active ingredient (in each tablet) Cimetidine 200 mg	Purpose Acid reducer
Uses <ul style="list-style-type: none"> relieves heartburn associated with acid indigestion and sour stomach prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages 	
Warnings Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers. Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. <ul style="list-style-type: none"> with other acid reducers 	
Ask a doctor before use if you have <ul style="list-style-type: none"> frequent chest pain frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain had heartburn over 3 months. This may be a sign of a more serious condition. heartburn with lightheadedness, sweating or dizziness chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness kidney disease liver disease 	
Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.	
Stop use and ask a doctor if <ul style="list-style-type: none"> you need to take this product for more than 14 days stomach pain continues your heartburn continues or worsens 	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	
Directions Adults and children 12 years and over: <ul style="list-style-type: none"> to relieve symptoms, swallow 1 tablet with a glass of water to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn do not take more than 2 tablets in 24 hours Children under 12 years: ask a doctor	
Other information store at 15°-30°C (59°-86°F)	
Inactive ingredients cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polyorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide	
Questions? Call toll-free 1-800-482-4394 weekdays.	

PRINCIPAL DISPLAY PANEL

Tagamet® HB 200
Cimetidine Tablets 200 mg
Acid Reducer

6 tablets (6 doses)

TAKE ANYTIME YOU NEED IT:
before meal • during meal • after meal

Tagamet
HB 200
Cimetidine Tablets 200 mg
Acid Reducer

JUST ONE TABLET
RELIEVES and PREVENTS
Heartburn and Acid Indigestion

Actual size

6 TABLETS
(6 DOSES)

Read and retain the important information and drug interaction warnings printed on the inside of this carton

Tagamet HB 200
Just ONE TABLET RELIEVES
and PREVENTS Heartburn
and Acid Indigestion

Flamper Evidence: Each tablet individually sealed in a blister pack. Do not use if blister or foil is open or torn.

Drug Facts READ THE DIRECTIONS AND WARNINGS, INCLUDING THOSE PRINTED ON THE INSIDE OF THE CARTON, BEFORE TAKING THIS PRODUCT.

Active ingredient (in each tablet)
Cimetidine 200 mg

Purpose
Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- frequent chest pain
- unexplained weight loss
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- you need to take this product for more than 14 days
- stomach pain continues
- your heartburn continues or worsens

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water
 - to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn
 - do not take more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information store at 15°-30°C (59°-86°F)

Inactive ingredients cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions? Call toll-free 1-800-482-4394 weekdays tagamet.com

TAGAMET

cimetidine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63029-222
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CIMETIDINE (UNII: 80061L1WGD) (CIMETIDINE - UNII:80061L1WGD)	CIMETIDINE	200 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	

POVIDONE (UNII: FZ989GH94E)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	

Product Characteristics

Color	white	Score	no score
Shape	DIAMOND	Size	13mm
Flavor		Imprint Code	TAGAMET;200
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63029-222-02	1 in 1 BOX	06/01/2012	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:63029-222-01	1 in 1 BOX	06/01/2012	
2		30 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:63029-222-03	1 in 1 BOX	06/01/2012	
3		50 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:63029-222-04	4 in 1 BOX	06/01/2019	
4		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:63029-222-05	4 in 1 BOX	06/01/2019	
5		9 in 1 BLISTER PACK; Type 0: Not a Combination Product		
6	NDC:63029-222-70	1 in 1 CARTON	06/01/2012	
6		70 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:63029-222-06	7 in 1 BOX	06/01/2012	
7		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
8	NDC:63029-222-09	1 in 1 CARTON	06/01/2012	
8		90 in 1 BOTTLE; Type 0: Not a Combination Product		
9	NDC:63029-222-11	1 in 1 CARTON	06/01/2012	
9		100 in 1 BOTTLE; Type 0: Not a Combination Product		
10	NDC:63029-	2 in 1 BOX	06/01/2012	

10	222-12	2 III 1 BDA	06/01/2012
10		6 in 1 BLISTER PACK; Type 0: Not a Combination Product	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020238	06/01/2012	

TAGAMET

cimetidine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63029-223
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CIMETIDINE (UNII: 80061L1WGD) (CIMETIDINE - UNII:80061L1WGD)	CIMETIDINE	200 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	

Product Characteristics

Color	blue	Score	2 pieces
Shape	DIAMOND	Size	13mm
Flavor	PEPPERMINT	Imprint Code	TAGAMET;200
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63029-223-30	3 in 1 BOX	02/01/2023	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020238	02/01/2023	

Labeler - Medtech Products Inc. (122715688)

Revised: 6/2024

Medtech Products Inc.