

**ONLY FOR FIRST AID MEDICINE- diphenhydramine hcl, acetaminophen, aspirin, caffeine
O4 Global Trading Usa, Llc**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Only For™ FIRST AID + MEDICINE KIT

ONLY FOR HEADACHE- acetaminophen, aspirin, caffeine tablet

Drug Facts

Active ingredient (in each geltab)

Acetaminophen 250mg

Aspirin 250mg (NSAID*)

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Pain reliever

Pain reliever aid

Uses:

Temporarily relieves minor aches and pains due to:

• headache • a cold • arthritis • muscular aches • toothache • premenstrual & menstrual cramps

Warnings:

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flue-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 8 geltabs in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing

prescription or nonprescription NSAIDs (aspirin, ibuprofen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if • you have liver disease • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma

Ask a doctor or pharmacist before use if you are taking

- a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if • an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • ringing in the ears or loss of hearing occurs • painful area is red or swollen • pain gets worse or lasts for more than 10 days • fever gets worse or lasts for more than 3 days • any new symptoms occur

These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

OVERDOSE WARNING: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

- do not use more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 gels every 6 hours; not more than 8 gels in 24 hours
- children under 12 years: ask a doctor

Other Information:

- store at room temperature between 20-25°C (68 to 77°F) • avoid excessive heat, cold and humidity • close cap tightly after use • read all product information before using.

Inactive Ingredients:

ammonium hydroxide, black iron oxide, brilliant blue FCF, corn starch, D&C yellow No.10 lake, ethyl cellulose, FD&C blue No.1 lake, FD&C red No.40 lake, gelatin, glycerin, hydroxypropyl methyl cellulose, isopropyl alcohol, isopropyl alcohol - butyl alcohol, lactose monohydrate, light liquid paraffin, microcrystalline cellulose, polysorb 85/70/00, povidone K30, pregelatinised starch, propylene glycol, purified water, shellac glaze, stearic acid, talc, titanium dioxide, triacetin

Questions or comments?

1-833-665-9444. (Mon-Fri 9am-5pm)

ONLY FOR ALLERGY- diphenhydramine hcl tablet

Drug Facts

Active ingredient (in each caplet)

Diphenhydramine HCl 25mg

Purpose

Antihistamine

Uses:

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily relieves these symptoms due to the common cold:
 - runny nose
 - sneezing

Warnings:

Do not use

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

- take every 4 to 6 hours, or as directed by doctor
- do not take more than 6 doses in 24 hours

adults and children 12 years and over	• take 1 to 2 caplets
children 6 to under 12 years	• take 1 caplet

Other Information:

- store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F).
- avoid excessive heat and light.
- store in dry place.

Inactive Ingredients:

croscarmellose sodium, dicalcium phosphate, D&C red No.27 aluminium lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, talc, titanium dioxide

Questions or comments?

1-833-665-9444.

(Mon-Fri 9am - 5pm)

ONLY FOR BODY PAIN- acetaminophen, diphenhydramine hcl tablet

Drug Facts**Active ingredient (in each caplet)**

Acetaminophen 500mg

Diphenhydramine HCl 25mg

Purpose

Pain reliever

Nighttime sleep aid

Uses:

- temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- if you have ever had any allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma.

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin
- taking sedatives or tranquilizers.

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery.

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
- pain gets worse or lasts more than 10 days
- fever gets worse or last more than 3 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

OVERDOSE WARNINGS: In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions:

- take with a full glass of water
- Do not take more than directed. **SEE OVERDOSE WARNING.**
- do not take more than 2 caplets in 24 hours

adults and children 12 years of age and over	<ul style="list-style-type: none">• take 2 caplets at bedtime• do not take more than 2 caplets of this product in 24 hours
children under 12 years of age	<ul style="list-style-type: none">• do not use

Other Information:

- Store at room temperature between 20-25°C (68-77°F).
- avoid excessive heat, cold and humidity.
- close cap tightly after use.

Inactive Ingredients:

colloidal silicon dioxide, croscarmellose sodium, FD&C blue No.1, hydroxy propyl methyl cellulose, magnesium stearate, starch, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide.

Questions or comments?

1-833-665-9444. (Mon-Fri 9am-5pm)

ONLY FOR FEVER- acetaminophen tablet**Drug Facts****Active ingredient (in each caplet)**

Acetaminophen 500mg

Purpose

Pain reliever / Fever reducer

Uses:

- temporary relief minor aches and pains due to:
 - the common cold
 - backache
 - toothache
 - premenstrual and menstrual cramps

- temporarily reduces fever.
- headache
- minor pain of arthritis
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever.

Warnings:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had any allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or last more than 3 days
- new symptoms occur.
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. OVERDOSE WARNINGS: In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions:

Do not take more than directed. **SEE OVERDOSE WARNING.**

adults and children 12 years of age and over	<ul style="list-style-type: none"> • take 2 caplets every 6 hours while symptoms last • do not take more than 6 caplets in 24 hours, unless directed by doctor • do not take more than 10 days, unless directed by a doctor
children under 12 years of age	<ul style="list-style-type: none"> • ask a doctor

Other Information:

- Store at room temperature between 20-25°C (68-77°F).
- avoid excessive heat, cold and humidity.
- close cap tightly after use.

Inactive Ingredients:

hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions or comments?

1-833-665-9444. (Mon-Fri 9am-5pm)

Only For™ FIRST AID + MEDICINE KIT

4 SOLUTIONS

HEADACHE 25 Geltabs

ACETAMINOPHEN 250mg / Aspirin (NSAID) 250mg / Caffeine 65mg

ALLERGY 25 Caplets

Diphenhydramine HCl 25mg

BODY PAIN 25 Caplets

ACETAMINOPHEN 500mg / Diphenhydramine HCl 25mg

FEVER 25 Caplets

ACETAMINOPHEN 500mg

FOR DRUG INFORMATION REVIEW, CHECK BOTTLES THAT COME INSIDE THE BOX

Distributed By: O4 GLOBAL TRADING USA, LLC.

12222 Merit Drive, Suite 1900

Dallas, TX 75251-3209

www.onlyfor.com

Packaged By: Strive Pharmaceuticals Inc.

East Brunswick, NJ 08816.

Kit Components Label

25 GELTABS
NDC: 73006-132-95

ADULT Only For

**HEADACHE
PAIN RELIEVER
PAIN RELIEVER AID**

**ACETAMINOPHEN 250mg
Aspirin (NSAID) 250mg / Caffeine 65mg**

Pain reliever
Pain reliever aid

TAMPER EVIDENT Do not use if tamper safety seal under cap is broken or missing

8 60003 57646 5

Drug Facts

Active ingredient (in each geltab) Purpose
 Acetaminophen 250mg Pain reliever
 Aspirin 250mg (NSAID)* Pain reliever
 Caffeine 65 mg Pain reliever aid
 *nonsteroidal anti-inflammatory drug

Uses:
 Temporarily relieves minor aches and pains due to:
 ■ headache ■ cold ■ arthritis ■ muscular aches ■ toothache
 ■ premenstrual & menstrual cramps

Warnings:
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include ■ skin reddening ■ hives ■ rash

Lot No. _____
Exp. Date _____

LIFT HERE

Drug Facts

Active ingredient (in each geltab)	Purpose
Acetaminophen 250mg	Pain reliever
Aspirin 250mg (NSAID)*	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses:
 Temporarily relieves minor aches and pains due to:
 ■ headache ■ cold ■ arthritis ■ muscular aches ■ toothache
 ■ premenstrual & menstrual cramps

Warnings:
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include ■ skin reddening ■ hives ■ rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 geltabs in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally rapid heart beat.

Do not use if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking ■ a prescription drug for diabetes, gout, or arthritis ■ any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms occur

These could be signs of a serious condition

Drug Facts (continued)

■ pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.**

OVERDOSE WARNING: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:
 ■ do not use more than directed
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 2 geltabs every 6 hours; not more than 8 geltabs in 24 hours
 ■ children under 12 years: ask a doctor

Other information:
 ■ store at room temperature between 20-25°C (68 to 77°F) ■ avoid excessive heat, cold and humidity ■ close cap tightly after use ■ read all product information before using.

Inactive ingredients:
 ammonium hydroxide, black iron oxide, brilliant blue FCF, corn starch, D&C yellow No. 10 lake, ethyl cellulose, FD&C blue No. 1 lake, FD&C red No. 40 lake, gelatin, glycerin, hydroxypropyl methyl cellulose, isopropyl alcohol, isopropyl alcohol - butyl alcohol, lactose monohydrate, light liquid paraffin, microcrystalline cellulose, polyorb 85/70/00, povidone K30, pregelatinized starch, propylene glycol, purified water, shellac, silica, stearic acid, talc, titanium dioxide, triacetin

Questions or comments?
 1-833-865-0444, (Mon-Fri 9am - 5pm)

Distributed By: O4 GLOBAL TRADING USA, LLC,
 12222 West Drive, Suite 1900
 Dallas, TX 75254-3209
www.onlyfor.com

Packaged By: Strive Pharmaceuticals Inc.
 East Brunswick, NJ 08816

25 CAPLETS
NDC: 73006-143-95

Only For.

ALLERGY ANTIHISTAMINE NIGHTTIME SLEEP-AID

Diphenhydramine HCl 25mg
Antihistamine

TAKE WITH CAUTION: Do not use if impaired safety and under cap is broken or missing.

8 60003 57649 6

Drug Facts

Active ingredient (in each caplet) Purpose
Diphenhydramine HCl 25mg.....Antihistamine

Uses:

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily relieves these symptoms due to the common cold:
- runny nose
- sneezing

Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Temporarily relieves these symptoms due to the common cold:

- runny nose
- sneezing

Let No. Exp. Date

LIFT HERE

Drug Facts

Active ingredient (in each caplet) Purpose
Diphenhydramine HCl 25mg.....Antihistamine

Uses:

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily relieves these symptoms due to the common cold:
- runny nose
- sneezing

1▼

Drug Facts (continued)

Warnings:

Do not use

- to make a child sleep
- with any other product containing diphenhydramine, even one used on skin.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness.
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

- take every 4 to 6 hours, or as directed by doctor
- do not take more than 6 doses in 24 hours

adults and children 12 years and over	take 1 to 2 caplets
children 6 to under 12 years	take 1 caplet
children under 6 years	do not use

2▼

Drug Facts (continued)

Other Information:

- store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F).
- avoid excessive heat and light.
- store in dry place.

Inactive Ingredients:
croscarmellose sodium, dicalcium phosphate, D&C red No.27 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, talc, titanium dioxide

Questions or comments?
1-833-865-9444.
(Mon-Fri 9am - 5pm)

Distributed By: O4 GLOBAL TRADING USA, LLC.
12222 Merit Drive, Suite 1900
Dallas, TX 75251-3209

www.onlyfor.com

Packaged By: Strive Pharmaceuticals Inc.
East Brunswick, NJ 08816.

3

LB7300614895-REV.00-072020

ADULT

25 CAPLETS
NDC: 73006-120-95

Only For.

BODY PAIN

pl) Purpose
Pain reliever
Nighttime sleep aid

headaches and minor
sleeplessness

acetaminophen.
Take:
1 caplet in 24 hours
then 1 caplet
every 4 to 6 hours
while using this

3▼


**BODY PAIN
PAIN RELIEVER
NIGHTTIME SLEEP AID**

**ACETAMINOPHEN 500mg
Diphenhydramine HCl 25mg**

Pain reliever
Nighttime sleep aid



TAMPER EVIDENT Do not use if imprint is broken or missing



Drug Facts

Active ingredient (in each caplet)
Acetaminophen 500mg.....Pain reliever
Diphenhydramine HCl 25mg.....Nighttime sleep aid

Uses:
■ temporary relief of occasional aches and pains with accompanying sleeplessness

Warnings:
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you:
■ take more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Lot No.
Exp. Date

LIFT HERE

Drug Facts

Active ingredient (in each caplet) **Purpose**
Acetaminophen 500mg.....Pain reliever
Diphenhydramine HCl 25mg.....Nighttime sleep aid

Uses:
■ temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings:
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

1▼

Drug Facts (continued)

Allergy Alert:
Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening
■ blisters
■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ with any other product containing diphenhydramine, even one used on skin
■ in children under 12 years of age
■ if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have
■ liver disease
■ a breathing problem such as emphysema or chronic bronchitis
■ trouble urinating due to an enlarged prostate gland
■ glaucoma.

Ask a doctor or pharmacist before use if
■ you are taking the blood thinning drug warfarin
■ taking sedatives or tranquilizers.

When using this product
■ drowsiness will occur
■ avoid alcoholic drinks
■ do not drive a motor vehicle or operate machinery.

Stop use and ask a doctor if
■ sleeplessness persists continuously for more than 2 weeks.
■ Insomnia maybe a symptom of serious underlying medical illness
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur.
These could be signs of a serious condition.

2▼

Drug Facts (continued)

■ pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.
OVERDOSE WARNING: In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions:
■ take with a full glass of water
■ Do not take more than directed. SEE OVERDOSE WARNING.
■ do not take more than 2 caplets in 24 hours

adults and children 12 years of age and over	■ take 2 caplets at bedtime ■ do not take more than 2 caplets of this product in 24 hours
children under 12 years of age	■ do not use

Other Information:
■ Store at room temperature between 20-25°C (68-77°F).
■ avoid excessive heat, cold and humidity.
■ close cap tightly after use.

Inactive Ingredients:
colloidal silicon dioxide, croscarmellose sodium, FD&C blue No.1, hydroxy propyl methyl cellulose, magnesium stearate, starch, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide.

Questions or comments?
1-833-665-9444, (Mon-Fri 9am - 5pm)

Distributed By: O4 GLOBAL TRADING USA, LLC.
12222 Merit Drive, Suite 1900
Dallas, TX 75251-3209
www.onlyfor.com

Packaged By: Strive Pharmaceuticals Inc.
East Brunswick, NJ 08816.

3

LB7300611995-REV.00-072020



ADULT

Only For

**FEVER
PAIN RELIEVER
FEVER REDUCER**

ACETAMINOPHEN 500mg
Pain reliever / Fever reducer

25 CAPLETS
NDC: 73006-143-95

TAMPER EVIDENT Do not use if imprint is broken or missing



Drug Facts

Active ingredient (in each caplet) **Purpose**
Acetaminophen 500mg.....Pain reliever / Fever reducer

Uses:
temporarily relieves minor aches and pains due to:
■ headache
■ backache
■ toothache
■ minor pain of arthritis
■ muscular aches
premenstrual and menstrual cramps
temporarily reduces fever.

Warnings:
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Lot No.
Exp. Date

LIFT

Drug Facts

Active ingredient (in each caplet) **Purpose**
 Acetaminophen 500mg.....Pain reliever / Fever reducer

Uses:

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
 - temporarily reduces fever.

Warnings:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product



Drug Facts (continued)

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present.
- These could be signs of serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. OVERDOSE WARNING: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.



Drug Facts (continued)

Directions:

Do not take more than directed. **SEE OVERDOSE WARNING.**

adults and children 12 years of age and over	<ul style="list-style-type: none"> ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by doctor ■ do not take more than 10 days, unless directed by a doctor
--	--

children under 12 years of age	<ul style="list-style-type: none"> ■ ask a doctor
--------------------------------	--

Other Information:

- Store at room temperature between 20-25°C (68-77°F).
- avoid excessive heat, cold and humidity.
- close cap tightly after use.

Inactive Ingredients:

hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions or comments?

1-833-665-9444. (Mon-Fri 9am - 5pm)

Distributed By: O4 GLOBAL TRADING USA, LLC,
 12222 Merit Drive, Suite 1900
 Dallas, TX 75251-3209

www.onlyfor.com

Packaged By: Strive Pharmaceuticals Inc.
 East Brunswick, NJ 08816.



LB7300614395-REV.00-072020

KIT LABEL



Only For.

Only For.

Only For.

Only For.



NDC 72006-320-08



ALLERGY

Drug Facts

Active ingredient (in each caplet) Purpose
 Diphenhydramine HCl 25mg.....Antihistamine

Uses:
 ■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ runny nose
 ■ sneezing
 ■ itchy, watery eyes
 ■ itching of the nose or throat
 ■ temporarily relieves these symptoms due to the common cold:
 ■ runny nose
 ■ sneezing

Drug Facts (continued)

Warnings:
Do not use
 ■ to make a child sleepy
 ■ with any other product containing diphenhydramine, even one used on skin.

Ask a doctor before use if you have
 ■ a breathing problem such as emphysema or chronic bronchitis
 ■ glaucoma
 ■ trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product
 ■ marked drowsiness may occur
 ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery
 ■ excitability may occur, especially in children.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:
 ■ take every 4 to 6 hours, or as directed by doctor
 ■ do not take more than 6 doses in 24 hours

adults and children 12 years and over	■ take 1 to 2 caplets
children 6 to under 12 years	■ take 1 caplet
children under 6 years	■ do not use

Drug Facts (continued)

Other Information:
 ■ store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F).
 ■ avoid excessive heat and light.
 ■ store in dry place.

Inactive Ingredients:
 croscarmellose sodium, dicalcium phosphate, D&C red No.27 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, talc, titanium dioxide

Questions or comments?
 1-833-665-9444.
 (Mon-Fri 9am - 5pm)

Distributed By: 04 GLOBAL TRADING USA, LLC.
 12222 Merit Drive, Suite 1900
 Dallas, TX 75251-3209

www.onlyfor.com

Packaged By: Strive Pharmaceuticals Inc.
 East Brunswick, NJ 08816.

FEVER

Drug Facts

Active ingredient (in each caplet) Purpose
 Acetaminophen 500mg.....Pain reliever / Fever reducer

Uses:
 ■ temporarily relieves minor aches and pains due to:
 ■ the common cold ■ headache
 ■ backache ■ minor pain of arthritis
 ■ toothache ■ muscular aches
 ■ premenstrual and menstrual cramps
 ■ temporarily reduces fever.

Warnings:
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product

Drug Facts (continued)

Allergy Alert:
 Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening
 ■ blisters
 ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver disease
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ new symptoms occur
 ■ redness or swelling is present.
 These could be signs of serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. OVERDOSE WARNING: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions:
 Do not take more than directed. SEE OVERDOSE WARNING.

adults and children 12 years of age and over	■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by doctor ■ do not take more than 10 days, unless directed by a doctor
children under 12 years of age	■ ask a doctor

Other Information:
 ■ Store at room temperature between 20-25°C (68-77°F).
 ■ avoid excessive heat, cold and humidity.
 ■ close cap tightly after use.

Inactive Ingredients:
 hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions or comments?
 1-833-665-9444. (Mon-Fri 9am - 5pm)

Distributed By: 04 GLOBAL TRADING USA, LLC.
 12222 Merit Drive, Suite 1900

BODY PAIN

Drug Facts

Active ingredient (in each caplet) Purpose
 Acetaminophen 500mg.....Pain reliever
 Diphenhydramine HCl 25mg.....Nighttime sleep aid

Uses:
 ■ temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings:
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product

Drug Facts (continued)

Allergy Alert:
 Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening
 ■ blisters
 ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ with any other product containing diphenhydramine, even one used on skin
 ■ in children under 12 years of age
 ■ if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have
liver disease
 ■ a breathing problem such as emphysema or chronic bronchitis
 ■ trouble urinating due to an enlarged prostate gland
 ■ glaucoma.

Ask a doctor or pharmacist before use if
 ■ you are taking the blood thinning drug warfarin
 ■ taking sedatives or tranquilizers.

When using this product
 ■ drowsiness will occur
 ■ avoid alcoholic drinks
 ■ do not drive a motor vehicle or operate machinery.

Stop use and ask a doctor if
 ■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days

Drug Facts (continued)

Directions:
 ■ take with a full glass of water
 ■ Do not take more than directed. SEE OVERDOSE WARNING.
 ■ do not take more than 2 caplets in 24 hours

adults and children 12 years of age and over	■ take 2 caplets at bedtime ■ do not take more than 2 caplets of this product in 24 hours
children under 12 years of age	■ do not use

Other Information:
 ■ Store at room temperature between 20-25°C (68-77°F).
 ■ avoid excessive heat, cold and humidity.
 ■ close cap tightly after use.

Inactive Ingredients:
 colloidal silicon dioxide, croscarmellose sodium, FD&C blue No.1, hydroxypropyl methyl cellulose, magnesium stearate, starch, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide.

HEADACHE

Drug Facts

Active ingredient (in each geltab) Purpose
 Acetaminophen 250mg.....Pain reliever
 Aspirin 250mg (NSAID)*.....Pain reliever
 Caffeine 65 mg.....Pain reliever aid
 *nonsteroidal anti-inflammatory drug

Uses:
 Temporarily relieves minor aches and pains due to:
 ■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache
 ■ premenstrual & menstrual cramps

Warnings:
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 geltabs in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally rapid heart beat.

Do not use if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking
 ■ a prescription drug for diabetes, gout, or arthritis
 ■ any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools
 ■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms occur
 These could be signs of a serious condition

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
OVERDOSE WARNING: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:
 ■ do not use more than directed
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 2 geltabs every 6 hours; not more than 8 geltabs in 24 hours
 ■ children under 12 years: ask a doctor

Other Information:
 ■ store at room temperature between 20-25°C (68 to 77°F) ■ avoid excessive heat, cold and humidity ■ close cap tightly after use ■ read all product information before using.

Inactive Ingredients:
 ammonium hydroxide, black iron oxide, brilliant blue FCF, corn starch, D&C yellow No.10 lake, ethyl cellulose, FD&C blue No.1 lake, FD&C red No.40 lake, nebutin, aluminum hydroxide/methylcellulose, isomalt

Dallas, TX 75251-3209
www.onlyfor.com
Packaged By: Strive Pharmaceuticals Inc.
East Brunswick, NJ 08816. 3

Questions or comments?
1-833-665-9444. (Mon-Fri 9am - 5pm)
Distributed By: O4 GLOBAL TRADING USA, LLC.
12222 Merit Drive, Suite 1900
Dallas, TX 75251-3209
www.onlyfor.com
Packaged By: Strive Pharmaceuticals Inc.
East Brunswick, NJ 08816. 3

alcohol, isopropyl alcohol - butyl alcohol, lactose monohydrate, light liquid paraffin, microcrystalline cellulose, polysorb 85/70/00, povidone K30, pregelatinised starch, propylene glycol, purified water, shellac glaze, stearic acid, talc, titanium dioxide, triacetin
Questions or comments?
1-833-665-9444. (Mon-Fri 9am - 5pm)
Distributed By: O4 GLOBAL TRADING USA, LLC.
12222 Merit Drive, Suite 1900
Dallas, TX 75251-3209
www.onlyfor.com
Packaged By: Strive Pharmaceuticals Inc.
East Brunswick, NJ 08816. 3

ONLY FOR FIRST AID MEDICINE

diphenhydramine hcl, acetaminophen, aspirin, caffeine kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73006-350
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73006-350-98	1 in 1 BOX; Type 0: Not a Combination Product	11/15/2020	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE, PLASTIC	25
Part 2	1 BOTTLE, PLASTIC	25
Part 3	1 BOTTLE, PLASTIC	25
Part 4	1 BOTTLE, PLASTIC	25

Part 1 of 4

ONLY FOR HEADACHE

acetaminophen, aspirin, caffeine tablet

Product Information

Item Code (Source)	NDC:73006-132
---------------------------	---------------

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients	
Ingredient Name	Strength
AMMONIA (UNII: 5138Q19F1X)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
BUTYL ALCOHOL (UNII: 8PJ61P6TS3)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SHELLAC (UNII: 46N107B71O)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics			
Color	green (Dark Green - White)	Score	no score
Shape	ROUND	Size	13mm
Flavor		Imprint Code	S77
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73006-132-95	1 in 1 CARTON		
1		25 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	11/15/2020	

Part 2 of 4

ONLY FOR ALLERGY

diphenhydramine hcl tablet

Product Information

Item Code (Source) NDC:73006-148

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
Shape	OVAL (Caplet)	Size	11mm
Flavor		Imprint Code	DP
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73006-148-95	1 in 1 CARTON		
1		25 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part338	11/15/2020	

Part 3 of 4

ONLY FOR BODY PAIN

acetaminophen, diphenhydramine hcl tablet

Product Information

Item Code (Source)	NDC:73006-120
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY3SJ)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue (LIGHT BLUE)	Score	no score
Shape	capsule	Size	18mm
Flavor		Imprint Code	AD01
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73006-120-	1 in 1 CARTON		

95	1 in 1 CARTON		
1	25 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	11/15/2020	

Part 4 of 4

ONLY FOR FEVER

acetaminophen tablet

Product Information

Item Code (Source)	NDC:73006-143
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	capsule	Size	17mm
Flavor		Imprint Code	S500
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73006-143-95	1 in 1 CARTON		
1		25 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	11/15/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part338	11/15/2020	

Labeler - O4 Global Trading Usa, Llc (081226861)

Revised: 12/2020

O4 Global Trading Usa, Llc