

MEDIQUE AT HOME DECOREL FORTE- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride tablet, film coated
Unifirst First Aid Corporation

Medique @ Home Decorel Forte

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 100 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Uses

Temporarily relieves these symptoms due to the common cold:

- headache
- nasal congestion
- cough
- minor aches and pains
- sore throat
- sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make cough more productive

Temporarily reduces fever.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is

accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.

If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough that lasts as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

Adults and children: (12 years and over):

- take 2 caplets every 4 hours
- not to take more than 10 caplets in 24 hours

Children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- **TAMPER EVIDENT** child resistant packets
- do not use any opened or torn packets

Inactive ingredients

corn starch, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments? 1-800-634-7680

Medique® @ Home

Decorel Forte

Acetaminophen 325 mg • **Pain Reliever/Fever Reducer**

Dextromethorphan HBr 10 mg • **Cough Suppressant**

Guaifenesin 100 mg • **Expectorant**

Phenylephrine HCl 5 mg • **Nasal Decongestant**

Pull to Open

Child-Resistant Packets

*not actual size

50 Packets

(2 Caplets Per Packet)

| | | | | |
|--|--|--|----------------------|--------------------|
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | | | | |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) | | | | |
| TALC (UNII: 7SEV7J4R1U) | | | | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | | | | |
| MAGNESIUM STEARATE (UNII: 70097M6I3O) | | | | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | | | | |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) | | | | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | | | | |
| | | | | |
| Product Characteristics | | | | |
| Color | orange | Score | no score | |
| Shape | CAPSULE (Caplet) | Size | 17mm | |
| Flavor | | Imprint Code | 44;546 | |
| Contains | | | | |
| | | | | |
| Packaging | | | | |
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:47682-735-33 | 50 in 1 BOX | 09/26/2022 | |
| 1 | NDC:47682-735-99 | 2 in 1 PACKET; Type 0: Not a Combination Product | | |
| 2 | NDC:47682-735-99 | 2 in 1 PACKET; Type 0: Not a Combination Product | 09/26/2022 | |
| | | | | |
| | | | | |
| Marketing Information | | | | |
| Marketing Category | Application Number or Monograph Citation | | Marketing Start Date | Marketing End Date |
| OTC Monograph Drug | M012 | | 09/26/2022 | |

Labeler - Unifirst First Aid Corporation (832947092)