

ACETAMINOPHEN- acetaminophen tablet
TARGET CORPORATION

Extra strength
Acetaminophen gelcaps, 500mg
Rapid release
Pain reliever/fever reducer
For Adults

Active ingredient (in each gelcap)

Acetaminophen USP, 500mg

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Keep out of reach of children.

Overdose Warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over

- take 2 gelscaps every 6 hours while symptoms last
- do not take more than 6 gelscaps in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

children under 12 years

- ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F). See USP Controlled Room Temperature
- avoid high humidity
- see end panel for expiration date and lot number

Inactive Ingredients

ammonium hydroxide, black iron oxide, colloidal silicon dioxide, croscarmellose sodium, D&C red #33, FD&C blue #1, FD&C red #40, FD&C yellow #6, gelatin, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, hypromellose, iron oxide red, isopropyl alcohol, n-butyl alcohol, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, stearic acid, titanium dioxide, yellow iron oxide.

Questions or Comments?

call **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST.

Principal Display Panel

size	:1+3/4 X 1+3/4 X 3+3/8	via
ref #	:PPI180604B	da:
material	:.016 SBS	

COATING
FREE AREA



Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

- | | |
|----------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>adults and children 12 years and over</p> | <ul style="list-style-type: none"> ■ take 2 gelscaps every 6 hours while symptoms last ■ do not take more than 6 gelscaps in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor |
| <p>children under 12 years</p> | <ul style="list-style-type: none"> ■ ask a doctor |

Other information

- store at 20° - 25°C (68° - 77°F). See USP Controlled Room Temperature
- avoid high humidity
- see end panel for expiration date and lot number

COATING
FREE AREA



Drug Facts (continued)

Inactive ingredients: ammonium hydroxide, black iron oxide, calcium silicate, croscarmellose sodium, D&C red #33, FD&C blue #1, FD&C red #40, FD&C yellow #6, gelatin, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, hypromellose, iron oxide red, isopropyl alcohol, n-butyl alcohol, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, stannic oxide, stearic acid, titanium dioxide, yellow iron oxide.

Questions or comments?

call 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST.

Satisfaction guaranteed - Love it or your money back.

NDC 11673-187-05

Distributed by Target Corporation
Minneapolis, MN 55403

Made in India
TM & ©2024 Target Brands, Inc.

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Extra Strength Tylenol® Rapid Release Gels.

ASPIRIN FREE



**Extra Strength
Acetaminophen**
500 mg Gelscaps, Rapid Release
Pain Reliever / Fever Reducer



**500 mg Gelscaps, Rapid Release
Pain Reliever / Fever Reducer**
For Adults
50 GELCAPS



**Extra Strength
Acetaminophen**
Compare to active ingredient in Extra Strength Tylenol® Rapid Release Gels*



Lot
Exp.

COATING
FREE AREA

READ AND KEEP CARTON FOR COMPLETE
WARNINGS AND INFORMATION

TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient	Purpose
(in each gelscap)	
Acetaminophen USP 500 mg	Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - backache
 - minor pain of arthritis
 - the common cold
 - toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.

Compare to active ingredient
in Extra Strength Tylenol®
Rapid Release Gels*

Extra Strength Acetaminophen

500 mg Gelcaps, Rapid Release
Pain Reliever / Fever Reducer

For Adults



Actual Size
100 Gelcaps

100 GELCAPS

Important: Read all product information before using.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**

Drug Facts

Active ingredient (in each gelcap) Acetaminophen USP, 500 mg. Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - the common cold
 - premenstrual and menstrual cramps
 - temporarily reduces fever
- backache
- toothache

Warnings

- Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
 - more than 4,000 mg of acetaminophen in 24 hours
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks every day while using this product
- Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
 - skin redness
 - hives
 - skin rash
 - itching
 - swelling of the face, lips, tongue, or throat
 - difficulty breathing

Directions

- do not take more than directed (see overdose warning)
- adults and children 12 years and over
 - take 2 gelcaps every 6 hours while symptoms last
 - do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
 - do not use for more than 10 days unless directed by a doctor
- children under 12 years
 - ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F). See USP Controlled Room Temperature
- avoid high humidity
- see end panel for expiration date and lot number

Inactive ingredients

ammonium hydroxide, black iron oxide, calcium stearate, croscarmellose sodium, D&G red #33, FD&C blue #1, FD&C red #40, FD&C yellow #6, gelatin, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, hypromellose, iron oxide red, isopropyl alcohol, n-butyl alcohol, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, propylene glycol, stearic acid, titanium dioxide, yellow iron oxide

Questions or comments?

call 1-877-776-5183 Mon-Fri 8:00 AM EST to 5:00 PM PST.

(CONTINUED ON BACK OF LABEL)

Dist. by Target Corporation

Minneapolis, MN 55403

Made in India

TM & ©2024

Target Brands, Inc.

All trademarks are property of their respective owners.

This product is not affiliated with the makers/owners of Extra Strength Tylenol® Rapid Release Gels.

NDC 11673-167-10

094-01 0547 R00 C-002262-01-034-0000

8 13874 02065 0

202000004938
702000002321

LOT EXP

ASPIRIN FREE

Inside (adhesive side)

Drug Facts (continued)

Ask a doctor before use if you have liver disease, bleeding drug warfarin.

Stop use and ask a doctor if

your pain gets worse or lasts more than 10 days

your fever gets worse or lasts more than 3 days

new symptoms occur

redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

BUTYL ALCOHOL (UNII: 8PJ61P6TS3)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
HYPROMELLOSE 2910 (3 MPA.S) (UNII: 0VUT3PMY82)
FERROSO FERRIC OXIDE (UNII: XM0M87F357)
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)
POVIDONE K30 (UNII: U725QWY32X)
STARCH, CORN (UNII: O8232NY3SJ)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
HYDROXYMETHYL CELLULOSE (UNII: 273FM27VK1)
STEARIC ACID (UNII: 4ELV7Z65AP)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
AMMONIA (UNII: 5138Q19F1X)
SHELLAC (UNII: 46N107B71O)
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)
D&C RED NO. 33 (UNII: 9DBA0SBB0L)
FD&C RED NO. 40 (UNII: WZB9127XOA)
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)
GELATIN (UNII: 2G86QN327L)

Product Characteristics

Color	gray (Encapsulated with red opaque and blue gray opaque hard gelatin shells)	Score	2 pieces
Shape	OVAL	Size	19mm
Flavor		Imprint Code	G1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-167-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	02/27/2023	
2	NDC:11673-167-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/27/2023	
3	NDC:11673-167-21	225 in 1 BOTTLE; Type 0: Not a Combination Product	02/27/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/27/2023	

Labeler - TARGET CORPORATION (006961700)