

**ISOXSUPRINE HYDROCHLORIDE- isoxsuprine hydrochloride tablet**  
**Vista Pharmaceuticals, Inc.**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*


-----  
**ISOXSUPRINEHYDROCHLORIDE TABLETS, USP**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA.*

CAUTION: Federal Law prohibits dispensing without prescription

**DESCRIPTION**

Isoxsuprine HCl occurs as a white odorless, crystalline powder, having a bitter taste, It has a following structural formula



**INDICATIONS**

Based on a review of this drug by the National Academy of Sciences -National Research Council and / or other information, the FDA has classified the medications as follows :

Possibly Effective :

1. For the relief of symptoms associated with cerebral vascular insufficiency
2. In Peripheral vascular disease of arteriosclerosis obliterans, thromboangitis obliterans (Buerger's Disease) and Raynaud's disease.

Final classification of the less-than-effective indications requires further investigation.

**COMPOSITION**

Each tablet contains Isoxsuprine HCl 20 mg.

These tablets contain the following inactive ingredients: dibasic calcium phosphate (anhydrous), lactose, magnesium stearate. microcrystalline cellulose, povidone k30, and sodium starch glycolate.

**DOSAGEANDADMINISTRATION**

Oral:10 to 20 mg three or four times daily

**CONTRAINdicATIONS AND CAUTIONS**

**Oral**

There are no known contraindications to oral use when administered in recommended doses. Should not be given immediately postpartum or in the presence of arterial bleeding.

## ADVERSE REACTIONS

On rare occasion, oral administration of the drug has been associated in time with the occurrences of hypotension, tachycardia, chest pain, nausea, vomiting, dizziness, abdominal distress, and severe rash. If rash appears, the drug should be discontinued.

Although available evidence suggests a temporal association of these reactions with isoxsuprine, a casual relationship can be neither confirmed nor refuted.

$\beta$ -Adrenergic receptor stimulants such as isoxsuprine hydrochloride have been used to inhibit pre-term labor. Maternal and fetal tachycardia may occur under such use. Hypocalcemia, hypoglycemia, hypotension and ileus have been reported to occur in infants whose mothers received isoxsuprine. Pulmonary edema has been reported in mothers treated with  $\beta$ -stimulants. Isoxsuprine HCl tablets, USP is neither approved nor recommended for use in the treatment of premature labor.

## HOW SUPPLIED

Isoxsuprine HCl tablets, USP are supplied in HDPE bottles.

20 mg Bottles of 1,000's: NDC61971-065-10

Manufactured in India by  
Vista Pharmaceuticals, Limited.

For  
Vista Pharmaceuticals, Inc.  
Revised:07/2017

**VISTA**  
NDC 61971-065-10  
**Isoxsuprine Hydrochloride Tablets, USP**  
**20 mg**  
**Rx Only**  
**1000 Tablets**

Each Tablet Contains:  
Isoxsuprine HCl, USP ..... 20 mg  
**WARNING: AS WITH ALL MEDICATIONS, KEEP OUT OF REACH OF CHILDREN.**  
Store at controlled room temperature 15° - 30°C (59° - 86°F) [see USP].  
**Usual Dosage:** See accompanying literature.  
**Pharmacist:** Dispense in a tight, light resistant container as defined in the USP.

Manufactured by:  
Vista Pharmaceuticals, Ltd.  
Gopalaipalli - 508254, India  
For:  
Vista Pharmaceuticals, Inc.  
West Orange, NJ 07052

Lot No.:  
Exp. Date:

304011-02  
N 3 61971-065-10 9

## ISOXSUPRINE HYDROCHLORIDE

isoxsuprine hydrochloride tablet

### Product Information

Product Type

HUMAN PRESCRIPTION DRUG

Item Code (Source)

NDC:61971-065

<b>Route of Administration</b>	ORAL			
<b>Active Ingredient/Active Moiety</b>				
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
	Isoxsuprine hydrochloride (UNII: V74TEQ36CO) (Isoxsuprine - UNII:R15UIB245N)	Isoxsuprine hydrochloride	20 mg	
<b>Inactive Ingredients</b>				
	<b>Ingredient Name</b>	<b>Strength</b>		
	LACTOSE (UNII: J2B2A4N98G)			
	CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
	CALCIUM PHOSPHATE, DIBASIC, ANHYDROUS (UNII: L11K75P92J)			
	POVIDONE K30 (UNII: U725QWY32X)			
	SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)			
	MAGNESIUM STEARATE (UNII: 70097M6I30)			
<b>Product Characteristics</b>				
<b>Color</b>	white	<b>Score</b>	2 pieces	
<b>Shape</b>	ROUND	<b>Size</b>	10mm	
<b>Flavor</b>		<b>Imprint Code</b>	20;VISTA065	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:61971-065-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	09/19/1997	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
unapproved drug other		09/19/1997		

**Labeler** - Vista Pharmaceuticals, Inc. (943932806)

### Establishment

Name	Address	ID/FEI	Business Operations
Vista Pharmaceuticals, Limited.		916648541	manufacture(61971-065) , analysis(61971-065)