

NOTTS - MUSCLE ACHES AND PAIN- acetaminophen tablet
VIVUNT PHARMA LLC

NOTTS - Muscle Aches & Pain

Drug Facts

<i>Active ingredient (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 650 mg	Pain Reliever/Fever Reducer

Uses

For temporary relief of minor aches and pains due to:

- common cold
- headache
- toothache
- muscular aches
- backaches
- premenstrual and menstrual cramps
- minor pain from arthritis

Temporarily reduces fever

Warnings

Liver Warning

This product contains acetaminophen. Severe damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours, which is the maximum daily amount.
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- rash
- blisters

If a skin reaction occurs, stop use and seek medical attention immediately.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients.

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if you are taking

the blood thinning drug warfarin

Stop use and ask a doctor if

- pain lasts more than 10 days
- pain lasts more than 3 days
- pain or fever gets worse
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use

Keep out of reach of children.

Overdose warning

Do not exceed recommended dosage. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- Do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 1 caplets every 4 to 6 hours▪ do not take more than 6 caplets in 24 hours, or as directed by a doctor▪ do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- Store between 20-25°C (68-77°F)
- Tamper-evident: Do not use if carton is open or if the foil inner seal is broken or missing.

Inactive ingredients

Corn Starch, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol, Polyvinyl Alcohol, Povidone, Stearic Acid, Sodium Starch Glycolate, Soy Lecithin, Talc, Titanium Dioxide.

* Divided into 2 intakes of one caplet every 4 hours. Use only as directed in Drug Facts.

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark Tylenol® 8HR.

Product of China

Distributed by:

VIVUNT PHARMA LLC

8950 SW 74th Court, Suite 1901

Miami, Florida Z.C. 33156-3175

PRINCIPAL DISPLAY PANEL - 24 Caplets

NOTTS™ 8HR*

Muscle Aches & Pain

Acetaminophen

- Minor Muscle Aches
- Fever Reducer

Compare to Tylenol® 8HR active ingredients**

NDC 82706-008-01

650 mg each caplet

24 CAPLETS



PRINCIPAL DISPLAY PANEL - 100 Caplets

NOTTS™ 8HR*

Muscle Aches & Pain

Acetaminophen

Drug Facts (continued)
 doctor or pharmacist ■ If you are allergic to acetaminophen or any of the inactive ingredients.
Ask a doctor before use if you have liver disease.
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
Stop use and ask a doctor if
 ■ pain lasts more than 10 days
 ■ fever lasts more than 3 days
 ■ pain or fever gets worse
 ■ new symptoms occur
 ■ redness or swelling is present.
 These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: Do not exceed recommended dosage. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
 ■ Do not take more than directed (see overdose warning).

Drug Facts
Active ingredient (in each caplet) Purpose
 Acetaminophen 650 mg Pain Reliever/Fever Reducer
Uses
 For temporary relief of minor aches and pains due to:
 ■ common cold ■ headache ■ toothache
 ■ muscular aches ■ backaches ■ premenstrual and menstrual cramps ■ minor pain from arthritis.
 Temporarily reduces fever.
Warnings
Liver warning: This product contains acetaminophen. Severe damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours, which is the maximum daily amount
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ rash ■ blisters. If a skin reaction occurs, stop use and seek medical attention immediately.
Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a

*Divided into 2 flakes of one caplet every 4 hours. Use only as directed in Drug Facts.

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark Tylenol® 8HR.

Drug Facts (continued)
 adults and children 12 years and over
 ■ take 1 caplet every 4 to 6 hours
 ■ do not take more than 6 caplets in 24 hours, or as directed by a doctor
 ■ do not use for more than 10 days unless directed by a doctor
 children under 12 years
 ■ ask a doctor
Other information
 ■ Store between 20°-25°C (68°-77°F)
 ■ Tamper-evident: Do not use if carton is open or if the foil inner seal is broken or missing.
Inactive ingredients
 Corn Starch, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol, Polyvinyl Alcohol, Povidone, Stearic Acid, Sodium Starch Glycolate, Soy Lecithin, Talc, Titanium Dioxide.

Product of China.
 Distributed by:
 VIVUNT PHARM LLC
 8950 SW 74th Court, Suite 1901
 Miami, Florida Z.C. 33156-3178



- Minor Muscle Aches
- Fever Reducer

Compare to Tylenol® 8HR active ingredients**

NDC 82706-008-02

650 mg each caplet

100 CAPLETS



Drug Facts (continued)

- new symptoms occur
- redness or swelling is present.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Over-the-counter drugs do not need recommended dosage or usage directions. Get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- Do not take more than directed (see overdose warning).
- Take 1 caplet every 4 to 6 hours
- do not take more than 6 caplets in 24 hours, or as directed by a doctor
- Do not take more than 10 days unless directed by a doctor
- ask a doctor

adults and children 12 years and over

children under 12 years

Other information

- Store between 20°-25°C (68°-77°F)
- Tamper-evident. Do not use if carton is open or if the foil inner seal is broken or missing.

Inactive ingredients

Com Starch, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol, Polyvinyl Alcohol, Povidone, Stearic Acid, Sodium Starch Glycolate, Soy Lecithin, Talc, Titanium Dioxide.

Drug Facts

Active ingredient (in each caplet) Acetaminophen 650mg

Purpose Pain Reliever/Fever Reducer

Uses

For temporary relief of minor aches and pains due to:

- fever
- headache
- toothache
- aches
- backaches
- menstrual and menstrual cramps
- minor pain from arthritis.

 Temporarily reduces fever.

Warnings

Liver warning: This product contains acetaminophen. Severe damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- rash
- blisters.

 If a skin reaction occurs, stop use and seek medical attention immediately.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- If you are allergic to acetaminophen or any of the inactive ingredients.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain lasts more than 10 days
- fever lasts more than 3 days
- pain or fever gets worse

Notts

8HR

Muscle Aches & Pain

Acetaminophen • Minor Muscle Pains • Fever Reducer

650 mg each caplet

Actual size

Compare to Tylenol® 8HR active ingredient**

NDC 82706-008-02

100 CAPLETS



* Divided into 2 intakes of one caplet every 4 hours. Use only as directed in Drug Facts.

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark Tylenol® 8HR.

Product of China.

Distributed by:
PIKARI, LLC
 8950 SW 74th Court, Suite 1901
 Miami, Florida Z.C. 33156-3178

NOTTS - MUSCLE ACHES AND PAIN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82706-008
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	Notts;650
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82706-008-02	1 in 1 CARTON	08/17/2022	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:82706-008-01	1 in 1 CARTON	08/17/2022	
2		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/17/2022	

Labeler - VIVUNT PHARMA LLC (045829437)

Revised: 10/2023

VIVUNT PHARMA LLC