

**NOTTS - EXTRA STRENGTH 500 MG- acetaminophen tablet**  
**VIVUNT PHARMA LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**NOTTS - Extra Strength 500 mg**

**Drug Facts**

---

<i>Active ingredient (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 500 mg	Pain Reliever/Fever Reducer

---

**Uses**

Temporary relief of minor aches and pains due to:

- common cold
- headache
- toothache
- muscular aches
- backaches
- premenstrual and menstrual cramps
- minor pain from arthritis

Temporarily reduces fever

**Warnings**

**Liver Warning**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours, which is the maximum daily amount.
- with other drugs containing acetaminophen.
- 3 or more alcoholic drinks every day while using this product.

**Allergy Alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- rash
- blisters

If a skin reaction occurs, stop use and seek medical attention immediately.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients.

**Ask a doctor before use if you have**

liver disease

**Ask a doctor or pharmacist before use if you are taking**

the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain lasts more than 10 days
- fever lasts more than 3 days
- pain or fever gets worse
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

Keep out of reach of children.

**Overdose warning**

Do not take more than recommended dosage. In case of overdose, get medical help or contact a Poison Control Center. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **Do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"><li>▪ take 2 caplets every 6 hours</li><li>▪ do not take more than 8 caplets in 24 hours, or as directed by a doctor</li><li>▪ do not use for more than 10 days unless directed by a doctor</li></ul>
children under 12 years	ask a doctor

**Other information**

- Store between 20-25°C (68-77°F)
- Tamper-evident: Do not use if carton is open or if the foil inner seal is broken or

missing.

- Tamper-evident: Do not use if the bottle cap is open or if the foil inner seal is broken or missing.

### **Inactive ingredients**

Corn Starch, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol, Polyvinyl Alcohol, Povidone, Stearic Acid, Sodium Starch Glycolate, Soy Lecithin, Talc, Titanium Dioxide.

\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark Tylenol® Extra Strength.

Product of China

Distributed by:

VIVUNT PHARMA LLC

8950 SW 74th Court, Suite 1901

Miami, Florida Z.C. 33156-3175

### **PRINCIPAL DISPLAY PANEL - 24 Caplets**

NOTTS™

Extra Strength

Acetaminophen

- Pain Reliever
- Fever Reducer

Compare to Tylenol® Extra Strength active ingredients\*

NDC 82706-007-01

500 mg each caplet

24 CAPLETS

Extra Strength.



**PRINCIPAL DISPLAY PANEL - 100 Caplets**

**NOTTS™**

Extra Strength

Acetaminophen

**Drug Facts (continued)**

- do not take more than 8 caplets in 24 hours, or as directed by a doctor
- do not use for more than 10 days unless directed by a doctor
- ask a doctor

children under 12 years

**Other information**

- Store between 20°-25°C (68°-77°F)
- Tamper-evident: Do not use if carton is open or if the foil inner seal is broken or missing.

**Inactive ingredients**

Com Starch, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol, Polyvinyl Alcohol, Povidone, Stearic Acid, Sodium Starch Glycolate, Soy Lecithin, Talc, Titanium Dioxide.

**Drug Facts**

**Active ingredient (in each caplet)** Purpose  
Acetaminophen 500 mg • Pain Reliever/Fever Reducer

**Uses**

Temporary relief of minor aches and pains due to:  
 ■ common cold ■ headache ■ toothache  
 ■ muscular aches ■ backaches ■ premenstrual and menstrual cramps ■ minor pain from arthritis.  
 Temporarily reduces fever.

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ rash ■ blisters. If a skin reaction occurs, stop use and seek medical attention immediately.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to

**Drug Facts (continued)**

acetaminophen or any of the inactive ingredients. Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

- Stop use and ask a doctor if**
- pain lasts more than 10 days
  - fever lasts more than 3 days
  - pain or fever gets worse
  - new symptoms occur
  - redness or swelling is present
- These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Do not take more than recommended dosage. In case of overdose, get medical help or contact a Poison Control Center. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- Do not take more than directed (see overdose warning).
- adults and children ■ take 2 caplets every 12 hours and over ■ 6 hours

\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark Tylenol® Extra Strength.

Product of China.  
 Distributed by:  
**VIVUNT PHARM LLC**  
 8950 SW 74th Court, Suite 1901  
 Miami, Florida Z.C. 33156-3178



**Extra Strength**

**Acetaminophen**

500 mg each caplet

- Pain Reliever
- Fever Reducer



Compare to Tylenol® Extra Strength active ingredient\*  
 NDC 82706-007-01

**24** CAPLETS



- Pain Reliever
- Fever Reducer

Compare to Tylenol® Extra Strength active ingredients\*

NDC 82706-007-02

500 mg each caplet

100 CAPLETS

Extra Strength.



## PRINCIPAL DISPLAY PANEL - 400 Caplets

NOTTS™

Extra Strength

Acetaminophen

- Pain Reliever
- Fever Reducer

NDC 82706-007-03  
 500 mg each caplet  
 400 CAPLETS  
 Extra Strength.

**Notts**  
**Extra Strength**  
**Acetaminophen**  
 500 mg each caplet  
 • Pain Reliever  
 • Fever Reducer  
 Compare to Tylenol®  
 Extra Strength active ingredient\*  
 NDC 82706-007-03  
**400** CAPLETS

**TO OPEN: PUSH AND TURN CAP**

**Drug Facts**  
**Active Ingredient (in each caplet)**  
 Acetaminophen 500 mg

**Purpose**  
 Pain Reliever/Fever Reducer

**Uses**  
 Temporary relief of minor aches and pains due to: ■ common cold ■ headache ■ toothache  
 ■ muscular aches ■ backaches ■ premenstrual and menstrual cramps ■ minor pain from arthritis.  
 Temporarily reduces fever.

**Warnings**  
 Do not take this product with acetaminophen. Severe damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ rash ■ blisters. If a skin reaction occurs, stop use and seek medical attention immediately.  
**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are allergic to acetaminophen or any of the inactive ingredients.  
**Ask a doctor before use if you have liver disease.**  
**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**  
**Stop use and ask a doctor if** ■ pain lasts more than 10 days ■ fever lasts more than 3 days ■ pain or fever gets worse ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.  
**If pregnant or breast feeding,** ask a health professional before use.  
**Keep out of reach of children.**  
**Overdose warning:** Do not take more than recommended dosage. In case of overdose, get medical help or contact a Poison Control Center. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
 ■ Do not take more than directed (see overdose warning).  
 ■ Adults and children 12 years and over: ■ take 2 caplets every 6 hours ■ do not take more than 8 caplets in 24 hours, or as directed by a doctor ■ do not use for more than 10 days unless directed by a doctor.  
 ■ Children under 12 years: ask a doctor.

**Other information**  
 ■ Store between 20°-25° C (68°-77° F).  
 ■ Tamper-evident: Do not use if the bottle cap is open or if the foil inner seal is broken or missing.

**Inactive ingredients**  
 Croscarmellose Sodium, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol, Polyvinyl Alcohol, Potassium Stearate, Sodium Stearate, Soy Lecithin, Talc, Titanium Dioxide.

\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark Tylenol® Extra Strength.

Product of China.  
 Distributed by:  
 VIVUNT PHARM LLC  
 8950 SW 74th Court, Suite 1901  
 Miami, Florida Z.C. 33156-3178

**PRINCIPAL DISPLAY PANEL - 6 Caplets**

NOTTS™  
 Extra Strength  
 Acetaminophen  
 Pain Reliever  
 Fever Reducer  
 NDC 82706-007-03  
 500 mg each caplet  
 400 CAPLETS  
 Extra Strength

• Relieves Pain  
 • Reduces Fever  
 Compare to Tylenol® Extra Strength active ingredient\*  
 NDC 82706-007-04

**3 PORTABLE POUCHES**  
 (2 Softgels per pouch)



# Extra Strength





# Extra Strength

## Acetaminophen

500 mg each caplet



**6** CAPLETS



**Drug Facts** (continued)

**Inactive ingredients**  
 Corn Starch, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol, Polyvinyl Alcohol, Povidone, Sodium Starch Glycolate, Soy Lecithin, Stearic Acid, Talc, Titanium Dioxide.

**Ask a doctor before use** If you have liver disease

**Ask a doctor or pharmacist before use** if you are taking the blood thinning drug warfarin.

**Stop use and ask a doctor if**  
 ■ pain lasts more than 10 days ■ fever lasts more than 3 days ■ pain or fever gets worse  
 ■ new symptoms occur ■ redness or swelling is present  
 These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Do not take more than recommended dosage. In case of overdose, get medical help or contact a Poison Control Center. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ Do not take more than directed (see overdose warning).  
 adults and children 12 years and over  
 ■ take 2 caplets every 6 hours ■ do not take more than 8 caplets in 24 hours, or as directed by a doctor ■ do not use for more than 10 days unless directed by a doctor  
 children under 12 years  
 ■ ask a doctor

**Other information**  
 ■ Store between 20°-25°C (68°-77°F)  
 ■ Tamper evident: Do not use if carton or pouch is open.


\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark Tylenol® Extra Strength.

ence

Barcode

**VIVUNT**  
 Product of Inc  
 Distributed by  
**VIVUNT PHAR**  
 8950 SW 74th  
 Miami, Florida





## NOTTS - EXTRA STRENGTH 500 MG

acetaminophen tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:82706-007
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TALC (UNII: 7SEV7J4R1U)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	Notts;500
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82706-007-01	1 in 1 CARTON	08/17/2022	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:82706-007-02	1 in 1 CARTON	08/17/2022	
2		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:82706-007-03	400 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/17/2022	
4	NDC:82706-007-04	3 in 1 CARTON	09/20/2023	
4		2 in 1 POUCH; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	08/17/2022	

**Labeler** - VIVUNT PHARMA LLC (045829437)

Revised: 9/2023

VIVUNT PHARMA LLC