

**BISOLVINE ADULT- guaifenesin, dextromethorphan liquid  
MENPER DISTRIBUTORS INC.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Active Ingredients (in each 10 mL) Purpose**

Dextromethorphan 20 mg..... Cough Suppressant  
Guaifenesin 200 mg ..... Expectorant

**Purpose**

Cough Suppressant  
Expectorant

**Uses**

temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold and helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive.

**Warnings**

Do not exceed recommended dosage.  
A persistent or chronic cough may be a sign of a serious condition. If cough persists for more than one week, tends to recur, or is accompanied by fever, rash, or persistent headache, consult a doctor.

**Do not use:**

- this product for persistent or chronic cough, such as occurs with smoking, asthma, chronic bronchitis or emphysema or if cough is accompanied by excessive phlegm (mucus), unless directed by a doctor.
- \* If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's Disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Keep out of reach of children

If pregnant or breast-feeding, ask a health professional before use. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

**Directions**

Do not exceed 6 doses in 24 hours

---

Age	Dose
Adults and Children 12 years of age and over	2 teaspoonfuls (10 ml) every 4 hours
Children under 12 years of age	Do not use

---

**Inactive ingredients**

artificial strawberry-banana flavor, citric acid, fd&c red #40, glycerin, methyl paraben, propylene glycol, propylparaben, sucrose, water

**Questions?**

Questions? call 1-800-560-5523 M-F 9 AM to 4 PM Eastern.



DEXTROMETHORPHAN HBR, GUAIFENESINA

# Bisolvine

ADULT

SOLUTION

JARABE

ADULTO

# Bisolvine

DEXTROMETORFAN HBR, GUAIFENESINA



Calma La Tos / Expectorante

Para el Alivio Temporal de La Tos y Congestión Causada por el Catarro Común



Contenido: 4 FL Oz (120 mL)

### Factores Farmacológicos

**Ingredientes activos (en cada 10 mL) Propósitos**  
Dextrometorfan Hbr 20 mg.....Alivio de la Tos  
Guafenesina 200 mg.....Expectorante

### Usos

Alivio temporal de la tos debido a irritación de la garganta y bronquios, asociada al resfriado común o gripa y ayuda a aliviar la flemas (mucosa) y secreciones bronquiales para hacer la tos más productiva.

### Advertencias

No exceder la dosis recomendada.

Una tos persistente puede ser señal de una condición seria. Si la tos perdura por más de una semana, regresa o está acompañada por fiebre, dolor de cabeza o erupciones en la piel, consulte a su médico.

### No usar

este producto para tos persistente o crónica, tal como ocurre si fuma, tiene asma, bronquitis crónica o enfisema o si la tos es acompañada por flemas excesiva (secreción) sola por indicación médica.  
si usted está tomando ahora alguna medicina bajo prescripción que contenga inhibidores de la monoaminooxidasa (MAOI) (ciertas drogas para la depresión, esquizofrenia, o condiciones emocionales, o enfermedad de Parkinson) o pasadas 2 semanas después de haber ingerido medicamentos que contengan MAOI. Si usted no está seguro que el medicamento que usted está tomando contiene MAOI, consulte a su doctor o a un farmacéutico antes de usarlo.

Mantener fuera del alcance de los niños.

Si está embarazada o lactando, pregunte a un profesional de la salud antes de usarlo. En caso de sobredosis accidental, busque ayuda médica o contacte a un Centro de Control de Envenenamiento inmediatamente.

**Instrucciones** No exceda más de 6 dosis en 24 horas.

edad	dosis
Adultos y niños 12 años de edad o más	2 cucharaditas (10 mL) cada 4 horas
Niños menores de 12 años de edad	No use

### Otra información

conservar este producto entre 15° a 30°C (59° a 86°F)

**Ingredientes inactivos** sabor artificial de fresa-banano, ácido cítrico, FD&C red #40, glicerina, metilparabeno, propilglicol, propilparabeno, sucrosa, agua.

**Preguntas?** llame 1-800-560-5223 L-V 9 AM a 4 PM Este.

EVIDENCIA DE SEGURIDAD: NO USAR ESTE PRODUCTO SI EL SELLO IMPRESO SOBRE LA TAPA DE LA BOTELLA ESTÁ TORCIDO, ROTO O FALTA.

SOLUTION

ADULT

# Bisolvine

DEXTROMETHORPHAN HBR, GUAIFENESINA



Cough Suppressant / Expectorant

For the Temporary Relief of Cough and Congestion due to the Common Cold



Contents: 4 FL Oz (120 mL)

### Drug Facts

**Active ingredients (in each 10 mL) Purposes**  
Dextromethorphan HBr 20 mg.....Cough suppressant  
Guafenesin 200 mg.....Expectorant

### Uses

Temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold and helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive.

### Warnings

Do not exceed recommended dosage.

A persistent cough may be a sign of a serious condition. If cough persists for more than one week, tends to recur, or is accompanied by fever, rash or persistent headache, consult a doctor.

### Do not use

this product for persistent or chronic cough, such as occurs with smoking, asthma, chronic bronchitis or emphysema or if cough is accompanied by excessive phlegm (mucus), unless directed by a doctor.  
if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's Disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Keep out of reach of children.

If pregnant or breast-feeding, ask a health professional before use. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

**Directions** Do not exceed 6 doses in 24 hours.

age	dose
Adults and children 12 years of age and over	2 teaspoons (10 mL) every 4 hours
Children under 12 years of age	Do not use

### Other information

store this product between 15° to 30°C (59° to 86°F)

**Inactive ingredients** artificial strawberry-banana flavor, citric acid, FD&C red #40, glycerin, methylparaben, propylene glycol, propylparaben, sucrose, water.

**Questions?** call 1-800-560-5223 M-F 9 AM to 4 PM Eastern.

TAMPER-EVIDENT: DO NOT USE THIS PRODUCT IF PRINTED CELLO-BAND OVER CAP IS TORN, BROKEN OR MISSING.

Distributed by/Distribuido por:  
Menper Distributors, Inc.  
11421 N.W. 107 Street Suite #24  
Miami, FL 33178  
www.menperdistributors.com



0 42279 80704 1



DEXTROMETHORPHAN HBR, GUAIFENESIN

**Bisolvine**

ADULT

SOLUTION

ADULTO  
**Bisolvine**  
DEXTROMETORFAN HBR, GUAIFENESINA

JARABE



Calma La Tos / Expectorante

Para el Alivio Temporal de  
La Tos y Congestión Causada  
por el Catarro Común



Contenido: 4 FL Oz (120 mL)

**Factores Farmacológicos**

**Ingredientes activos (en cada 10 mL) Propósitos**  
Dextrometorfan Hbr 20 mg.....Alivio de la Tos  
Guafenesina 200 mg.....Expectorante

**Usos**

Alivio temporal de la tos debido a irritación de la garganta y bronquios, asociado al resfriado común o gripa y ayuda a aliviar la flemas (mucosa) y secreciones bronquiales para hacer la tos más productiva.

**Advertencias**

No exceder la dosis recomendada. Una tos persistente puede ser señal de una condición seria. Si la tos perdura por más de una semana, regresa o está acompañada por fiebre, dolor de cabeza o erupciones en la piel, consulte a su médico.

**No usar**

este producto para tos persistente o crónica, tal como ocurre si fuma, tiene asma, bronquitis crónica o enfisema o si la tos es acompañada por flemas excesivas (secreción) sola por indicación médica. si usted está tomando ahora alguna medicina bajo prescripción que contenga inhibidores de la monoaminooxidasa (MAOI) (ciertas drogas para la depresión, alquitirinas, o condiciones emocionales, o enfermedad de Parkinson) o pasadas 2 semanas después de haber ingerido medicamentos que contengan MAOI. Si usted no está seguro que el medicamento que usted está tomando contiene MAOI, consulte a su doctor o a un farmacéutico antes de usarlo.

**Mantener fuera del alcance de los niños.**

Si está embarazada o lactando, pregunte a un profesional de la salud antes de usarlo. En caso de sobredosis accidental, busque ayuda médica o contacte a un Centro de Control de Envenenamiento inmediatamente.

**Instrucciones** No exceda más de 6 dosis en 24 horas.

edad	dosis
Adultos y niños 12 años de edad o más	2 cucharaditas (10 mL) cada 4 horas
Niños menores de 12 años de edad	No use

**Otra información**

conservar este producto entre 15° a 30°C (59° a 86°F)

**Ingredientes inactivos** sabor artificial de fresa-banano, ácido cítrico, FD&C red #40, glicerina, metilparabeno, propilglicol, propilparabeno, sucrosa, agua.

**Preguntas?** llame 1-800-560-5223 L-V 9 AM a 4 PM Este.

**EVIDENCIA DE SEGURIDAD:** NO USAR ESTE PRODUCTO SI EL SELLO IMPRESO SOBRE LA TAPA DE LA BOTELLA ESTÁ TORCIDO, ROTO O FALTA.

ADULT  
**Bisolvine**  
DEXTROMETHORPHAN HBR, GUAIFENESIN

SOLUTION



Cough Suppressant / Expectorant

For the Temporary Relief  
of Cough and Congestion  
due to the Common Cold



Contents: 4 FL Oz (120 mL)

**Drug Facts**

**Active ingredients (in each 10 mL) Purposes**  
Dextromethorphan HBr 20 mg.....Cough suppressant  
Guafenesin 200 mg.....Expectorant

**Uses**

Temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold and helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive.

**Warnings**

Do not exceed recommended dosage. A persistent cough may be a sign of a serious condition. If cough persists for more than one week, tends to recur, or is accompanied by fever, rash or persistent headache, consult a doctor.

**Do not use**

this product for persistent or chronic cough, such as occurs with smoking, asthma, chronic bronchitis or emphysema or if cough is accompanied by excessive phlegm (mucus), unless directed by a doctor. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's Disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Keep out of reach of children.**

If pregnant or breast-feeding, ask a health professional before use. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

**Directions** Do not exceed 6 doses in 24 hours.

age	dose
Adults and children 12 years of age and over	2 teaspoons (10 mL) every 4 hours
Children under 12 years of age	Do not use

**Other information**

store this product between 15° to 30°C (59° to 86°F)

**Inactive ingredients** artificial strawberry-banana flavor, citric acid, FD&C red #40, glycerin, methylparaben, propylene glycol, propylparaben, sucrose, water.

**Questions?** call 1-800-560-5223 M-F 9 AM to 4 PM Eastern.

**TAMPER-EVIDENT:** DO NOT USE THIS PRODUCT IF PRINTED CELLO-BAND OVER CAP IS TORN, BROKEN OR MISSING.

Distributed by/Distribuido por:  
**Menper Distributors, Inc.**  
11421 N.W. 107 Street Suite #24  
Miami, FL 33178  
www.menperdistributors.com



**BISOLVINE ADULT**

guaifenesin, dextromethorphan liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53145-325
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL

GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)

GUAIFENESIN

200 mg  
in 10 mL**Inactive Ingredients**

Ingredient Name	Strength
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SUCROSE</b> (UNII: C151H8M554)	

**Product Characteristics**

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	STRAWBERRY (Strawberry -Banana Flavor)	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53145-325-04	1 in 1 CARTON	02/01/2010	
1		120 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	02/01/2010	

**Labeler** - MENPER DISTRIBUTORS INC. (101947166)

Revised: 2/2022

MENPER DISTRIBUTORS INC.