

**ENTERIC COATED ASPIRIN REGULAR STRENGTH- aspirin tablet, coated**  
**Best Choice (Valu Merchandisers Company)**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredient (in each tablet)**

**Aspirin 325 mg (NSAID)\***

**\*nonsteroidal anti-inflammatory drug**

**Purpose**

**Pain reliever**

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - menstrual pain
  - minor arthritis pain
  - muscle pain
  - toothache
  - colds
- or as recommended by a doctor

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of **Reye's syndrome**, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

if you are allergic to aspirin or any other pain reliever/fever reducer.

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 12 years of age and over: take 1 to 2 tablets every 4 hours, while symptoms persist.
- drink a full glass of water with each dose.
- do not take more than 12 tablets in 24 hours unless directed by a doctor
- children under 12 years of age: do not use

**Other information**

- store at controlled room temperature 20°-25°C (68°-77°F)

**Inactive ingredients**

corn starch, croscarmellose sodium, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

## Questions or comments?

Call 1-877-753-3935 Monday-friday 9AM-5PM EST

## Principal Display Panel

COMPARE TO THE ACTIVE INGREDIENT IN ECOTRIN®†

Regular Strength

Enteric Coated Aspirin

Pain Reliever (NSAID)

Aspirin Regimen\*\*

Tablets

†This product is not manufactured or distributed by Prestige Brands, Inc., distributor of Ecotrin®.

\*\*Talk to your doctor before starting an aspirin regimen.

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BORKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

PROUDLY DISTRIBUTED BY:

VALU MERCHANDISERS

5000 KANSAS AVE

KANSAS CITY, KS 66106

www.bestchoicebrand.com

## Package label

**Drug Facts (continued)**  
Get medical help or contact a Poison Control Center right away.

**Directions**  
■ adults and children 12 years of age and over: take 1 to 2 tablets every 4 hours, while symptoms persist  
■ drink a full glass of water with each dose  
■ do not take more than 12 tablets in 24 hours unless directed by a doctor  
■ children under 12 years of age: do not use

**Other information**  
■ store at controlled room temperature 20°-25°C (68°-77°F)

**Inactive ingredients** corn starch, croscarmellose sodium, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

**Questions or comments?** Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

†This product is not manufactured or distributed by Prestige Brands, Inc., distributor of Ecotrin®.  
\*\*Talk to your doctor before starting an aspirin regimen.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BORKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

ected or any  
you ms, such as  
isease, liver  
e asthma  
if you are  
gns of  
at faint  
mit blood  
it better  
ical help right  
occurs  
days  
days  
tion.  
alth  
important not  
of pregnancy  
doctor  
born child  
of overdose.

PLD-81351  
Lot No.:  
Exp. Date:  
0  
Best Choice  
100% Guaranteed  
www.bestcho

B FC001735



**BEST CHOICE**

ibcbrand.com



PRODUCT DISTRIBUTED BY:  
WALLI MEDICATIONS, CO.  
5000 KANSAS AVE  
KANSAS CITY, KS 66106

**Drug Facts (continued)**

- take more or for a longer time than directed
- Do not use if you are allergic to aspirin or other pain reliever/fever reducer.

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problem or heartburn
- you have high blood pressure, heart disease, cirrhosis, or kidney disease
- you are taking a diuretic
- you have a history of stomach problem

**Ask a doctor or pharmacist before use** taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- you experience any of the following signs or symptoms
- stomach bleeding:
  - you have bloody or black stools
  - you have stomach pain that does not go away
- an allergic reaction occurs. Seek medical attention if:
  - new symptoms occur
  - ringing in the ears or loss of hearing (tinnitus) occurs
  - pain gets worse or lasts more than 3 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present

These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. It is especially important to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a health professional because it may cause problems in the fetus or complications during delivery. **Keep out of reach of children.** In case of emergency, call 1-800-368-8763.

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 325 mg (NSAID)\* ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscle pain
  - toothache
  - menstrual pain
  - minor arthritis pain
- or as recommended by a doctor

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

COMPARE TO THE ACTIVE INGREDIENT IN ECOTRIN®†

**REGULAR STRENGTH**

**Enteric Coated Aspirin**

**Best Choice®**

Actual Size

**Pain Reliever (NSAID)**  
Aspirin Regimen\*\*

**125 Tablets**  
325 mg EACH

0 3 2 6 1 5 0 1

**Best choice Regular Strength Enteric Coated Aspirin**

**ENTERIC COATED ASPIRIN REGULAR STRENGTH**

aspirin tablet, coated

**Product Information**

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:63941-370 |
| <b>Route of Administration</b> | ORAL           |                           |               |

**Active Ingredient/Active Moiety**

| Ingredient Name | Basis of Strength | Strength |
|-----------------|-------------------|----------|
|-----------------|-------------------|----------|

|  |         |        |
|--|---------|--------|
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | ASPIRIN | 325 mg |
|--|---------|--------|

### Inactive Ingredients

| Ingredient Name   | Strength |
|---|----------|
| STARCH, CORN (UNII: O8232NY3SJ)   |          |
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48)                                   |          |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)                                      |          |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8)                                      |          |
| HYPROMELLOSES (UNII: 3NXW29V3WO)  |          |
| METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13) |          |
| CELLULOSE, MICROCRYSTALLINE (UNII: OPIR32D61U)                            |          |
| MINERAL OIL (UNII: T5L8T28FGP)  |          |
| POLYSORBATE 80 (UNII: 6OZP39ZG8H)   |          |
| DIMETHICONE (UNII: 92RU3N3Y1O)  |          |
| SODIUM HYDROXIDE (UNII: 55X04QC32I)                                       |          |
| SODIUM LAURYL SULFATE (UNII: 368GB5141J)                                  |          |
| TALC (UNII: 7SEV7J4R1U)   |          |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP)                                       |          |
| TRIETHYL CITRATE (UNII: 8Z96QXD6UM)                                       |          |
| ALUMINUM OXIDE (UNII: LM26O6933)  |          |

### Product Characteristics

|          |        |              |          |
|----------|--------|--------------|----------|
| Color    | ORANGE | Score        | no score |
| Shape    | ROUND  | Size         | 11mm     |
| Flavor   |        | Imprint Code | T        |
| Contains |        |              |          |

### Packaging

| # | Item Code        | Package Description                                | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:63941-370-12 | 1 in 1 BOX   | 11/11/2011           | 11/11/2020         |
| 1 |                  | 125 in 1 BOTTLE; Type 0: Not a Combination Product |                      |                    |

### Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part343                                  | 11/11/2011           | 11/11/2020         |

**Labeler** - Best Choice (Valu Merchandisers Company) (868703513)

Revised: 7/2019

Best Choice (Valu Merchandisers Company)