

PYRANTEL PAMOATE- pyrantel pamoate suspension
ProMex LLC

Pyrantel Pamoate

Drug Facts

Active Ingredient (in each 1mL)

Pyrantel Pamoate 144 mg/mL
(50mg pyrantel base per mL)

Purpose

Anthelmintic

Use

for the treatment of pinworms.

Warnings

*Abdominal cramps, nausea, vomiting, diarrhea, headache, or dizziness sometimes occurs after taking this drug. If any of these conditions persist, consult a doctor.

*If you are pregnant or have liver disease, do not take this product unless directed by a doctor.

Directions

Shake well before use. Adults, children 12 years of age and over, and children 2 years to under 12 years of age: Oral dosage is a single dose of 5 milligrams of pyrantel base per pound, or 11 milligrams per kilogram, of body weight. Do not exceed 1 gram (1000 mg).

Weight	Dosage (Taken as a single dose)
Under 25 lbs or 2 yrs.	Don't use unless directed by a doctor
25-37 lbs	½ teaspoonful
38-62 lbs	1 teaspoonful
63-87 lbs	1 ½ teaspoonful
88-112 lbs	2 teaspoonful
113-137 lbs	2 ½ teaspoonful
138-162 lbs	3 teaspoonful
163-187lbs	3 ½ teaspoonful

over 188 lbs

4 teaspoonful

***Read package insert carefully before taking this medication.** Take only according to directions and do not exceed the recommended dosage unless directed by a doctor. Medication should only be taken one time as a single dose; do not repeat treatment unless directed by a doctor. When one individual in a household has pinworms, the entire household should be treated unless otherwise advised. See Warnings. If any worms other than pinworms are present before or after treatment consult a doctor. If any symptoms or pinworms are still present after treatment, consult a doctor.

*This product can be taken any time of day, with or without meals, it may be taken alone or with milk or fruit juice. Use of a laxative is not necessary prior to, during or after medication.

Keep out of Reach of Children. In case of accidental overdose/ingestion contact a poison control center immediately.

Inactive Ingredients

Purified Water, Glycerin, Sugar, Flavor, Xanthan Gum, Citric Acid, Sodium Benzoate and Potassium Sorbate

PRINCIPAL DISPLAY PANEL - 30 ml Bottle Carton

VERMISOL

SUSPENSION


1 oz (30 ml)

VERMISOL

DATOS DEL MEDICAMENTO

Ingrediente Activo (en cada mL)	Propósito
Pamoato de Pirante 144 mg/mL (Equivalente a 50mg de pirante base por mL)	Antiparasitario
Uso: para el tratamiento de lombrices intestinales	
Advertencia: Retortijones, náusea, vómito, diarrea, dolor de cabeza o mareo ocasionalmente puede ocurrir después de tomar este producto. Consulte a un médico si persiste alguno de estos síntomas.	
Si esta embarazada o tiene enfermedad de hígado no use si no es indicado por su médico	
Instrucciones: Agitar antes de usar. Adultos, niños de 12 años de edad o más, y niños mayores de 2 a menores de 12 años de edad: la dosis oral es una sola dosis de 5 miligramos de la base de pirante por libra, o 11 miligramos por kilogramo de peso corporal. No exceda de 1 gramo (1,000 mg).	

Peen Finca (fome una sola dosis)



S U S P E N S I O N

1 oz (30 ml)

Almacénese a temperaturas entre 15°-30°C (59°-86°F)

DATOS DEL MEDICAMENTO (continuado)

Mantener fuera del alcance de los niños. En caso de sobredosis / ingestión accidental contacte un centro de control de envenenamiento inmediatamente.

Ingredientes inactivos: Agua purificada, glicerina, azúcar, goma Xantán, sabor, ácido cítrico, benzoato de sodio, sorbato de potasio

Distributed By:

PRO-MEX®

P.O. Box 63-0716 - Miami, FL. 33163

UPC
184125-00026

Menos de 25 lbs o 2 años	No usar sin aviso de un médico
25-37 lb	½ cucharadita
38-62 lbs	1 cucharadita
63-87 lbs	1 ½ cucharadita
88-112 lbs	2 cucharadita
113-137 lbs	2 ½ cucharadita
138-162 lbs	3 cucharadita
163-187 lbs	3 ½ cucharadita
188 lbs o mas	4 cucharadita

■ **Lea el inserto cuidadosamente antes de tomar este medicamento.** Tome solamente según las instrucciones y no exceder la dosis recomendada a menos que lo indique un médico. Medicamento sólo debe tomarse una vez en una sola dosis, no repetir el tratamiento a menos que lo indique un médico. Cuando una persona en un hogar tiene oxiuros, toda la familia debe ser tratada a menos que se aconseje lo contrario. Véase Advertencias. Si algún gusano que no sea pinworms están presentes antes o después del tratamiento, consulte a un médico. Si los síntomas o los oxiuros todavía están presentes después del tratamiento, consulte a un médico.

■ Este producto se puede tomar a cualquier hora del día, con o sin comidas. Se puede tomar solo o con leche o jugo de frutas. El uso de un laxante no es necesario antes, durante o después de la medicación.

Tamper Evident: Do not use if seal is broken
No Usar Si el Sello esta roto

S U S P E N S I O N

(Pyrantel Pamoate)

VERMISOL

VERMISOL



S U S P E N S I O N

1 oz (30 ml)

Store at 15°-30°C (59°-86°F)



P
V006_009

Drug Facts

Active Ingredient (in each 1mL) **PURPOSE**
Pyrantel Pamoate 144 mg/mL Anthelmintic
(50mg pyrantel base per mL)

Use: for the treatment of pinworms.

Warnings: ■ Abdominal cramps, nausea, vomiting, diarrhea, headache, or dizziness sometimes occur after taking this drug. If any of these conditions persist consult a doctor.
■ If you are pregnant or have liver disease, do not take this product unless directed by a doctor.

Directions: Shake well before use. Adults, children 12 years of age and over, and children 2 years to under 12 years of age: Oral dosage is a single dose of 5 milligrams of pyrantel base per pound, or 11 milligrams per kilogram, of body weight. Do not exceed 1 gram (1000 mg).

Weight	Dosage (Taken as a single dose)
Under 25 lbs or 2 yrs	Don't use unless directed by a doctor
25-37 lbs	½ teaspoonful
38-62 lbs	1 teaspoonful
63-87 lbs	1 ½ teaspoonful
88-112 lbs	2 teaspoonful
113-137 lbs	2 ½ teaspoonful
138-162 lbs	3 teaspoonful
163-187 lbs	3 ½ teaspoonful
over 188 lbs	4 teaspoonful

■ Read package insert carefully before taking this medication. Take only according to directions and do not exceed the recommended dosage unless directed by a doctor. Medication should only be taken one time as a single dose; do not repeat treatment unless directed by a doctor. When one individual in a household has pinworms, the entire household should be treated unless otherwise advised. See Warnings. If any worms other than pinworms are present before or after treatment, consult a doctor. If any symptoms or pinworms are still present after treatment, consult a doctor.

■ This product can be taken any time of day, with or without meals. It may be taken alone or with milk or fruit juice. Use of a laxative is not necessary prior to, during, or after medication.

Keep out of Reach of Children. In case of accidental overdose/ingestion contact a poison control center immediately.

Inactive Ingredients: Purified Water, Glycerin, Sugar, Flavor, Xanthan Gum, Citric Acid, Sodium Benzoate and Potassium Sorbate

PC-134.00

PYRANTEL PAMOATE

pyrantel pamoate suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58988-1770
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Pyrantel Pamoate (UNII: 81BK194Z5M) (Pyrantel - UNII:4QIH0N49E7)	Pyrantel	144 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
Water (UNII: 059QF0KO0R)	
Citric Acid Monohydrate (UNII: 2968PHW8QP)	
Glycerin (UNII: PDC6A3C0OX)	
Sucrose (UNII: C151H8M554)	
Xanthan Gum (UNII: TTV12P4NEE)	
Sodium Benzoate (UNII: OJ245FE5EU)	
Potassium Sorbate (UNII: 1VPU26JZZ4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58988-1770-1	1 in 1 CARTON	09/27/2002	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M024	09/27/2002	

Labeler - ProMex LLC (789974388)

Revised: 12/2024

ProMex LLC