

COLD-EEZE DAYTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride liquid
ProPhase Labs, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Cold-EEZE®
Daytime Cold & Flu
Drug Facts

Active Ingredients (in each 20 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - sinus congestion and pressure
 - cough due to minor throat and bronchial irritation
 - minor aches and pains
 - sore throat
 - headache
 - temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- **with any other drug containing acetaminophen** (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
(see **Overdose warning**)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- **Adults and children 12 years and older:** 20 mL orally every 4 hours, not to exceed 6 doses in 24

hours.

- **Children under 12 years of age:** Do not use unless directed by a doctor

Other information

- each 20 mL contains:
sodium 12 mg
- tamper evident: do not use if foil seal under bottle cap is open or missing
- store between 15-30°C (59-86°F)
- do not refrigerate
- dosing cup provided

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions?

1-800-505- 2653

(M-F: 9AM-5PM EST)

You may also report side effects to this phone number

Distributed by:

ProPhase Labs, Inc.

PO Box 1349

Doylestown, PA 18901

PRINCIPAL DISPLAY PANEL - 266 mL Bottle Label

MAXIMUM STRENGTH

Cold-EEZE®

Daytime Cold & Flu

Acetaminophen 650 mg • Pain Reliever/Fever Reducer

Phenylephrine HCl 10 mg • Nasal Decongestant

Dextromethorphan HBr 20 mg • Cough Suppressant

Guaifenesin 400 mg • Expectorant

MULTI-SYMP TOM

Relief of:

- ☐ **Headache, Fever, Sore Throat & Aches & Pains**
- ☐ **Nasal/Sinus Congestion, Sinus Pressure**
- ☐ **Cough, Chest Congestion**
- ☐ **Thins & Loosens Mucus**

For Ages 12+

9 fl oz (266 mL)

3001000211-47005

NDC 61941-0300-1

MAXIMUM STRENGTH

Cold-EEZE®

Daytime Cold & Flu

Acetaminophen 650 mg • Pain Reliever/Fever Reducer
Phenylephrine HCl 10 mg • Nasal Decongestant
Dextromethorphan HBr 20 mg • Cough Suppressant
Guaifenesin 400 mg • Expectorant

MULTI-SYMPTOM

Relief of:

- ✓ Headache, Fever, Sore Throat & Aches & Pains
- ✓ Nasal/Sinus Congestion, Sinus Pressure
- ✓ Cough, Chest Congestion
- ✓ Thins & Loosens Mucus

For Ages 12+
9 fl oz (266 mL)

3001000211-47005

NDC 61941-0300-1

Drug Facts

Active Ingredients (in each 20 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - sinus congestion and pressure
 - cough due to minor throat and bronchial irritation
 - minor aches and pains
 - sore throat
 - headache
 - temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Drug Facts (continued)

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Drug Facts (continued)

or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

When using this product

Drug Facts (continued)

- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions,

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur

- do not use more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products

Drug Facts (continued)

- dose as follows or as directed by a doctor
- mL = milliliter
- **Adults and children 12 years and older:** 20 mL orally every 4 hours, not to exceed 6 doses in 24 hours.
- **Children under 12 years of age:** Do not use unless directed by a doctor

Other information

- each 20 mL contains: **sodium 12 mg**
- tamper evident: do not use if foil seal under bottle cap is open or missing
- store between 15-30°C (59-86°F)
- do not refrigerate
- dosing cup provided

Inactive ingredients
 anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions?
 1-800-505-2653 (M-F: 9AM-5PM EST)
 You may also report side effects to this phone number

Please visit our website at www.coldeeze.com

Questions or Comments:
Call: 1-800-505-2653 (M-F: 9AM-5PM EST)

Manufactured for and Distributed by:
 ProPhase Labs, Inc.
 PO Box 1349
 Doylestown, PA 18901

info@ProPhaseLabs.com

© 2015

PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

*Maximum Strength per 4 hour dose.

Tamper evident: do not use if foil seal under bottle cap is open or missing.

Dose every 4 hours.



3801000212-479006

LOT:

EXP:

COLD-EEZE DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:61941-0300
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Acetaminophen (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	650 mg in 20 mL	
Dextromethorphan Hydrobromide (UNII: 9D2RTI9KYH) (Dextromethorphan - UNII:7355X3ROTS)	Dextromethorphan Hydrobromide	20 mg in 20 mL	
Guaifenesin (UNII: 495W7451VQ) (Guaifenesin - UNII:495W7451VQ)	Guaifenesin	400 mg in 20 mL	
Phenylephrine Hydrochloride (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine Hydrochloride	10 mg in 20 mL	
Inactive Ingredients			

Ingredient Name	Strength
Anhydrous Citric Acid (UNII: XF417D3PSL)	
Edetate Disodium (UNII: 7FLD91C86K)	
FD&C Blue NO. 1 (UNII: H3R47K3TBD)	
FD&C Red NO. 40 (UNII: WZB9127XOA)	
Glycerin (UNII: PDC6A3C0OX)	
Propylene Glycol (UNII: 6DC9Q167V3)	
Propyl Gallate (UNII: 8D4SNN7V92)	
Water (UNII: 059QF0K00R)	
Sodium Benzoate (UNII: OJ245FE5EU)	
Sodium Citrate, Unspecified Form (UNII: 1Q73Q2JULR)	
Sorbitol (UNII: 506T60A25R)	
Sucralose (UNII: 96K6UQ3ZD4)	
Xanthan Gum (UNII: TTV12P4NEE)	

Product Characteristics

Color	BLUE	Score	
Shape		Size	
Flavor	CITRUS	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61941-0300-1	260 mL in 1 BOTTLE, UNIT-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	01/01/2016	

Labeler - ProPhase Labs, Inc. (620557298)

Establishment

Name	Address	ID/FEI	Business Operations
ProPhase Labs, Inc.		620557298	LABEL(61941-0300) , ANALYSIS(61941-0300) , REPACK(61941-0300)

Establishment

Name	Address	ID/FEI	Business Operations
Pharmaloz Manufacturing, Inc.		067101998	MANUFACTURE(61941-0300) , PACK(61941-0300) , REPACK(61941-0300)