

**VICKS DAYQUIL HIGH BLOOD PRESSURE COLD AND FLU- acetaminophen,
dextromethorphan liquid**
The Procter & Gamble Manufacturing Company

VICKS® DayQuil™
HIGH BLOOD PRESSURE COLD & FLU

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 20 mg

Purpose

Pain reliever/fever reducer
Cough suppressant

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses (30 mL each dose) in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over
 children 4 to under 12 yrs
 children under 4 yrs

30 mL per dose every 4 hrs
 ask a doctor
 do not use

Other information

- **each 30 mL contains:**sodium 26.06 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients

citric acid, FD&C yellow no. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions?**1-800-362-1683****TAMPER EVIDENT:****Do not use if printed shrinkband seal around the neck is broken or missing****DIST. BY****PROCTER & GAMBLE****CINCINNATI, OH 45202****PRINCIPAL DISPLAY PANEL - 236 ml Bottle Label**

VICKS®

DayQuil™

HIGH BLOOD PRESSURE

COLD & FLU

Acetaminophen,Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Cough

SUGAR & ALCOHOL FREE

DECONGESTANT FREE

Non-Drowsy

8 FL OZ (236 mL)

Drug Facts (continued)

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL per dose every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 30 mL contains: sodium 26.06 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients citric acid, FD&C yellow no. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions? 1-800-362-1683

Drug Facts (continued)

- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain or cough get worse or last more than 7 days

TAMPER EVIDENT:
Do not use if printed shrinkband seal around the neck is broken or missing
← PEEL BACK FOR DRUG FACTS

Drug Facts

Active ingredients (in each 30mL) **Purpose**
 Acetaminophen 650 mg Pain reliever/fever reducer
 Dextromethorphan HBr 20 mg Cough suppressant

Uses temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- minor aches & pains
- headache
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses (30 mL each dose) in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen

PARENTS:
Learn about how medicine abuse
www.StopMedicineAbuse.org

DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202
 Patents: www.pg.com/patents
 vicks.com

P&G
www.pg.com
 90321160

PLASTIC BOTTLE
 100% RECYCLED

2390004488



VICKS DAYQUIL HIGH BLOOD PRESSURE COLD AND FLU

acetaminophen, dextromethorphan liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-972
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	

SORBITOL (UNII: 506T60A25R)	
XANTHAN GUM (UNII: TTV12P4NEE)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	CITRUS	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-972-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/12/2022	
2	NDC:69423-972-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/12/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/12/2022	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2023

The Procter & Gamble Manufacturing Company