

**VICKS DAYQUIL HIGH BLOOD PRESSURE COLD AND FLU- acetaminophen,  
dextromethorphan liquid**  
**The Procter & Gamble Manufacturing Company**

-----  
**VICKS® DayQuil™**  
**HIGH BLOOD PRESSURE COLD & FLU**

***Drug Facts***

**Active ingredients (in each 30 mL)**

**Acetaminophen 650 mg**  
Dextromethorphan HBr 20 mg

**Purpose**

Pain reliever/fever reducer  
Cough suppressant

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses (30 mL each dose) in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- only use dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL per dose every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- **each 30 mL contains:** sodium 29 mg
- store at no greater than 25°C and do not refrigerate

**Inactive ingredients**

citric acid, FD&C yellow no. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

**Questions?**

**1-800-362-1683**

**TAMPER EVIDENT:**

**Do not use if printed shrinkband seal around the neck is broken or missing**

**DIST. BY**

**PROCTER & GAMBLE**

**CINCINNATI, OH 45202**

**PRINCIPAL DISPLAY PANEL - 236 ml Bottle Label**

VICKS®

DayQuil™

HIGH BLOOD PRESSURE

COLD & FLU

**Acetaminophen**, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Cough

SUGAR & ALCOHOL FREE

DECONGESTANT FREE

Non-Drowsy

8 FL OZ (236 mL)

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- pain or cough get worse or last more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- only use dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL per dose every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- each 30 mL dose contains: sodium 29 mg
- store at no greater than 25°C and do not refrigerate

**Inactive ingredients** citric acid, FD&C yellow no. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

**Questions?** 1-800-362-1683

**Drug Facts (continued)**

- more than 4 doses (30 mL each dose) in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**TAMPER EVIDENT:**  
Do not use if printed shrinkband seal around the neck is broken or missing  
→ **PEEL BACK FOR DRUG FACTS**  
ALSO SOLD AS DAYQUIL FOR PEOPLE WITH DIABETES

**Drug Facts**

**Active ingredients (in each 30mL)**      **Purpose**

Acetaminophen 650 mg..... Pain reliever/fever reducer

Dextromethorphan HBr 20 mg..... Cough suppressant

**Uses** temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202  
Patents: www.pg.com/patents

**PARENTS:**  
Learn how to keep kids safe from medicine abuse  
www.StopMedicineAbuse.org

Empty & Replace Cap

PLASTIC BOTTLE      PLASTIC CUP

how2recycle.info

P&G 21065560

3 23900 04488 3



## VICKS DAYQUIL HIGH BLOOD PRESSURE COLD AND FLU

acetaminophen, dextromethorphan liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69423-972
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	

