

PENAQUA SOL-G- penicillin g potassium powder, for solution
Bimeda Inc.

PenAqua Sol-G (penicillin G potassium)

DESCRIPTION

0.5 BU Front Panel

Penicillin G Potassium

Antibiotic for drinking water

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

For oral use in turkeys only

For Veterinary use only

Store at room temperature between 20° to 25°C (68° to 77°F), with excursions up to 40°C (104°F).

Keep out of the reach of children.

Restricted drug (California) -Use only as directed

Active Ingredient:

0.500 Billion Units of penicillin G potassium

Nonsterile

Net Weight: 313 grams (11 ounces)

Approved by FDA under ANADA #200-372

INDICATIONS

For the treatment of erysipelas in turkeys (caused by *Erysipelothrix rhusiopathiae*).

DOSAGE AND ADMINISTRATION

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

Directions

Combine contents and approximately 1-1/2 pints (710 mL) of water in a glass or plastic container. Stir to dissolve. Allow concentrated solution to stand until the foam disappears. The concentrated solution should be used up or discarded within one hour after preparation.

Automatic Watering Systems - Pour the concentrated solution into a glass or plastic container then add enough water to make 2.6 gallons (9.9 liters) of stock solution. [This amount of solution will medicate 333 gallons (1260 liters) of drinking water]. The

automatic waterer should be adjusted to deliver 1 ounce (30 mL) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions daily.

Gravity Flow Watering Systems - Pour the concentrated solution into enough water to make 333 gallons (1260 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

WARNINGS

Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment. Do not use in turkeys producing eggs for human consumption.

PRECAUTIONS

For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian.

DESCRIPTION

1 BU Front Panel

Penicillin G Potassium

Antibiotic for drinking water

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

For oral use in turkeys only

For Veterinary use only

Store at room temperature between 20° to 25°C (68° to 77°F), with excursions up to 40°C (104°F).

Keep out of the reach of children.

Restricted drug (California) -Use only as directed

Active Ingredient:

1 Billion Units of penicillin G potassium

Nonsterile

Net Weight: 626 grams (22 ounces)

Approved by FDA under ANADA #200-372

INDICATIONS AND USAGE

For the treatment of erysipelas in turkeys (caused by *Erysipelothrix rhusiopathiae*).

DOSAGE AND ADMINISTRATION

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

Directions

Combine contents and approximately 3 pints (1420 mL) of water in a glass or plastic container. Stir to dissolve. Allow concentrated solution to stand until the foam disappears. The concentrated solution should be used up or discarded within one hour after preparation.

Automatic Watering Systems - Pour the concentrated solution into a glass or plastic container then add enough water to make 5.2 gallons (19.7 liters) of stock solution. [This amount of solution will medicate 666 gallons (2521 liters) of drinking water]. The automatic waterer should be adjusted to deliver 1 ounce (30 mL) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions daily.

Gravity Flow Watering Systems - Pour the concentrated solution into enough water to make 666 gallons (2521 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

WARNINGS

Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment. Do not use in turkeys producing eggs for human consumption.

PRECAUTIONS

For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian.

Manufactured for:

Bimeda, Inc.

Le Sueur, MN 56058

www.bimeda.com

Rev. 08/2024

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

NDC 61133-4019-1

 **PEN AQUA SOL-G[®]**
(penicillin G potassium)

Antibiotic for drinking water

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

For oral use in turkeys only

For Veterinary use only

Store at room temperature between 20° to 25°C (68° to 77°F), with excursions up to 40°C (104°F).

Keep out of the reach of children.
Restricted drug (California)- Use only as directed

Active Ingredient:
0.500 Billion Units penicillin G potassium
Nonsterile

Net Weight: 313 grams (11 ounces)
Approved by FDA under ANADA # 200-372

 Bimeda

NDC 61133-4019-1
PenAqua Sol-G[®]
(penicillin G potassium)
Antibiotic for drinking water
.500 Billion Units penicillin G potassium

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

NDC 61133-4019-2

 **PEN AQUA SOL-G[®]**
(penicillin G potassium)

Antibiotic for drinking water

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

For oral use in turkeys only

For Veterinary use only

Store at room temperature between 20° to 25°C (68° to 77°F), with excursions up to 40°C (104°F).

Keep out of the reach of children.
Restricted drug (California)- Use only as directed

Active Ingredient:
1 Billion Units penicillin G potassium
Nonsterile

Net Weight: 626 grams (22 ounces)
Approved by FDA under ANADA # 200-372

 Bimeda

NDC 61133-4019-2

PenAqua Sol-G®
(penicillin G potassium)
Antibiotic for drinking water

1 Billion Units penicillin G potassium

PENAQUA SOL-G

penicillin g potassium powder, for solution

Product Information

| | | | |
|--------------------------------|--------------------------|---------------------------|----------------|
| Product Type | PRESCRIPTION ANIMAL DRUG | Item Code (Source) | NDC:61133-4019 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|--------------------------|-------------------------------------|
| Penicillin G Potassium (UNII: VL775ZTH4C) (Penicillin G - UNII:Q42T66VG0C) | Penicillin G | 500000000 [iU] in 500000000 [iU] |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|----------|------------------|----------------------------|-----------------------------|---------------------------|
| 1 | NDC:61133-4019-1 | 500000000 [iU] in 1 POUCH | | |
| 2 | NDC:61133-4019-2 | 1000000000 [iU] in 1 POUCH | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------------|---|-----------------------------|---------------------------|
| ANADA | ANADA200372 | 03/14/2025 | |

Labeler - Bimeda Inc. (060492923)

Revised: 3/2025

Bimeda Inc.