

**NYQUIL SEVERE PLUS VAPOCOOL COLD AND FLU- acetaminophen, phenylephrine hydrochloride, doxylamine succinate, and dextromethorphan hydrobromide tablet, coated**  
**NYQUIL SEVERE PLUS VAPOCOOL COLD AND FLU, CVP HEALTH- acetaminophen, phenylephrine hydrochloride, doxylamine succinate, and dextromethorphan hydrobromide tablet, coated**  
**NYQUIL SEVERE PLUS VAPOCOOL COLD AND FLU, TRAVEL BASIX- acetaminophen, phenylephrine hydrochloride, doxylamine succinate, and dextromethorphan hydrobromide tablet, coated**  
**Lil' Drug Store Products, Inc.**

-----  
**NyQuil™ Severe+ VapoCOOL™ Cold & Flu**

***Drug Facts***

| <b><i>Active ingredients (in each caplet)</i></b> | <b><i>Purpose</i></b>       |
|---|-----------------------------|
| Acetaminophen 325 mg                              | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg                        | Cough suppressant           |
| Doxylamine succinate 6.25 mg                      | Antihistamine               |
| Phenylephrine HCl 5 mg                            | Nasal decongestant          |

**Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Allergy Alert**

acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- do not exceed 8 caplets per 24 hrs

|                                 |                                  |
|---------------------------------|----------------------------------|
| adults & children 12 yrs & over | 2 caplets with water every 4 hrs |
| children 4 to under 12 yrs      | ask a doctor                     |
| children under 4 yrs            | do not use                       |

**Other information**

- do not exceed 25°C

**Inactive ingredients**

crospovidone, D&C Yellow No. 10 Aluminum Lake, FD&C Blue No. 1 Aluminum Lake, FD&C Blue No. 2 Aluminum Lake, FD&C Yellow No. 6 Aluminum Lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

**Questions?**

**1-800-362-1683**

**Product repackaged and distributed with permission of manufacturer by:**  
Lil' Drug Store Products, Inc., 9300 Earhart Lane SW, Cedar Rapids, IA 52404

# NyQuil™ Severe Plus VapoCOOL™ C & F - Lil' Drug Store PDP/Package

NyQuil™

SEVERE+

VICKS® VapoCOOL™

COLD & FLU

Acetaminophen, Phenylephrine HCl,

Doxylamine Succinate, Dextromethorphan HBr

- Minor Aches & Pains, Fever
- Nasal Congestion & Sinus Pressure
- Sneezing, Runny Nose
- Cough

Nighttime Relief

4

Coated Caplets

Lil'

DrugStore®

**NyQuil™ SEVERE+**  
**VICKS® VapoCOOL™**

**COLD & FLU**

**Acetaminophen, Phenylephrine HCl,  
Doxylamine Succinate, Dextromethorphan HBr**

- Minor Aches & Pains, Fever
- Nasal Congestion & Sinus Pressure
- Sneezing, Runny Nose
- Cough

Nighttime Relief

4 Coated Caplets

Lil' DrugStore

**PARENTS:**  
Learn about how medicines affect children  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**VapoCOOL™**  
COLD & FLU

**NyQuil™ SEVERE+**

**DO NOT USE IF THE POUCH IS BROKEN OR TORN.**  
**FAILURE TO FOLLOW THESE WARNINGS COULD RESULT IN SERIOUS CONSEQUENCES.**

**Drug Facts**

| Active Ingredients (In each caplet) | Purpose                     |
|-------------------------------------|-----------------------------|
| Acetaminophen 325 mg                | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg          | Cough suppressant           |
| Doxylamine succinate 6.25 mg        | Antihistamine               |
| Phenylephrine HCl 5 mg              | Nasal decongestant          |

**Uses** temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product



**Allergy Alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**LIFT PANEL FOR CONTINUED DRUG FACTS**



## NyQuil™ Severe + VapoCOOL™ C & F - CVP Health PDP/Package

NyQuil™

SEVERE+

VICKS® VapoCOOL™

COLD & FLU

Acetaminophen, Phenylephrine HCl,

Doxylamine Succinate, Dextromethorphan HBr

- Minor Aches & Pains, Fever
- Nasal Congestion & Sinus Pressure
- Sneezing, Runny Nose
- Cough


[caplet image]

**Nighttime Relief**



Coated Caplets

[CVP® HEALTH logo]



**NyQuil™ SEVERE+ VICKS® VapoCOOL™ - TRAVEL BASIX PDP/Package**

**NyQuil™**

**SEVERE+**

**VICKS® VapoCOOL™**

**COLD & FLU**

**Acetaminophen, Phenylephrine HCl,**

**Doxylamine Succinate, Dextromethorphan HBr**

- **Minor Aches & Pains, Fever**
- **Nasal Congestion & Sinus Pressure**
- **Sneezing, Runny Nose**

• Cough

Nighttime Relief

[caplet image]

4

Coated Caplets

[TRAVEL BASIX logo]



**NYQUIL SEVERE PLUS VAPOCOOL COLD AND FLU**  
 acetaminophen, phenylephrine hydrochloride, doxylamine succinate, and dextromethorphan hydrobromide tablet, coated

| Product Information   |                             |                    |                |
|---|-----------------------------|--------------------|----------------|
| Product Type  | HUMAN OTC DRUG              | Item Code (Source) | NDC:66715-9905 |
| Route of Administration   | ORAL                        |                    |                |
| Active Ingredient/Active Moiety   |                             |                    |                |
| Ingredient Name   | Basis of Strength           | Strength           |                |
| <b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)               | ACETAMINOPHEN               | 325 mg             |                |
| <b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg               |                |

|  |                               |         |
|--|-------------------------------|---------|
| <b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)                | DOXYLAMINE SUCCINATE          | 6.25 mg |
| <b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg   |

### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| <b>CROSPVIDONE (15 MPA.S AT 5%)</b> (UNII: 68401960MK)     |          |
| <b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)            |          |
| <b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)              |          |
| <b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)              |          |
| <b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)            |          |
| <b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)               |          |
| <b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)       |          |
| <b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A) |          |
| <b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)   |          |
| <b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)            |          |
| <b>STARCH, CORN</b> (UNII: O8232NY3SJ)                     |          |
| <b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)                  |          |
| <b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)                     |          |
| <b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)                        |          |
| <b>TALC</b> (UNII: 7SEV7J4R1U)                             |          |
| <b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)                 |          |

### Product Characteristics

|                 |       |                     |          |
|-----------------|-------|---------------------|----------|
| <b>Color</b>    | green | <b>Score</b>        | no score |
| <b>Shape</b>    | OVAL  | <b>Size</b>         | 19mm     |
| <b>Flavor</b>   |       | <b>Imprint Code</b> | NQ       |
| <b>Contains</b> |       |                     |          |

### Packaging

| # | Item Code        | Package Description                             | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:66715-9905-2 | 2 in 1 CARTON                                   | 05/01/2019           |                    |
| 1 |                  | 2 in 1 POUCH; Type 0: Not a Combination Product |                      |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012                                     | 05/01/2019           |                    |

**NYQUIL SEVERE PLUS VAPOCOOL COLD AND FLU, CVP HEALTH**



acetaminophen, phenylephrine hydrochloride, doxylamine succinate, and dextromethorphan hydrobromide tablet, coated

### Product Information

|                                |                |                           |                |
|--------------------------------|----------------|---------------------------|----------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:66715-6581 |
| <b>Route of Administration</b> | ORAL           |                           |                |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength             | Strength |
|--|-------------------------------|----------|
| <b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)                    | ACETAMINOPHEN                 | 325 mg   |
| <b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)      | PHENYLEPHRINE HYDROCHLORIDE   | 5 mg     |
| <b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)                | DOXYLAMINE SUCCINATE          | 6.25 mg  |
| <b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg    |

### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| <b>CROSPVIDONE (15 MPA.S AT 5%)</b> (UNII: 68401960MK)     |          |
| <b>D&amp;C YELLOW NO. 10</b> (UNII: 355W5USQ3G)            |          |
| <b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)              |          |
| <b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)              |          |
| <b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)            |          |
| <b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)               |          |
| <b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)       |          |
| <b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A) |          |
| <b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)   |          |
| <b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)            |          |
| <b>STARCH, CORN</b> (UNII: O8232NY3SJ)                     |          |
| <b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)                  |          |
| <b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)                     |          |
| <b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)                        |          |
| <b>TALC</b> (UNII: 7SEV7J4R1U)                             |          |
| <b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)                 |          |

### Product Characteristics

|                 |       |                     |          |
|-----------------|-------|---------------------|----------|
| <b>Color</b>    | green | <b>Score</b>        | no score |
| <b>Shape</b>    | OVAL  | <b>Size</b>         | 19mm     |
| <b>Flavor</b>   |       | <b>Imprint Code</b> | NQ       |
| <b>Contains</b> |       |                     |          |

### Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-----------|---------------------|----------------------|--------------------|
|---|-----------|---------------------|----------------------|--------------------|

|   |                  |   |            |
|---|------------------|---|------------|
| 1 | NDC:66715-6581-2 | 2 in 1 CARTON                                   | 08/01/2022 |
| 1 |                  | 2 in 1 POUCH; Type 0: Not a Combination Product |            |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012                                     | 08/01/2022           |                    |

## NYQUIL SEVERE PLUS VAPOCOOL COLD AND FLU, TRAVEL BASIX

acetaminophen, phenylephrine hydrochloride, doxylamine succinate, and dextromethorphan hydrobromide tablet, coated

### Product Information

|                         |                |                    |                |
|-------------------------|----------------|--------------------|----------------|
| Product Type            | HUMAN OTC DRUG | Item Code (Source) | NDC:66715-6405 |
| Route of Administration | ORAL           |                    |                |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength             | Strength |
|--|-------------------------------|----------|
| <b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)                    | ACETAMINOPHEN                 | 325 mg   |
| <b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)      | PHENYLEPHRINE HYDROCHLORIDE   | 5 mg     |
| <b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)                | DOXYLAMINE SUCCINATE          | 6.25 mg  |
| <b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg    |

### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| <b>CROSPVIDONE (15 MPA.S AT 5%)</b> (UNII: 68401960MK)     |          |
| <b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)            |          |
| <b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)              |          |
| <b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)              |          |
| <b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)            |          |
| <b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)               |          |
| <b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)       |          |
| <b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A) |          |
| <b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)   |          |
| <b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)            |          |
| <b>STARCH, CORN</b> (UNII: O8232NY3S)                      |          |
| <b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)                  |          |
| <b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)                     |          |
| <b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)                        |          |
| <b>TALC</b> (UNII: 7SEV7J4R1U)                             |          |

**TITANIUM DIOXIDE** (UNII: 15FIX9V2JP)

### Product Characteristics

|                 |       |                     |          |
|-----------------|-------|---------------------|----------|
| <b>Color</b>    | green | <b>Score</b>        | no score |
| <b>Shape</b>    | OVAL  | <b>Size</b>         | 19mm     |
| <b>Flavor</b>   |       | <b>Imprint Code</b> | NQ       |
| <b>Contains</b> |       |                     |          |

### Packaging

| # | Item Code        | Package Description                             | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:66715-6405-2 | 2 in 1 CARTON                                   | 11/29/2022           |                    |
| 1 |                  | 2 in 1 POUCH; Type 0: Not a Combination Product |                      |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012                                     | 11/29/2022           |                    |

**Labeler** - Lil' Drug Store Products, Inc. (093103646)

Revised: 10/2023

Lil' Drug Store Products, Inc.