



**SUCRALFATE- sucralfate tablet**  
**American Health Packaging**

-----  
**Sucralfate Tablets, USP**  
**8469501/0622**  
**Rx only**

**DESCRIPTION**

Sucralfate is an  $\alpha$ -D-glucopyranoside,  $\beta$ -D-fructofuranosyl-, octakis-(hydrogen sulfate), aluminum complex. It has the following structural formula:



( $x = 8$  to  $10$  and  $y = 22$  to  $31$ )  $\text{R} = \text{SO}_3\text{Al}(\text{OH})_2$

Tablets for oral administration contain 1 g of sucralfate, USP.

Also contain: povidone, magnesium stearate, and colloidal silicon dioxide.

Therapeutic category: antiulcer.

**CLINICAL PHARMACOLOGY**

Sucralfate is only minimally absorbed from the gastrointestinal tract. The small amounts of the sulfated disaccharide that are absorbed are excreted primarily in the urine.

Although the mechanism of sucralfate's ability to accelerate healing of duodenal ulcers remains to be fully defined, it is known that it exerts its effect through a local, rather than systemic, action. The following observations also appear pertinent:

1. Studies in human subjects and with animal models of ulcer disease have shown that sucralfate forms an ulcer-adherent complex with proteinaceous exudate at the ulcer site.
2. *In vitro*, a sucralfate-albumin film provides a barrier to diffusion of hydrogen ions.
3. In human subjects, sucralfate given in doses recommended for ulcer therapy inhibits pepsin activity in gastric juice by 32%.
4. *In vitro*, sucralfate adsorbs bile salts.

These observations suggest that sucralfate's antiulcer activity is the result of formation of an ulceradherent complex that covers the ulcer site and protects it against further attack by acid, pepsin, and bile salts. There are approximately 14 to 16 mEq of acid-neutralizing capacity per 1 g dose of sucralfate.

## Clinical Trials

### Acute Duodenal Ulcer

Over 600 patients have participated in well-controlled clinical trials worldwide. Multicenter trials conducted in the United States, both of them placebo-controlled studies with endoscopic evaluation at 2 and 4 weeks, showed:

#### STUDY 1

| Treatment Groups | Ulcer Healing/ No. Patients |                |
|------------------|-----------------------------|----------------|
|                  | 2 wk                        | 4 wk (Overall) |
| Sucralfate       | 37/105 (35.2%)              | 82/109 (75.2%) |
| Placebo          | 26/106 (24.5%)              | 68/107 (63.6%) |

#### STUDY 2

| Treatment Groups | Ulcer Healing/ No. Patients |                |
|------------------|-----------------------------|----------------|
|                  | 2 wk                        | 4 wk (Overall) |
| Sucralfate       | 8/24 (33%)                  | 22/24 (92%)    |
| Placebo          | 4/31 (13%)                  | 18/31 (58%)    |

The sucralfate-placebo differences were statistically significant in both studies at 4 weeks but not at 2 weeks. The poorer result in the first study may have occurred because sucralfate was given 2 hours after meals and at bedtime rather than 1 hour before meals and at bedtime, the regimen used in international studies and in the second United States study. In addition, in the first study liquid antacid was utilized as needed, whereas in the second study antacid tablets were used.

### Maintenance Therapy After Healing of Duodenal Ulcer

Two double-blind randomized placebo-controlled U.S. multicenter trials have demonstrated that sucralfate (1 g bid) is effective as maintenance therapy following healing of duodenal ulcers.

In one study, endoscopies were performed monthly for 4 months. Of the 254 patients who enrolled, 239 were analyzed in the intention-to-treat life table analysis presented below.

| Duodenal Ulcer Recurrence Rate (%) |                   |      |      |      |      |
|------------------------------------|-------------------|------|------|------|------|
| Drug                               | Months of Therapy |      |      |      |      |
|                                    | n                 | 1    | 2    | 3    | 4    |
| Sucralfate                         | 122               | 20 * | 30 * | 38 † | 42 † |
| Placebo                            | 117               | 33   | 46   | 55   | 63   |

\* P<0.05

† P<0.01

In this study, prn antacids were not permitted.

In the other study, scheduled endoscopies were performed at 6 and 12 months, but for-cause endoscopies were permitted as symptoms dictated. Median symptom scores between the sucralfate and placebo groups were not significantly different. A life table intention-to-treat analysis for the 94 patients enrolled in the trial had the following results:

| <b>Duodenal Ulcer Recurrence Rate (%)</b> |    |          |           |
|---|----|----------|-----------|
| Drug                                      | n  | 6 months | 12 months |
| Sucralfate                                | 48 | 19 *     | 27 *      |
| Placebo                                   | 46 | 54       | 65        |

\* P<0.002

In this study, prn antacids were permitted.

Data from placebo-controlled studies longer than 1 year are not available.

## **INDICATIONS AND USAGE**

Sucralfate tablets are indicated in:

- Short-term treatment (up to 8 weeks) of active duodenal ulcer. While healing with sucralfate may occur during the first week or two, treatment should be continued for 4 to 8 weeks unless healing has been demonstrated by x-ray or endoscopic examination.
- Maintenance therapy for duodenal ulcer patients at reduced dosage after healing of acute ulcers.

## **CONTRAINDICATIONS**

Sucralfate tablets are contraindicated in patients with known hypersensitivity reactions to the active substance or to any of the excipients.

## **PRECAUTIONS**

The physician should read the " **PRECAUTIONS**" section when considering the use of this drug in pregnant or pediatric patients, or patients of childbearing potential.

Duodenal ulcer is a chronic, recurrent disease. While short-term treatment with sucralfate can result in complete healing of the ulcer, a successful course of treatment with sucralfate should not be expected to alter the post healing frequency or severity of duodenal ulceration.

Isolated reports of sucralfate tablet aspiration with accompanying respiratory complications have been received. Therefore, sucralfate tablets should be used with caution by patients who have known conditions that may impair swallowing, such as

recent or prolonged intubation, tracheostomy, prior history of aspiration, dysphagia, or any other conditions that may alter gag and cough reflexes, or diminish oropharyngeal coordination or motility.

### **Special Populations: Chronic Renal Failure and Dialysis Patients**

When sucralfate is administered orally, small amounts of aluminum are absorbed from the gastrointestinal tract. Concomitant use of sucralfate with other products that contain aluminum, such as aluminum-containing antacids, may increase the total body burden of aluminum.

Patients with normal renal function receiving the recommended doses of sucralfate and aluminum-containing products adequately excrete aluminum in the urine. Patients with chronic renal failure or those receiving dialysis have impaired excretion of absorbed aluminum. In addition, aluminum does not cross dialysis membranes because it is bound to albumin and transferrin plasma proteins. Aluminum accumulation and toxicity (aluminum osteodystrophy, osteomalacia, encephalopathy) have been described in patients with renal impairment. Sucralfate should be used with caution in patients with chronic renal failure.

### **Drug Interactions**

Some studies have shown that simultaneous sucralfate administration in healthy volunteers reduced the extent of absorption (bioavailability) of single doses of the following: cimetidine, digoxin, fluoroquinolone antibiotics, ketoconazole, l-thyroxine, phenytoin, quinidine, ranitidine, tetracycline, and theophylline. Subtherapeutic prothrombin times with concomitant warfarin and sucralfate therapy have been reported in spontaneous and published case reports. However, two clinical studies have demonstrated no change in either serum warfarin concentration or prothrombin time with the addition of sucralfate to chronic warfarin therapy.

The mechanism of these interactions appears to be nonsystemic in nature, presumably resulting from sucralfate binding to the concomitant agent in the gastrointestinal tract. In all case studies to date (cimetidine, ciprofloxacin, digoxin, norfloxacin, ofloxacin, and ranitidine), dosing the concomitant medication 2 hours before sucralfate eliminated the interaction. Because of the potential of Sucralfate to alter the absorption of some drugs, Sucralfate tablets should be administered separately from other drugs when alterations in bioavailability are felt to be critical. In these cases, patients should be monitored appropriately.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Chronic oral toxicity studies of 24 months' duration were conducted in mice and rats at doses up to 1 g/kg (12 times the human dose).

There was no evidence of drug-related tumorigenicity. A reproduction study in rats at doses up to 38 times the human dose did not reveal any indication of fertility impairment. Mutagenicity studies were not conducted.

### **Pregnancy**

#### **Teratogenic effects**

Teratogenicity studies have been performed in mice, rats, and rabbits at doses up to 50

times the human dose and have revealed no evidence of harm to the fetus due to sucralfate. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

### **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when sucralfate is administered to a nursing woman.

### **Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

### **Geriatric Use**

Clinical studies of sucralfate tablets did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy (See **DOSAGE AND ADMINISTRATION**).

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function (See **PRECAUTIONS Special Populations: Chronic Renal Failure and Dialysis Patients**). Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

## **ADVERSE REACTIONS**

Adverse reactions to sucralfate in clinical trials were minor and only rarely led to discontinuation of the drug. In studies involving over 2700 patients treated with sucralfate tablets, adverse effects were reported in 129 (4.7%).

Constipation was the most frequent complaint (2%). Other adverse effects reported in less than 0.5% of the patients are listed below by body system:

**Gastrointestinal:** diarrhea, nausea, vomiting, gastric discomfort, indigestion, flatulence, dry mouth

**Dermatological:** pruritus, rash

**Nervous System:** dizziness, insomnia, sleepiness, vertigo

**Other:** back pain, headache

**Post-marketing:** cases of hypersensitivity have been reported with the use of sucralfate tablets, including dyspnea, lip swelling, pruritus, rash, and urticaria.

Cases of anaphylactic reactions, bronchospasm, laryngeal edema, edema of the mouth, pharyngeal edema, respiratory tract edema and swelling of the face have been reported with an unknown oral formulation of sucralfate.

Bezoars have been reported in patients treated with sucralfate. The majority of patients had underlying medical conditions that may predispose to bezoar formation (such as delayed gastric emptying) or were receiving concomitant enteral tube feedings.

Inadvertent injection of insoluble sucralfate and its insoluble excipients has led to fatal complications, including pulmonary and cerebral emboli. Sucralfate is **not** intended for intravenous administration.

## OVERDOSAGE

Due to limited experience in humans with overdosage of sucralfate, no specific treatment recommendations can be given. Acute oral toxicity studies in animals, however, using doses up to 12 g/kg body weight, could not find a lethal dose. Sucralfate is only minimally absorbed from the gastrointestinal tract. Risks associated with acute overdosage should, therefore, be minimal. In rare reports describing sucralfate overdose, most patients remained asymptomatic. Those few reports where adverse events were described included symptoms of dyspepsia, abdominal pain, nausea, and vomiting.

## DOSAGE AND ADMINISTRATION

**Active Duodenal Ulcer.** The recommended adult oral dosage for duodenal ulcer is 1 g four times per day on an empty stomach.

Antacids may be prescribed as needed for relief of pain but should not be taken within one-half hour before or after sucralfate.

While healing with sucralfate may occur during the first week or two, treatment should be continued for 4 to 8 weeks unless healing has been demonstrated by x-ray or endoscopic examination.

**Maintenance Therapy:** The recommended adult oral dosage is 1 g twice a day.

**Elderly:** In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy (See **PRECAUTIONS, Geriatric Use**).

Call your doctor for medical advice about side effects. You may report side effects to Nostrum Laboratories, Inc. at [quality@nostrumpharma.com](mailto:quality@nostrumpharma.com) or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

## HOW SUPPLIED

Sucralfate 1 g tablets, USP are supplied in unit dose packages of 100 (10 x 10). White, oblong, bisected tablets debossed with "N" and "S1" on one side.

Unit dose packages of 100 (10 x 10) NDC 60687-695-01

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

**FOR YOUR PROTECTION:** Do not use if blister is torn or broken.

## **PACKAGING INFORMATION**

American Health Packaging unit dose blisters (see How Supplied section) contain drug product from Nostrum Laboratories, Inc. as follows:

(1 g / 100 UD) NDC 60687-695-01 packaged from NDC 29033-003

Distributed by:

**American Health Packaging**

Columbus, OH 43217

**8469501/0622**

**Package/Label Display Panel - Carton - 1 g**

NDC 60687-**695**-01

**Sucralfate**

Tablets, USP

**1 gram**

100 Tablets (10 x 10)

Rx Only



(01) 0 03 60687 695 01 3

NDC 60687-**695**-01

**Sucralfate**

Tablets, USP

**1 gram**

100 Tablets (10 x 10)

Rx Only

**Each Tablet Contains:**

Sucralfate, USP ..... 1 gram

**Usual Dosage:** See full prescribing information.

**Store** at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

**FOR YOUR PROTECTION:** Do not use if blister is torn or broken.

The drug product contained in this package is from NDC # 29033-003, Nostrum Laboratories, Inc.

Distributed by: American Health Packaging, Columbus, Ohio 43217

Scan for current  
Prescribing Information →



769501  
0469501/0924

NDC 60687- **695**-01

**Sucralfate**  
Tablets, USP

**1 gram**

**100 Tablets (10 x 10)**

**Rx Only**

**Each Tablet Contains:**

Sucralfate, USP ..... 1 gram

**Usual Dosage:** See full prescribing information.

**Store** at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

**FOR YOUR PROTECTION:** Do not use if blister is torn or broken.

The drug product contained in this package is from NDC # 29033-003, Nostrum Laboratories, Inc.

Distributed by: American Health Packaging, Columbus, Ohio 43217

769501

0469501/0924

**Package/Label Display Panel - Blister - 1 g**



Sucralfate  
Tablet, USP

1 gram

| SUCRALFATE   |                         |  |                              |                    |
|--|-------------------------|--|------------------------------|--------------------|
| sucralfate tablet  |                         |  |                              |                    |
|  |                         |  |                              |                    |
| Product Information  |                         |  |                              |                    |
| Product Type   | HUMAN PRESCRIPTION DRUG | Item Code (Source)                                     | NDC:60687-695(NDC:29033-003) |                    |
| Route of Administration                                      | ORAL                    |  |                              |                    |
|  |                         |  |                              |                    |
| Active Ingredient/Active Moiety                              |                         |  |                              |                    |
| Ingredient Name  |                         |  | Basis of Strength            | Strength           |
| SUCRALFATE (UNII: XX73205DH5) (SUCRALFATE - UNII:XX73205DH5) |                         |  | SUCRALFATE                   | 1 g                |
|  |                         |  |                              |                    |
| Inactive Ingredients   |                         |  |                              |                    |
| Ingredient Name  |                         |  |                              | Strength           |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)                     |                         |  |                              |                    |
| MAGNESIUM STEARATE (UNII: 70097M6I30)                        |                         |  |                              |                    |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)                           |                         |  |                              |                    |
|  |                         |  |                              |                    |
| Product Characteristics                                      |                         |  |                              |                    |
| Color  | white (White)           | Score  | 2 pieces                     |                    |
| Shape  | OVAL (Oblong)           | Size   | 19mm                         |                    |
| Flavor   |                         | Imprint Code   | N;S1                         |                    |
| Contains   |                         |  |                              |                    |
|  |                         |  |                              |                    |
| Packaging  |                         |  |                              |                    |
| #  | Item Code               | Package Description                                    | Marketing Start Date         | Marketing End Date |
| 1  | NDC:60687-695-01        | 100 in 1 CARTON  | 05/11/2023                   | 07/31/2026         |
| 1  | NDC:60687-695-11        | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product |                              |                    |
|  |                         |  |                              |                    |



### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA               | ANDA074415                               | 05/11/2023           | 07/31/2026         |

**Labeler** - American Health Packaging (929561009)

### Establishment

| Name                      | Address | ID/FEI    | Business Operations |
|---------------------------|---------|-----------|---------------------|
| American Health Packaging |         | 929561009 | repack(60687-695)   |

Revised: 6/2025

American Health Packaging