

ARNICA GEL- arnica montana gel
Sled Distribution, LLC

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

81417-005 MY MEDIC ARNICA GEL

Active ingredients:

Arnica montana 7%

Purpose

Relieves muscle pain and stiffness, swelling from injuries, discoloration from bruises.

Uses Temporarily relieves muscle pain and stiffness due to:

- minor injuries
- overexertion
- falls
- reduces symptoms of bruising such as : discoloration, pain, swelling.

WarningsFor external use only.

Do not use if you are allergic to Arnica montana or any of this product's inactive ingredients.

When using this product

- avoid contact with eyes, mucous membranes, wounds, damaged or irritated skin
- dryness or irritation may occur
- use only as directed
- do not tightly wrap or bandage the treated area
- do not apply heat or ice to the treated area immediately before or after use.

Stop use and ask a doctor if

- Conditions persist for more than 3 days or worsens
- Symptoms clear up and occur again within a few days.

Keep out of reach of children

- If swallowed, get medical help or contact a Poison Control Center immediately.

Directions

Apply a thin layer of gel to affected area as soon as possible after minor injury. Repeat 3 times a day or as needed. If heat or ice is applied, wait 5 minutes before applying ointment.

Other information do not use if the sachet seal is broken.

Alcohol, carbomer, purified water, propylene glycol, sodium hydroxide


MY MEDIC™

TEAR HERE

Arnica Gel

Anti-inflammatory ointment which reduces swelling & pain

Topical



3.5 g
(.125 fl oz)



EXPERIENCE ELEVATED FIRST AID

Drug Facts

| Active ingredients | Purpose |
|--------------------|--|
| Arnica montana 7% | Relieves muscle pain & stiffness, swelling from injuries, discoloration from bruises |

Uses temporarily relieves muscle pain and stiffness due to:

- minor injuries
- overexertion
- falls
- reduces symptoms of bruising such as:
- discoloration
- pain
- swelling

Warnings For external use only.
Do not use if you are allergic to Arnica montana or any of this product's inactive ingredients.

When using this product

- avoid contact with eyes, mucous membranes, wounds, damaged or irritated skin
- dryness or irritation may occur
- use only as directed
- do not tightly wrap or bandage the treated area
- do not apply heat or ice to the treated area immediately before or after use.

Stop use and ask a doctor if

- conditions persist for more than 3 days or worsens
- symptoms clear up and occur again within a few days.

Keep out of reach of children
If swallowed, get medical help or contact a Poison Control Center right away.

Directions Apply a thin layer of gel to affected area as soon as possible after minor injury. Repeat 3 times a day or as needed. If heat or ice is applied, wait 5 minutes before applying ointment.

Other information do not use if the sachet seal is broken.

Inactive ingredients alcohol, carbomer, purified water, propylene glycol, sodium hydroxide

MFG: XXXXXXXX SCAN ME FOR MORE KNOWLEDGE
EXP: XXXXXXXX SKILLS & REPAIRS

MADE IN CHINA

DISTRIBUTED BY: SLED DISTRIBUTION LLC DBA MYMEDIC 120 North Redwood Rd. Ste. A, North Salt Lake, UT 84054



ARNICA GEL

arnica montana gel

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:81417-005 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|--------------|
| ARNICA MONTANA (UNII: O80TY208Z W) (ARNICA MONTANA - UNII:O80TY208Z W) | ARNICA MONTANA | 7 g in 100 g |

Inactive Ingredients

| Ingredient Name | Strength |
|----------------------------|----------|
| ALCOHOL (UNII: 3K9958V90M) | |

CARBOMER INTERPOLYMER TYPE A (ALLYL SUCROSE CROSSLINKED) (UNII: 59TL3WG5CO)

WATER (UNII: 059QF0KO0R)

SODIUM HYDROXIDE (UNII: 55X04QC32I)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:81417-005-01 | 3.5 g in 1 POUCH; Type 0: Not a Combination Product | 08/29/2022 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|------------------------|--|----------------------|--------------------|
| unapproved homeopathic | | 08/29/2022 | |

Labeler - Sled Distribution, LLC (079772888)

Revised: 1/2024

Sled Distribution, LLC