

STOMACH RELIEF MAXIMUM STRENGTH- bismuth subsalicylate liquid
Strategic Sourcing Services LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each 30 mL)

Bismuth subsalicylate 1050 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - fullness
 - belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- a sodium-restricted diet
- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over:
 - 30 mL (1 dose) every 1 hour as needed
 - do not take more than 4 doses (120 mL) in 24 hours
 - use until diarrhea stop but not more than 2 days
- children under 12 years of age:ask a doctor
- drink plenty of clear fluids to help prevent dehydration cause by diarrhea

Other information

- each 30 mL contains: **sodium 19 mg**
- each 30 mL contains: **salicylate 455 mg**
- sugar free
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

Compare to **Pepto-Bismol® active ingredient***

Stomach Relief

Bismuth subsalicylate 1050 mg

Upset Stomach Reliever/Antidiarrheal

Relieves upset stomach, heartburn, indigestion, nausea & diarrhea

Maximum strength

Original flavor

FL OZ (mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Another Quality Product Distributed by McKesson

One Post Street, San Francisco, CA 94104

www.healthmart.com/healthmartbrand

Package Label

Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ mL = milliliter ■ shake well before using
■ measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product ■ adults and children 12 years and over: ■ 30 mL (1 dose) every 1 hour as needed ■ do not take more than 4 doses (120 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years of age: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information ■ each 30 mL contains: sodium 19 mg
■ each 30 mL contains: salicylate 455 mg
■ sugar free ■ keep tightly closed ■ protect from freezing
■ avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

Another Quality Product Distributed by McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee
www.healthmart.com/healthmartbrand

PLD-A390A
LB003699

0 52569 13521 1

Health Mart PHARMACY
compare to Pepto-Bismol®
active ingredient*

stomach relief
bismuth subsalicylate 1050 mg
upset stomach reliever/antidiarrheal

relieves upset stomach, heartburn, indigestion, nausea & diarrhea

maximum strength
original flavor

8 fl oz (237 mL)

Health Mart Stomach Relief

Drug Facts TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Active ingredient (in each 30 mL) Bismuth subsalicylate 1050 mg.....Upset stomach reliever/Antidiarrheal **Purpose**

Uses relieves ■ travelers' diarrhea ■ diarrhea
■ upset stomach due to overindulgence in food and drink, including
■ heartburn ■ gas ■ indigestion
■ nausea ■ fullness ■ belching

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Contains salicylate. Do not take if you are
■ allergic to salicylates (including aspirin)
■ taking other salicylate products

Do not use if you have
■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have
■ a sodium-restricted diet ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
■ diabetes ■ gout ■ arthritis ■ anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs
■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. ▶

STOMACH RELIEF MAXIMUM STRENGTH

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62011-0127	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD118YE, SALICYLIC ACID - UNII:O414PZ4LPZ)		BISMUTH SUBSALICYLATE	1050 mg in 30 mL	
Inactive Ingredients				
Ingredient Name			Strength	
BENZOIC ACID (UNII: 8SKN0B0MIM)				
D&C RED NO. 22 (UNII: 1678RKX8RT)				
D&C RED NO. 28 (UNII: 767IP0Y5NH)				
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)				
WATER (UNII: 059QF0K00R)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
SALICYLIC ACID (UNII: O414PZ4LPZ)				
SODIUM SALICYLATE (UNII: WIQ1H85SYP)				
SORBIC ACID (UNII: X045WJ989B)				
METHYLCELLULOSE (1500 CPS) (UNII: P0NTE48364)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62011-0127-1	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/31/2014	12/30/2021
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC MONOGRAPH FINAL	part335	12/31/2014	12/30/2021	

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 12/2019

Strategic Sourcing Services LLC