

**IVERMECTIN- ivermectin paste**  
**Stellarnet Limited**

---

**PawBliss**

**Anthelmintic and Boticide**  
**Ivermectin Horse Paste 1.87**

**For Oral Use in Horses Only**

**IVERMECTIN**

(*ivermectin paste*) 1.87%

Net Wt. 0.21 oz (6.08 g)

Contents treat up to 1,250 lb body weight

Approved by FDA under ANADA # 200-326

Apple Flavored

*Removes worms and bots with a single dose.*

**INDICATIONS:** Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Ivermectin paste provides effective treatment and control of the following parasites in horses: large strongyles (adults), including *Strongylus vulgaris* (including early forms in blood vessels), *S. edentatus* (including tissue stages), *S. equinus*, *Triodontophorus* spp. including *T. brevicauda* and *T. serratus*, and *Craterostomum acuticaudatum*; small strongyles (adults, including those resistant to some benzimidazole-class compounds), including *Coronocyclus* spp. (*C. coronatus*, *C. labratus*, and *C. labiatus*), *Cyathostomum* spp. (*C. catinatum* and *C. pateratum*), *Cylicocyclus* spp. (*C. insignis*, *C. leptostomum*, *C. nassatus*, and *C. brevicapsulatus*), *Cylicodontophorus* spp., *Cylicostephanus* spp. (*C. calicatus*, *C. goldi*, *C. longibursatus*, and *C. minutus*), and *Poteriostomum* spp.; small strongyles (fourth-stage larvae); pinworms (adults and fourth-stage larvae), *Oxyuris equi*; ascarids (adults and third- and fourth-stage larvae), *Parascaris equorum*; hairworms (adults), *Trichostrongylus axei*; large-mouthed stomach worms (adults), *Habronema muscae*; bots (oral and gastric stages), *Gasterophilus* spp. including *G. intestinalis* and *G. nasalis*; lungworms (adults and fourth-stage larvae), *Dictyocaulus arnfieldi*; intestinal threadworms (adults), *Strongyloides westeri*; summer sores caused by *Habronema* and *Draschia* spp. (cutaneous third-stage larvae); and dermatitis caused by neck threadworm microfilariae (*Onchocerca* spp.).

**DOSAGE:** This syringe contains sufficient paste to treat one 1,250 lb horse at the recommended dose rate of 91 mcg ivermectin per lb (200 mcg/kg) of body weight. Each weight marking on the syringe plunger delivers sufficient paste to treat 250 lb of body weight. Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment and may encourage the development of parasite resistance.

**PARASITE CONTROL PROGRAM:** All horses should be included in a regular parasite control program, with particular attention given to mares, foals, and yearlings. Foals should be treated initially at 6 to 8 weeks of age, with routine treatments repeated as appropriate. Consult your veterinarian to develop a parasite control program that meets

your specific needs. Ivermectin paste effectively controls gastrointestinal nematodes and bots of horses. Regular treatment reduces the likelihood of verminous arteritis caused by *Strongylus vulgaris*.

**PRODUCT ADVANTAGES:** Broad-spectrum control — ivermectin paste eliminates important internal parasites of horses, including bots and the arterial stages of *Strongylus vulgaris*, with a single dose. Ivermectin is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

**ANIMAL SAFETY:** Ivermectin paste may be used in horses of all ages, including mares at any stage of pregnancy. Stallions may be treated without adversely affecting fertility.

**STORAGE INFORMATION:** Store at 68°F to 77°F (20°C to 25°C). Excursions between 59°F and 86°F (15°C to 30°C) are permitted.

**ADMINISTRATION:**

1. While holding the plunger, turn the knurled ring on the plunger one-quarter (1/4) turn to the left and slide it to the position corresponding to the prescribed weight, aligning the arrow on the ring with the line between the weight marking and "lb," as shown in the pictogram.
2. Lock the ring in place by making a one-quarter (1/4) turn to the right. Ensure that it is locked and no longer slides.
3. Ensure that the horse's mouth contains no feed.
4. Remove the cover from the tip of the syringe.
5. Insert the syringe tip into the horse's mouth at the space between the teeth.
6. Depress the plunger as far as it will go, depositing the paste on the back of the tongue.
7. Immediately raise the horse's head for a few seconds after dosing.



**WARNING:** Do not use in horses intended for human consumption. **Not for use in humans. Keep this and all drugs out of the reach of children.** Refrain from smoking or eating when handling this product. Wash hands after use. Avoid contact with eyes.

**PRECAUTIONS:** Ivermectin paste has been formulated specifically for use in horses only. This product should not be used in other animal species, as severe adverse reactions, including fatalities in dogs, may result.

**OTHER WARNINGS:** Parasite resistance may develop to any dewormer. Resistance has been reported for most classes of dewormers. Frequent treatment with a dewormer, used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated, may slow the development of parasite resistance. Clinical examinations, diagnostic tests, and parasite management history should be used to determine whether this product is appropriate for the herd prior to the use of any dewormer. Following the use of any dewormer, the effectiveness of treatment should be monitored (for example, through fecal egg count reduction tests or other appropriate methods). A decrease in a drug's effectiveness over time, as determined by fecal egg count reduction tests, may indicate the development of resistance to the dewormer administered. Parasite management programs should be adjusted accordingly based on regular monitoring.

**ENVIRONMENTAL SAFETY:** Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

**INFORMATION FOR HORSE OWNERS:** Swelling and itching reactions after treatment with ivermectin paste have occurred in horses carrying heavy infections of neck threadworm (*Onchocerca* spp.) microfilariae. These reactions are most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable. Consult your veterinarian should any such reactions occur. Healing of summer sores involving extensive tissue changes may require additional appropriate therapy in conjunction with treatment with ivermectin paste. Reinfection and measures for its prevention should also be considered. Consult your veterinarian if the condition does not improve.

**Distributed by:**

Y.L. Associates, Incorporated

9659 Limar Way  
San Diego, CA 92129-3508  
United States

**PRINCIPAL DISPLAY PANEL - 0.21 oz (6.08 g)**

Keep out of the reach of children.

For Oral Use in Horses Only


NDC: 86249-001-01

IVERMECTIN  
(ivermectin paste) 1.87%

Net Wt. 0.21 oz (6.08 g)  
Contents treat up to 1,250 lb body weight

Approved by FDA under ANADA # 200-326

Apple-Flavored



**Manufactured by:** Oscar Pet Co., Ltd.  
Building 3, No. 10, Fengxiang Middle Road, the Pearl River Village,  
Jianggao Town, Balyun District, Guangzhou, Guangdong, 510000, China(CHN)  
**Distributed by:** YI-Associates, Incorporated  
9659 Miramar Way, SAN DIEGO, CA 92129-3508, United States

**WARNING: Do not use in horses intended for human consumption.**  
Not for use in humans. Keep this and all drugs out of reach of children. Refrain from smoking and eating when handling. Wash hands after use. Avoid contact with eyes.

specifically for use in horses only. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

**OTHER WARNINGS:** Parasite resistance may develop to any dewormer.

## For Oral Use Horses Only

Removes worms and bots with a single dose  
Keep out of reach of children

# PawBliss

Apple Flavored!

Apple Flavored!

# IVERMECTIN (ivermectin PASTE 1.87%)

**ADMINISTRATION:**

- 1. While holding plunger, turn the knurled ring on the plunger 1/4 turn to the left and slide it to the side, then the barrel as at the prescribed weight, aligning the arrow on the ring to the line between the weight and lbs, as shown in the pictogram.
- 2. Lock the ring in place by making a 1/4 turn to the right. Ensure it is locked (it should not turn).
- 3. Make sure the plunger is fully compressed before use.
- 4. Remove the cover from the tip of the syringe.
- 5. Insert the syringe tip into the horse's mouth at the space between the upper and lower teeth.
- 6. Depress the plunger as far as it will go, depositing powder on the back of the tongue.
- 7. Immediately raise the horse's head for a few seconds after dosing.

**INFORMATION FOR HORSE OWNERS:** Swelling and itching reactions after treatment with *Pawlik's* (ivermectin paste) have occurred in horses carrying heavy infestations of neck headlureworm (*Oncophora sp.*) microfilariae. These reactions were most likely the result of microfilariae dying in large numbers in the blood stream. It is not known whether the manufacturer should any such reactions occur. Hearing of summer sales involving extensive tissue changes may require other appropriate therapy in conjunction with treatment with *Pawlik's*. Reinfestation, and measures for its prevention should also be considered. Consult your veterinarian if the condition does not improve.

## IVERMECTIN

## ivermectin paste

## Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:86249-001
<b>Route of Administration</b>	ORAL		

## **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>IVERMECTIN</b> (UNII: 8883YP2R6D) (IVERMECTIN - UNII:8883YP2R6D)	IVERMECTIN	18.7 mg in 1 g

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	APPLE	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:86249-001-01	6.08 g in 1 SYRINGE, PLASTIC		
2	NDC:86249-001-02	6.08 g in 1 SYRINGE, PLASTIC		
3	NDC:86249-001-03	6.08 g in 1 SYRINGE, PLASTIC		
4	NDC:86249-001-04	6.08 g in 1 SYRINGE, PLASTIC		
5	NDC:86249-001-05	6.08 g in 1 SYRINGE, PLASTIC		
6	NDC:86249-001-06	6.08 g in 1 SYRINGE, PLASTIC		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200326	01/12/2026	

**Labeler** - Stellarnet Limited (773255011)

**Registrant** - Stellarnet Limited (773255011)

## Establishment

Name	Address	ID/FEI	Business Operations
Oscar (Guangzhou) Pet Co., Ltd.		569802289	api manufacture

Revised: 1/2026

Stellarnet Limited