

**NON-DROWSY SINUS SEVERE, CIRCLE K- acetaminophen, guaifenesin, phenylephrine hcl caplet, film-coated tablet, film coated**  
**NON-DROWSY SINUS SEVERE, CIRCLE K- acetaminophen, guaifenesin, phenylephrine hcl caplet, film-coated tablet, film coated**  
**Lil' Drug Store products, Inc.**

-----  
**Non-Drowsy Sinus Severe, Circle K™**

## ***Drug Facts***

### **Active ingredients**

#### ***Active ingredients (in each caplet)***

Acetaminophen 325 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

### **Purpose**

#### ***Purpose***

Pain reliever/fever reducer

Expectorant

Nasal decongestant

### **Uses**

#### ***Uses***

- temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:
- headache
- nasal congestion
- minor aches and pains
- sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

### **Warnings**

#### **Liver warning**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

## **Allergy alert**

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

## **Do not use**

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

## **Ask a doctor**

### **Ask a doctor before use if you have**

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- heart disease
- thyroid disease
- diabetes
- liver disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**When using this product do not exceed recommended dosage.**

## **Stop use**

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

#### ***Directions***

- **do not take more than directed**
- adults and children 12 years and over
- take 2 caplets every 4 hours
- swallow whole - do not crush, chew, or dissolve
- do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

### ***Other information***

#### ***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

***Inactive ingredients*** corn starch, crospovidone, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

### ***Questions or comments***

***Questions or comments?*** 1-877-507-6516 (M-F 8AM-4:30PM CST)

**PDP/Package 6-count, NDC 66715-5632**

**TEMPORARILY RELIEVES SINUS HEADACHE & PRESSURE**

Compare to the active ingredients in

**Tylenol® SINUS SEVERE\***

**CIRCLE K™**

**non-drowsy**

**sinus severe**

**Acetaminophen,**

**Guaifenesin, Phenylephrine HCl**

**Pain reliever/Fever Reducer**

# Expectorant, Nasal Decongestant

[caplet image]

Actual Size

6 caplets



PDP/Package 24-count, NDC 66715-5544

## TEMPORARILY RELIEVES SINUS HEADACHE & PRESSURE

Compare to the active ingredients in  
**Tylenol® SINUS SEVERE\***

**CIRCLE K™**

**non-drowsy**

**sinus severe**

**Acetaminophen,**

**Guaifenesin, Phenylephrine HCl**

**Pain reliever/Fever Reducer**

**Expectorant,**

**Nasal Decongestant**

[caplet image]

Actual Size

24 caplets

**TEMPORARILY RELIEVES SINUS HEADACHE & PRESSURE**  
Compare to the active ingredients in  
**Tylenol® SINUS SEVERE®**

**CIRCLE K™**  
**non-drowsy**  
**sinus severe**

**Acetaminophen,  
Guaifenesin, Phenylephrine HCl  
Pain Reliever/Fever Reducer,  
Expectorant,  
Nasal Decongestant**



**24 caplets**

**KEEP OUTER PACKAGE FOR  
COMPLETE PRODUCT INFORMATION**

**Drug Facts**

**Active Ingredients (in each caplet)**

Acetaminophen 325 mg ..... Pain reliever/fever reducer  
Guaifenesin 200 mg ..... Expectorant  
Phenylephrine HCl 5 mg ..... Nasal decongestant

**Purpose**

**Uses** Temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:  
■ headache ■ nasal congestion ■ sore throat and pain ■ sinus congestion and pressure  
■ helps loosen phlegm (mucus) and thin bronchial secretions to make a coughs more productive

**Drug Facts (continued)**

**Warnings** Temporarily reduces fever  
■ **Warnings:** This product contains acetaminophen. Severe liver damage may occur if you take  
■ more than 4,000 mg of acetaminophen in 24 hours  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ hives ■ rash ■ skin redness  
If a skin reaction occurs, stop use and seek medical help right away.  
**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist  
■ if you are now taking a prescription nonsteroidal anti-inflammatory drug (NSAID) (aspirin, ibuprofen, naproxen, celecoxib, valdecoxib, or other NSAIDs), or for 2 weeks after stopping the NSAID drug, if you do not know if your prescription drug contains an NSAID, ask a doctor or pharmacist before taking this product.  
■ if you have ever had an allergic reaction to this product or any of its ingredients  
**Ask a doctor before use if you have**  
■ cough that occurs with too much phlegm (mucus)  
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema  
■ heart disease ■ thyroid disease ■ diabetes  
■ liver disease ■ high blood pressure

**Drug Facts (continued)**

**Warnings**

**Directions** ■ do not take more than directed  
■ adults and children 12 years and over  
■ take 2 caplets every 4 hours  
■ swallow whole - do not crush, chew, or dissolve  
■ do not take more than 10 caplets in 24 hours  
■ children under 12 years: ask a doctor  
**Other information**  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Drug Facts (continued)**

**Directions**

**Warnings** ■ difficulty in urination due to enlargement of the prostate gland  
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.  
When using this product do not exceed recommended dosage.  
Stop use and ask a doctor if  
■ nervousness, dizziness, or sleeplessness occur  
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present ■ new symptoms occur  
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.  
If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Warnings** ■ difficulty in urination due to enlargement of the prostate gland  
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.  
When using this product do not exceed recommended dosage.  
Stop use and ask a doctor if  
■ nervousness, dizziness, or sleeplessness occur  
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present ■ new symptoms occur  
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.  
If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

**Warnings** ■ difficulty in urination due to enlargement of the prostate gland  
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.  
When using this product do not exceed recommended dosage.  
Stop use and ask a doctor if  
■ nervousness, dizziness, or sleeplessness occur  
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present ■ new symptoms occur  
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.  
If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

**Warnings** ■ difficulty in urination due to enlargement of the prostate gland  
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.  
When using this product do not exceed recommended dosage.  
Stop use and ask a doctor if  
■ nervousness, dizziness, or sleeplessness occur  
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present ■ new symptoms occur  
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.  
If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

**Warnings** ■ difficulty in urination due to enlargement of the prostate gland  
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.  
When using this product do not exceed recommended dosage.  
Stop use and ask a doctor if  
■ nervousness, dizziness, or sleeplessness occur  
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present ■ new symptoms occur  
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.  
If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Product manufactured for:  
Lii Drug Store Products, Inc.  
9300 Earhart Lane SW  
Cedar Rapids, IA 52404  
Proudly distributed by  
Circle K Stores, Inc.



8-1108-527C-08-SS  
ORG0820C527XX  
55449C-CK-03-22

**NON-DROWSY SINUS SEVERE, CIRCLE K**

acetaminophen, guaifenesin, phenylephrine hcl caplet, film-coated tablet, film coated

<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-5632
<b>Route of Administration</b>	ORAL		

<b>Active Ingredient/Active Moiety</b>		<b>Basis of Strength</b>	<b>Strength</b>
<b>Ingredient Name</b>			
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN		200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1W5297W6MV)	PHENYLEPHRINE HYDROCHLORIDE		5 mg
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN		325 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>CROSPROVIDONE</b> (UNII: 2S7830E561)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>	MINT	<b>Imprint Code</b>	44;527
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5632-3	3 in 1 CARTON	08/01/2022	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/01/2022	

## NON-DROWSY SINUS SEVERE, CIRCLE K

acetaminophen, guaifenesin, phenylephrine hcl caplet, film-coated tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-5544
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE</b> (UNII: 2S7830E561)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>	MINT	<b>Imprint Code</b>	44;527
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5544-9	12 in 1 CARTON	08/01/2022	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/01/2022	

**Labeler** - Lil' Drug Store products, Inc. (093103646)