DAY TIME COLD/FLU RELIEF- acetaminohpen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled OUR FAMILY (Nash Finch Company)

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - fever
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

• liver disease

- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not exceed recommended dosage.

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose working: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other DAY TIME or NITE TIME products, carefully read each label to insure correct dosing

Other information

- store at room temperature 15°-30°C (59°-86°F)
- avoid excessive heat

Inactive ingredients

butylate hydroxyanisole, butylated hydroxytoluene, edible white ink, FD&C red #40*, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

*may contain this ingredient

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the Active Ingredients in Vicks® DayQuil® LiquiCaps®†

Pain Reliever/Fever Reducer

Cough Suppressant

Nasal Decongestant

ACETAMINOPHEN

Dextromethorphan HBr

Phenylephrine HCl

Multi-Symptom Cold/Flu Relief

DAY TIME LIQUID-CAPS

NON-DROWSY

LIQUID-CAPS

Does not contain Pseudoephedrine

[†]This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Vicks® DayQuil® LiquiCaps®.

DISTRIBUTED BY

NASH FINCH COMPANY ©2004, 1996

NFC BRANDS


7600 FRANCE AVE S, MPLS, MN 55435

www.ourfamilyfoods.com

TAMPER-EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Package Label



(hounitroo) stars mun	(heunitroo) stars mun
Active ingredients (in each softgel) Purpose Active ingredients (in each softgel) Pain releventievententieventententeeventieventeeventieventeevententeeventeeventieve	USes minor actives accommon cold and flu symptoms: minor actives and paires minor actives and poince in tratation minor actives minor ac
staa Facto	Drug Facts (continued)

Our Family

Quality Care Since 1904

R

Compare to the Active Ingredients in Vicks® DayQuil® LiquiCaps®†

Lot No.: Exp. Date:

tThis product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Vicks® DayQuil® LiquiCaps®,	When using this product, do not exceed recommended dosage. Stop use and ask a dottor if m pair, cough, or nasal congestion gets worse or lasts more than 7 days m ene symptoms occur mervousness, disciness, or sleeplessness occur effeven gets worse mervousness, disciness, or sleeplessness occur mereant mervousness, disciness, or serious condition. These could be signs of a serious condition.
Questions or comments? Call 5:753-3935 Monday-Friday 9AM-5PM EST	Ask a doctor or pharmacist before use if you are taking the blood thinning drug wattarin.
Inactive ingredients burylsted hydroxyanisole, burylsted hydroxytoluene, edible white ink, FD&C red #40*, FD&C yellow #6, gelstin, glycerin, polyethylene glycol, povidone, propylene glycol, purfied water, sorbitan, sorbitol *may contain this ingredient	Ask a doctor before use if you have inverdence on flop to the doctor before use if you have inverdence use if you have inverdence of the doctor betasting and thread due to an enlarged prostate gland in persistent or entronic coupt such as occurs with annotation, astimma, or emphysema couprimation of the doctor such and to be doctor as the doctor such and the doctor such and the doctor such and the doctor such as occurs with the doctor such as the
Other information = store at room temperature 15°-30°C (59°-86°F) = avoid excessive heat	conditions, or Parkinson's disease), or ior 2 weeks after stopping the MADI drug. If you do not know if your prescription drug contains an MADI, ask a doctor or pharmacist before taking this product.
UTERCIONS and not take more than directed (see Overdose varianting) and on take more than 4 doses in 24 hours warming) and or take more than 4 doses in 24 hours and children 12 years and over take 2 softgels with watter every a dhours and children under 12 years; do not use a when using other DAY TIME or NITE TIME products, carefully read each label to insure correct dosing the insure correct dosing the material dosing the correct dosing the correct dosing the material dosing the correct dosing the	vomiting, consult a doctor promptly. Do not use m with any other drug containing acetaminophen (prescription or nonpreacription). If you are not are whether a drug contains acetaminophen, ask a doctor or pharmacist. m f you are not wishing a prescription moramine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional (MAOI) (certain drugs for depression, psychiatric or emotional
Drug Facts (continued) If pregnant or breast-leeding, ask a health professional before use. Meep out of reast of hildren. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right avay. Quick medical attention is critical for equits as well as for children even if you do not notice any signs or symptoms. Directions medical attention is critical tor by the set of overdose and the recommended dose can by the set of overdose and the recent of the set of the set of the set of overdose and the set of overdose and signs or symptoms.	Drug Facts (continued) Warnings Liver warming. This product contains acetaminophen. Severe liver damage may occur if you take: more alcoholic drinks every day while using this product all or none alcoholic drinks every day while using this product Sore throat warming. If sore throat is severe, persists for more than 2 Sore throat warming. If sore throat is severe, persists for more than 2 Sore throat warming. If sore throat is severe, persists for more than 2 Sore throat warming. If sore throat is severe, persists for more than 2 Sore throat warming. If sore throat is severe, persists for more than 2 Sore throat warming of the severe persists for more than 2 Sore throat of the sectometer of the severe persists for more than 2 Sore throat of the sectometer of the
(heunitroo) stars any	(bengitno2) store min

Our Family Day Time Cold/Flu Liquid-Caps

Pain Reliever/Fever Reducer

Vmh

Cough Suppressant Nasal Decongestant

Dextromethorphan HBr Phenylephrine HCI

CETAMINOPHE

Product Informatio	n					
Product Type		OTC DRUG	Item Code (Source)	NDC:7025	3-470
				oource)		5 17 5
Route of Administratio	n ORAL					
Active Ingredient/A	ctive Moiety					
	Ingredient N	Name		Basis	of Strength	Strengt
ACETAMINO PHEN (UNII	•		NII:362O9ITL9D)	ACETAMINO	•	325 mg
DEXTROMETHORPHAN HYDRO BRO MIDE (UNII: 9 D2RTI9 KYH)DEXTROMETHORPHAN - UNII:7355X3ROTS)DEXTROMETHORPHAN - UNII:7355X3ROTS)						10 mg
PHENYLEPHRINE HYDR UNII:1WS297W6MV)	O CHLO RIDE (UNII: 0	4JA59TNSJ) (PHE	NYLEPHRINE -	PHENYLEPHE HYDROCHLO		5 mg
Inactive Ingredient	S					
Ingredient Name					S	trength
BUTYLATED HYDRO XY	ANISOLE (UNII: REK	4960K2U)				
BUTYLATED HYDRO XY	TOLUENE (UNII: 1P9	D0Z171K)				
FD&C RED NO.40 (UNII	WZB9127XOA)					
FD&C YELLOW NO.6 (UNII: H77VEI93A8)					
GELATIN (UNII: 2G86QN	327L)					
POLYETHYLENE GLYC	OLS (UNII: 3WJQ0SD	W1A)				
POVIDONES (UNII: FZ98	9 GH9 4E)					
PROPYLENE GLYCOL (UNII: 6 DC9 Q 16 7 V3)					
SORBITAN (UNII: 60921	CV9RU)					
SORBITOL (UNII: 506T6	0A25R)					
GLYCERIN (UNII: PDC6A	.3C0OX)					
WATER (UNII: 059QF0KC	00R)					
Product Characteri	stics					
Color	ORANGE	Sco	ore		no score	
Shape	CAPSULE		Size		19 mm	
Flavor		Imprint Code			P19;95A	
Contains		2111			.,	
contains						
Packaging						
# Item Code	Package De	scription	Marketing Sta	art Date	Marketing H	End Date
1 NDC:70253-470-16	2 in 1 CARTON					

Marketing Information							
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				
OTC MONOGRAPH FINAL	part341	0 5/3 1/20 13					

Labeler - OUR FAMILY (Nash Finch Company) (006962294)

Revised: 6/2013

OUR FAMILY (Nash Finch Company)