

MEIJER MULTI-SYMPTOM RELIEF- polyethylene glycol 400, tetrahydrozoline hcl, zinc sulfate solution/ drops
Meijer, Inc.

Meijer Multi-Symptom Relief (PLD)

Active ingredients

Polyethylene glycol 400....1%

Tetrahydrozoline HCl0.05%

Zinc sulfate.....0.25%

Purposes

Polyethylene glycol 400Lubricant

Tetrahydrozoline HClRedness reliever

Zinc sulfateAstringent

Uses

- Relieves dryness of the eye
- for temporary relief of discomfort and redness of the eye due to minor eye irritations
- for temporary relief of burning and irritation due to exposure to wind or sun
- for protection against further irritation

Warnings

For external use only

Ask a doctor before use if you have narrow angle glaucoma

When using this product

- pupils may become enlarged temporarily
- overuse may cause more eye redness
- remove contact lenses before using
- do not touch lip of container to any surface to avoid contamination
- replace cap after using
- do not use if this solution changes color or becomes cloudy

Stop use and ask a doctor if

- you feel eye pain
- changes in vision occur
- redness or irritation of the eye lasts
- condition worsens or lasts more than 72 hours

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- instill 1 to 2 drops in the affected eye(s) up to 4 times daily

- children under 6 years of age: ask a doctor

Other information

- some users may experience a brief tingling sensation
- store at 20°-25°C (68°-77°F)

Inactive ingredients

benzalkonium chloride, boric acid, edetate disodium, glycerin, hypromellose, purified water, sodium chloride, sodium citrate

Questions or comments?

Call 1-888-527-4276

meijer multi-symptom relief
Lubricant/Astringent/Redness Reliever Eye Drops

Drug Facts
Active ingredients
Polyethylene glycol 400 1%.....Lubricant
Tetrahydrozoline HCl 0.05%.....Redness reliever
Zinc sulfate 0.25%.....Astringent
Purposes

Uses
■ relieves dryness of the eye
■ for temporary relief of discomfort and redness of the eye due to minor eye irritations
■ for temporary relief of burning and irritation due to exposure to wind or sun
■ for protection against further irritation

Warnings
For external use only
Ask a doctor before use if you have narrow angle glaucoma
When using this product ■ pupils may become enlarged temporarily ■ overuse may cause more eye redness
■ remove contact lenses before using
■ do not touch tip of container to any surface to avoid contamination
■ replace cap after using
■ do not use if this solution changes color or becomes cloudy
Stop use and ask a doctor if ■ you feel eye pain
■ changes in vision occur ■ redness or irritation of the eye lasts
■ condition worsens or lasts more than 72 hours
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
■ instill 1 to 2 drops in the affected eye(s) up to 4 times daily
■ children under 6 years of age: ask a doctor

Drug Facts (continued)
Other information
■ some users may experience a brief tingling sensation
■ store at 20°- 25°C (68°- 77°F)
Inactive ingredients
benzalkonium chloride, boric acid, edetate disodium, glycerin, hypromellose, purified water, sodium chloride, sodium citrate
Questions or comments?
Call 1-888-527-4276

meijer Compare to Visine® Red Eye Total Comfort Multi-Symptom active ingredients*
NDC 79481-0250-1
DIST. BY MEIJER DISTRIBUTION, INC. GRAND RAPIDS, MI 49544 www.meijer.com
OUR QUALITY GUARANTEE The Meijer Family WWW.MEIJER.COM/SATISFACTION
PAPER BOX PLASTIC BOTTLE
how2recycle.info
7 13733 20139 1
CEDVT00151MEC
PD 5159810
0.5 FL OZ (15 mL)
Relieves red, burning, watery, itchy, gritty, dry, irritated eyes
STERILE
Lubricant/Astringent/Redness Reliever Eye Drops

TAMPER EVIDENT: DO NOT USE IF IMPRINTED NECKBAND IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

MEIJER MULTI-SYMPATOM RELIEF

polyethylene glycol 400, tetrahydrozoline hcl, zinc sulfate solution/ drops

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-0250
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TETRAHYDROZOLINE HYDROCHLORIDE (UNII: 0YZT43HS7D) (TETRAHYDROZOLINE - UNII:S9U025Y077)	TETRAHYDROZOLINE HYDROCHLORIDE	0.05 g in 100 mL
ZINC SULFATE (UNII: 89DS0H96TB) (ZINC CATION - UNII:13S1S8SF37)	ZINC SULFATE	0.25 g in 100 mL
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ) (POLYETHYLENE GLYCOL 400 - UNII:B697894SGQ)	POLYETHYLENE GLYCOL 400	1 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
BORIC ACID (UNII: R57ZHV85D4)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0250-1	1 in 1 CARTON	01/09/2023	
1		15 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M018	01/09/2023	

Labeler - Meijer, Inc. (006959555)**Registrant** - K.C. Pharmaceuticals, Inc. (174450460)**Establishment**

Name	Address	ID/FEI	Business Operations
K.C. Pharmaceuticals, Inc.		174450460	manufacture(79481-0250) , pack(79481-0250) , label(79481-0250)