

**EZ2GO LAXATIVE BISACODYL- bisacodyl capsule, coated
Valley Medical Products, LLC**

ez2go Laxative Bisacodyl

Drug Facts

Active Ingredients (In each tablet)

Bisacodyl USB 5 mg

Purpose

Stimulant laxative

Uses

- For temporary relief of occasional constipation and irregularity
- This product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use:

if you cannot swallow without chewing

Ask a doctor before use if you have

- Stomach pain, nausea or vomiting
- A sudden change in bowel habits that lasts more than 2 weeks

When using this product

- Do not chew or crush tablet(s)
- Do not use within 1 hour after taking an antacid or milk
- you may have stomach discomfort, faintness & cramps

Stop use and ask a doctor if

- You have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition
- You need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- Take with a glass of water

Adults and Children 12 years and over	Take 1 to 3 tablets in a single dose daily
Children 6 to under 12 years	Take 1 tablet in a single daily dose
Children under 6 years	ask a doctor

Other information**SEE BLISTER FOR EXPIRATION AND LOT NUMBER DO NOT USE IF BLISTER IS TORN OR SHOWS SIGNS OF TAMPERING**

- Store at 25°C(77°F); excursions permitted between 15°-30°C(59°-86°F)
- Avoid excessive humidity
- See end flap for expiration date and lot number


Inactive Ingredients

acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, oron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polydetrose,propylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac glaze, simethicone, bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or Comments?

1-800-403-9832 or WWW.ez2go.com

Package Labeling:

**Drug Facts**

Active Ingredients (In each tablet) Purpose
Bisacodyl USP, 5 mg Stimulant laxative

Uses

- For temporary relief of occasional constipation and irregularity
- This product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use: if you cannot swallow without chewing

Ask a doctor before use if you have

- Stomach pain, nausea or vomiting
- A sudden change in bowel habits that lasts more than 2 weeks

When using this product

- Do not chew or crush tablet(s)
- Do not use within 1 hour after taking an antacid or milk
- You may have stomach discomfort, faintness & cramps

Stop use and ask a doctor if

- You have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition
- You need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away →

Drug Facts (continued)

Directions	*Take with a glass of water
Adults and Children 12 years and over	Take 1 to 3 tablets in a single dose daily
Children 6 to under 12 years	Take 1 tablet in a single daily dose
Children under 6 years	ask a doctor

Other information

SEE BLISTER FOR EXPIRATION AND LOT NUMBER
DO NOT USE IF BLISTER IS TORN OR SHOWS SIGNS OF TAMPERING

- Store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- Avoid excessive humidity

Inactive Ingredients

acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, oron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polydextrose, propylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac glaze, simethicone, bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or Comments?

1-800-403-9832 or
www.ez2go.com

Manufactured for: © Copyright & Distributed by Valley Medical Products, LLC, Jacksonville, FL 32244 USA. All rights reserved.

EZ2GO LAXATIVE BISACODYL

bisacodyl capsule, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76470-059
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYL BISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
AMMONIA (UNII: 5138Q19F1X)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CARNAUBA WAX (UNII: R12CBM0E1Z)	
STARCH, CORN (UNII: 08232NY3SJ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	

MAGNESIUM STEARATE (UNII: 70097M6I30)
METHYLPARABEN (UNII: A2I8C7HI9T)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
POLYVINYL ACETATE PHTHALATE (UNII: 58QVG85GW3)
POVIDONE (UNII: FZ989GH94E)
PROPYLPARABEN (UNII: Z8IX2SC1OH)
BICARBONATE ION (UNII: HN1ZRA3Q20)
STEARIC ACID (UNII: 4ELV7Z65AP)
SUCROSE (UNII: C151H8M554)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TRIACETIN (UNII: XHX3C3X673)
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	5mm
Flavor		Imprint Code	5
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76470-059-59	4 in 1 BLISTER PACK; Type 0: Not a Combination Product	12/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)(A)	12/01/2025	

Labeler - Valley Medical Products, LLC (969389407)

Registrant - Valley Medical Products, LLC (969389407)

Revised: 1/2026

Valley Medical Products, LLC