

HEB STOMACH RELIEF MAX STRENGTH- bismuth subsalicylate suspension HEB

HEB Max Strength Bismuth Subsalicylate Drug Facts

Active ingredient (in each 30 mL dose cup)

Bismuth subsalicylate 1050 mg

Purposes

Upset stomach reliever and antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away at (1-800-222-1222).

Directions

- shake well before use
- for accurate dosing use dose cup
- **adults and children 12 years and over:** 1 dose (30 mL) every hour as needed
- do not exceed 4 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- each 30 mL contains: **sodium 20 mg**
- **salicylate 437 mg**
- low sodium
- sugar free
- store at room temperature.
- protect from freezing
- avoid excessive heat (more than 104°F or 40°C).

Inactive ingredients

carboxymethylcellulose sodium, D&C Red #22, D&C Red #28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

Questions or comments?

1-866-467-2748

Principal Display Panel

Compare to Pepto-Bismol® Maximum Strength active ingredient

HEB ®

Stomach Relief

Max

Bismuth Subsalicylate 1050 mg

Upset Stomach Reliever/Antidiarrheal

Multi-Symptom Relief

Relieves:

- Nausea
- Heartburn
- Indigestion
- Upset Stomach
- Diarrhea

MADE WITH PRIDE & CARE FOR H-E-B ®

SAN ANTONIA, TX 78204

100% GUARANTEE promise, if you aren't completely pleased with this product, we'll be happy to replace it or refund your money. You have our word on it.

***This product is not manufactured or distributed by Procter & Gamble, distributor of Pepto-Bismol® Maximum Strength.**

Package Label 473 mL

<p>Drug Facts (continued)</p> <p>Directions ■ shake well before use</p> <ul style="list-style-type: none"> ■ only use dose cup provided ■ adults and children 12 years and over <ul style="list-style-type: none"> ■ 15 mL (1 dose) every 1/2 hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea. ■ 15 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (120 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea <p>Other information</p> <ul style="list-style-type: none"> ■ each 15 mL dose cup contains: potassium 7 mg, sodium 5 mg ■ salicylate 230 mg ■ low sodium ■ sugar free ■ store at room temperature ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C) <p>Inactive ingredients carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, gelatin gum, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum.</p> <p>Questions or comments? 1-866-467-2748</p> <p>MADE WITH PRIDE & CARE FOR H-E-B® SAN ANTONIO, TX 78204</p> <p>100% GUARANTEE promise If you aren't completely pleased with this product, we'll be happy to replace it or refund your money. You have our word on it.</p> <p>LOT: 70716HEBLR 33497-2407 UNVARNISHED AREA EXP: 0 4 1220 54787 6</p>	<p>Compare to Pepto-Bismol® Ultra Strength active ingredient*</p> <p>NDC 37808-770-16</p> <p>H-E-B</p> <p>TEXAS-SIZE PACK *16 FL OZ*</p> <h1 style="font-size: 2em;">Stomach Relief Ultra</h1> <p>Bismuth Subsalicylate, 525 mg / Upset Stomach Reliever/Antidiarrheal</p> <p>Multi-Symptom Relief</p> <p>Relief of:</p> <ul style="list-style-type: none"> • Nausea • Heartburn • Indigestion • Upset Stomach • Diarrhea <p>2X CONCENTRATED FORMULA**</p> <p>16 FL OZ (1 PT) 473 mL</p>	<p>Do not use if imprinted shrink band is missing or broken</p> <p>Drug Facts</p> <table border="1"> <thead> <tr> <th>Active ingredient (in each 15 mL dose)</th> <th>Purpose</th> </tr> </thead> <tbody> <tr> <td>Bismuth subsalicylate 525 mg</td> <td>Upset stomach reliever and antidiarrheal</td> </tr> </tbody> </table> <p>Uses relieves ■ traveler's diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness</p> <p>Warnings</p> <p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Contains salicylate. Do not take if you are</p> <ul style="list-style-type: none"> ■ allergic to salicylates (including aspirin) ■ taking other salicylate products <p>Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool</p> <p>Ask a doctor before use if you have ■ fever ■ mucus in the stool</p> <p>Ask a doctor or pharmacist before use if you are taking any drug for ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis</p> <p>When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.</p> <p>Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.</p> <p><small>*This product is not manufactured or distributed by Procter & Gamble, distributor of Pepto-Bismol® Ultra Strength. **vs Original Pepto-Bismol®</small></p>	Active ingredient (in each 15 mL dose)	Purpose	Bismuth subsalicylate 525 mg	Upset stomach reliever and antidiarrheal
Active ingredient (in each 15 mL dose)	Purpose					
Bismuth subsalicylate 525 mg	Upset stomach reliever and antidiarrheal					

Package Label 236 mL

<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.</p> <p>Directions ■ shake well before use</p> <ul style="list-style-type: none"> ■ for accurate dosing, use dose cup ■ adults and children 12 years and over: 1 dose (30 mL) every hour as needed ■ do not exceed 4 doses (120 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea <p>Other information</p> <ul style="list-style-type: none"> ■ each 30 mL contains: sodium 20 mg ■ salicylate 437 mg ■ low sodium ■ sugar free ■ store at room temperature ■ protect from freezing ■ avoid excessive heat (more than 104°F or 40°C) <p>Inactive ingredients carboxymethylcellulose sodium, D&C Red #22, D&C Red #28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum</p> <p>Questions or comments? 1-866-467-2748</p> <p>MADE WITH PRIDE & CARE FOR H-E-B® SAN ANTONIO, TX 78204 8586-1810</p> <p>LOT: UNVARNISHED AREA EXP: 70708HEBLR 0 4 1220 54789 0</p>	<p>Compare to Pepto-Bismol® Maximum Strength active ingredient*</p> <p>H-E-B</p> <h1 style="font-size: 2em;">Stomach Relief Max</h1> <p>Bismuth Subsalicylate 1050 mg / Upset Stomach Reliever/Antidiarrheal</p> <p>Multi-Symptom Relief</p> <p>Relief of:</p> <ul style="list-style-type: none"> • Nausea • Heartburn • Indigestion • Upset Stomach • Diarrhea <p>8 FL OZ (236 mL)</p>	<p>Do not use if imprinted shrink band is missing or broken</p> <p>Drug Facts</p> <table border="1"> <thead> <tr> <th>Active ingredient (in each 30 mL dose cup)</th> <th>Purposes</th> </tr> </thead> <tbody> <tr> <td>Bismuth subsalicylate 1050 mg</td> <td>Upset stomach reliever and antidiarrheal</td> </tr> </tbody> </table> <p>Uses relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching</p> <p>Warnings</p> <p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Contains salicylate. Do not take if you are</p> <ul style="list-style-type: none"> ■ allergic to salicylates (including aspirin) ■ taking other salicylate products <p>Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool</p> <p>Ask a doctor before use if you have ■ fever ■ mucus in the stool</p> <p>Ask a doctor or pharmacist before use if you are taking any drug for ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis</p> <p>When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.</p> <p>Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days</p> <p><small>* This product is not manufactured or distributed by Procter & Gamble, distributor of Pepto-Bismol® Maximum Strength.</small></p>	Active ingredient (in each 30 mL dose cup)	Purposes	Bismuth subsalicylate 1050 mg	Upset stomach reliever and antidiarrheal
Active ingredient (in each 30 mL dose cup)	Purposes					
Bismuth subsalicylate 1050 mg	Upset stomach reliever and antidiarrheal					

HEB STOMACH RELIEF MAX STRENGTH

bismuth subsalicylate suspension

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-770
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ, BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	1050 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 7671P0Y5NH)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
WATER (UNII: 059QF0KO0R)	
SALICYLIC ACID (UNII: O414PZ4LPZ)	
DIMETHICONE, UNSPECIFIED (UNII: 92RU3N3Y1O)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	pink (viscous)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-770-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/23/2019	
2	NDC:37808-770-08	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/23/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	04/23/2019	

Labeler - HEB (007924756)

