

CLINDAMYCIN PHOSPHATE- clindamycin phosphate aerosol, foam Bryant Ranch Prepack

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Clindamycin Phosphate Foam, 1% safely and effectively. See full prescribing information for Clindamycin Phosphate Foam, 1%.

Clindamycin Phosphate Foam, 1%

For Topical Use

Initial U.S. Approval: 1970

INDICATIONS AND USAGE

Clindamycin Phosphate Foam, 1% is a lincosamide product indicated for acne vulgaris in patients 12 years and older. (1)

DOSAGE AND ADMINISTRATION

- For topical use only; not for oral, ophthalmic, or intravaginal use. (2)
- Apply Clindamycin Phosphate Foam, 1% once daily to affected areas. (2)
- Flammable; avoid fire, flame and/or smoking during and immediately following application. (2)

DOSAGE FORMS AND STRENGTHS

Foam containing 1% clindamycin as clindamycin phosphate. (3)

CONTRAINDICATIONS

Clindamycin Phosphate Foam, 1% is contraindicated in individuals with a history of regional enteritis or ulcerative colitis, or a history of antibiotic-associated colitis, (including pseudomembranous colitis). (4)

WARNINGS AND PRECAUTIONS

- Colitis: Clindamycin can cause severe colitis, which may result in death. Diarrhea, bloody diarrhea, and colitis (including pseudomembranous colitis) have been reported with the use of clindamycin. Clindamycin Phosphate Foam, 1% should be discontinued if significant diarrhea occurs. (5.1)

ADVERSE REACTIONS

The most common adverse reactions (>1%) are headache and application site reactions including burning, pruritus, and dryness. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Perrigo at 1-866-634-9120 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 2/2022

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Colitis

5.2 Irritation

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

7.1 Erythromycin

7.2 Neuromuscular Blocking Agents

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

17.1 Instructions for Use

17.2 Skin Irritation

17.3 Colitis

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Clindamycin Phosphate Foam, 1% is indicated for topical application in the treatment of acne vulgaris in patients 12 years and older.

2 DOSAGE AND ADMINISTRATION

Clindamycin Phosphate Foam, 1% is for topical use only, and not for oral, ophthalmic or intravaginal use.

Apply Clindamycin Phosphate Foam, 1% once daily to affected areas after the skin is washed with mild soap and allowed to fully dry. Use enough to cover the entire affected area.

If there has been no improvement after 6 to 8 weeks or if the condition becomes worse, treatment should be discontinued.

The contents of Clindamycin Phosphate Foam, 1% are flammable; avoid fire, flame and/or smoking during and immediately following application.

3 DOSAGE FORMS AND STRENGTHS

White to off-white thermolabile foam. Each gram of Clindamycin Phosphate Foam, 1%

contains, as dispensed, 12 mg (1.2%) of clindamycin phosphate, equivalent to 10 mg (1%) of clindamycin.

4 CONTRAINDICATIONS

Clindamycin Phosphate Foam, 1% is contraindicated in individuals with a history of regional enteritis or ulcerative colitis, or a history of antibiotic-associated colitis (including pseudomembranous colitis).

5 WARNINGS AND PRECAUTIONS

5.1 Colitis

Systemic absorption of clindamycin has been demonstrated following topical use of this product. Diarrhea, bloody diarrhea, and colitis (including pseudomembranous colitis) have been reported with the use of topical clindamycin. If significant diarrhea occurs, Clindamycin Phosphate Foam, 1% should be discontinued. [*See Adverse Reactions*(6.2).]

Severe colitis has occurred following oral or parenteral administration of clindamycin with an onset of up to several weeks following cessation of therapy. Antiperistaltic agents such as opiates and diphenoxylate with atropine may prolong and/or worsen severe colitis. Severe colitis may result in death.

Studies indicate a toxin(s) produced by *Clostridia* is one primary cause of antibiotic-associated colitis. The colitis is usually characterized by severe persistent diarrhea and severe abdominal cramps and may be associated with the passage of blood and mucus. Stool cultures for *Clostridium difficile* and stool assay for *C. difficile* toxin may be helpful diagnostically.

5.2 Irritation

Clindamycin Phosphate Foam, 1% can cause irritation. Concomitant topical acne therapy should be used with caution since a possible cumulative irritancy effect may occur, especially with the use of peeling, desquamating, or abrasive agents. If irritation or dermatitis occurs, clindamycin should be discontinued.

Avoid contact of Clindamycin Phosphate Foam, 1% with eyes, mouth, lips, other mucous membranes or areas of broken skin. If contact occurs, rinse thoroughly with water.

Clindamycin Phosphate Foam, 1% should be prescribed with caution in atopic individuals.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

A total of 439 subjects with mild to moderate acne vulgaris were treated once daily for 12 weeks with clindamycin phosphate foam, 1%.

The incidence of adverse reactions occurring in $\geq 1\%$ of the subjects in clinical trials comparing clindamycin phosphate foam, 1% and its vehicle is presented in Table 1.

Table 1: Adverse Reactions Occurring in $\geq 1\%$ of Subjects

Adverse Reactions	Number (%) of Subjects	
	Clindamycin Phosphate Foam, 1% N = 439	Vehicle Foam N = 154
Headache	12 (3%)	1 (1%)
Application site burning	27 (6%)	14 (9%)
Application site pruritus	5 (1%)	5 (3%)
Application site dryness	4 (1%)	5 (3%)
Application site reaction, not otherwise specified	3 (1%)	4 (3%)

In a contact sensitization study, none of the 203 subjects developed evidence of allergic contact sensitization to clindamycin phosphate foam, 1%.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post approval use of clindamycin phosphate foam, 1%: application site pain, application site erythema, diarrhea, urticaria, abdominal pain, hypersensitivity, rash, abdominal discomfort, nausea, seborrhea, application site rash, dizziness, pain of skin, colitis (including pseudomembranous colitis), and hemorrhagic diarrhea. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Abdominal pain and gastrointestinal disturbances, as well as gram-negative folliculitis, have also been reported in association with the use of topical formulations of clindamycin. Orally and parenterally administered clindamycin have been associated with severe colitis, which may end fatally.

7 DRUG INTERACTIONS

7.1 Erythromycin

Clindamycin Phosphate Foam, 1% should not be used in combination with topical or oral erythromycin-containing products due to possible antagonism to its clindamycin component. *In vitro* studies have shown antagonism between these two antimicrobials. The clinical significance of this *in vitro* antagonism is not known.

7.2 Neuromuscular Blocking Agents

Clindamycin has been shown to have neuromuscular blocking properties that may enhance the action of other neuromuscular blocking agents. Therefore, Clindamycin Phosphate Foam, 1% should be used with caution in patients receiving such agents.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category B: There are no adequate and well-controlled studies in pregnant women treated with clindamycin phosphate foam, 1%. Clindamycin Phosphate Foam, 1% should be used during pregnancy only if the potential benefit clearly outweighs the potential risk to the fetus.

Reproduction studies have been performed in rats and mice using subcutaneous and oral doses of clindamycin phosphate, clindamycin hydrochloride and clindamycin palmitate hydrochloride. These studies revealed no evidence of fetal harm.

The highest dose used in the rat and mouse teratogenicity studies was equivalent to a clindamycin phosphate dose of 432 mg/kg. For a rat, this dose is 84 fold higher, and for a mouse 42 fold higher, than the anticipated human dose of clindamycin phosphate from clindamycin phosphate foam, 1% based on a mg/m² comparison.

8.3 Nursing Mothers

It is not known whether clindamycin is excreted in human milk following use of Clindamycin Phosphate Foam, 1%. However, orally and parenterally administered clindamycin has been reported to appear in breast milk. Because of the potential for serious adverse reactions in nursing infants, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

If used during lactation and Clindamycin Phosphate Foam, 1% is applied to the chest, care should be taken to avoid accidental ingestion by the infant.

8.4 Pediatric Use

Safety and effectiveness of clindamycin phosphate foam, 1% in children under the age of 12 have not been studied.

8.5 Geriatric Use


The clinical study with clindamycin phosphate foam, 1% did not include sufficient numbers of subjects aged 65 and over to determine if they respond differently than younger subjects.

11 DESCRIPTION

Clindamycin Phosphate Foam, 1% contains clindamycin (1%) as clindamycin phosphate.

Clindamycin phosphate is a water-soluble ester of the semi-synthetic antibiotic produced by a 7(S)-chloro-substitution of the 7(R)-hydroxyl group of the parent antibiotic, lincomycin.

The chemical name for clindamycin phosphate is methyl 7-chloro-6,7,8-trideoxy-6-(1-methyl- *trans*-4-propyl-L-2-pyrrolidinecarboxamido)1-thio-L- *threo*- α -D- *galacto*-octopyranoside 2-(dihydrogen phosphate). The structural formula for clindamycin phosphate is represented below:



Molecular Formula: $C_{18}H_{34}ClN_2O_8PS$

Molecular Weight: 504.97 g/mol

Clindamycin Phosphate Foam, 1% contains clindamycin (1%) as clindamycin phosphate, at a concentration equivalent to 10 mg clindamycin per gram in a thermolabile hydroethanolic foam vehicle consisting of cetyl alcohol, ethanol (58%), polysorbate 60, propylene glycol, purified water, and stearyl alcohol pressurized with a hydrocarbon (propane/butane) propellant.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Mechanism of action in acne vulgaris is unknown. [See *Microbiology* (12.4)]

12.2 Pharmacodynamics

Pharmacodynamics of Clindamycin Phosphate Foam, 1% is unknown.

12.3 Pharmacokinetics

In an open label, parallel group study in 24 subjects with acne vulgaris, 12 subjects (3 male and 9 female) applied 4 grams of clindamycin phosphate foam, 1% once-daily for five days, and 12 subjects (7 male and 5 female) applied 4 grams of a clindamycin gel, 1%, once daily for five days. On Day 5, the mean C_{max} and $AUC_{(0-12)}$ were 23% and 9% lower, respectively, for clindamycin phosphate foam, 1% than for the clindamycin gel, 1%.

Following multiple applications of clindamycin phosphate foam, 1%, less than 0.024% of the total dose was excreted unchanged in the urine over 12 hours on Day 5.

12.4 Microbiology

No microbiology studies were conducted in the clinical trials with this product.

Clindamycin binds to the 50S ribosomal subunits of susceptible bacteria and prevents elongation of peptide chains by interfering with peptidyl transfer, thereby suppressing protein synthesis. Clindamycin has been shown to have *in vitro* activity against *Propionibacterium acnes* (*P. acnes*), an organism that has been associated with acne vulgaris; however, the clinical significance of this activity against *P. acnes* was not examined in clinical studies with Clindamycin Phosphate Foam, 1%. *P. acnes* resistance to clindamycin has been documented.

Inducible Clindamycin Resistance

The treatment of acne with antimicrobials is associated with the development of antimicrobial resistance in *P. acnes* as well as other bacteria (e.g. *Staphylococcus aureus*, *Streptococcus pyogenes*). The use of clindamycin may result in developing inducible resistance in these organisms. This resistance is not detected by routine susceptibility testing.

Cross Resistance

Resistance to clindamycin is often associated with resistance to erythromycin.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

The carcinogenicity of a 1.2% clindamycin phosphate gel similar to Clindamycin Phosphate Foam, 1% was evaluated by daily application to mice for two years. The daily doses used in this study were approximately 3 and 15 times higher than the human dose of clindamycin phosphate from 5 milliliters of Clindamycin Phosphate Foam, 1%, assuming complete absorption and based on a body surface area comparison. No significant increase in tumors was noted in the treated animals.

A 1.2% clindamycin phosphate gel similar to Clindamycin Phosphate Foam, 1% caused a statistically significant shortening of the median time to tumor onset in a study in hairless mice in which tumors were induced by exposure to simulated sunlight.

Genotoxicity tests performed included a rat micronucleus test and an Ames Salmonella reversion test. Both tests were negative.

Reproduction studies in rats using oral doses of clindamycin hydrochloride and clindamycin palmitate hydrochloride have revealed no evidence of impaired fertility.

14 CLINICAL STUDIES

In one multicenter, randomized, double-blind, vehicle-controlled clinical trial, subjects with mild to moderate acne vulgaris used clindamycin phosphate foam, 1% or the vehicle Foam once daily for twelve weeks. Treatment response, defined as the proportion of subjects clear or almost clear, based on the Investigator Static Global Assessment (ISGA), and the mean percent reductions in lesion counts at the end of treatment in this study are shown in Table 2.

Table 2: Efficacy Results at Week 12

Efficacy Parameters	Clindamycin Phosphate Foam, 1% N = 386	Vehicle Foam N = 127
Treatment response (ISGA)	31%	18%*
Percent reduction in lesion counts		
Inflammatory Lesions	49%	35%*
Noninflammatory Lesions	38%	27%*

Total Lesions	43%	31%*
---------------	-----	------

*P<0.05

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Clindamycin Phosphate Foam, 1% containing clindamycin phosphate equivalent to 10 mg clindamycin per gram, is white to off-white in color and thermolabile.

It is available in the following sizes:

- 100 gram aerosol can - NDC 63629-8629-1

16.2 Storage and Handling

Store at 20-25°C (68-77°F) [see USP controlled room temperature].

Flammable. Avoid fire, flame or smoking during and immediately following application.

Contents under pressure. Do not puncture or incinerate. Do not expose to heat or store at temperature above 120°F (49°C).

Keep out of reach of children.

17 PATIENT COUNSELING INFORMATION

See FDA-Approved patient labeling (Patient Information).

17.1 Instructions for Use

- Patients should be advised to wash their skin with mild soap and allow it to dry before applying Clindamycin Phosphate Foam, 1%.
- Patients should use enough Clindamycin Phosphate Foam, 1% to cover the face and to apply once daily.
- Patients should dispense Clindamycin Phosphate Foam, 1% directly into the cap or onto a cool surface.

Patients should wash their hands after applying Clindamycin Phosphate Foam, 1%.

17.2 Skin Irritation

Clindamycin Phosphate Foam, 1% may cause irritation such as erythema, scaling, itching, burning, or stinging. Patients should be advised to discontinue use if excessive irritancy or dermatitis occur.

17.3 Colitis

In the event a patient treated with Clindamycin Phosphate Foam, 1% experiences severe diarrhea or gastrointestinal discomfort, Clindamycin Phosphate Foam, 1% should be discontinued and a physician should be contacted.

Made in Israel

Manufactured by Perrigo

Yeruham, Israel

Distributed by Perrigo

Allegan, MI 49010

www.perrigo.com

Rev 08-19

4T500 RC J4

Pharmacist-Detach here and Give Instructions to Patient

Patient Information

Clindamycin Phosphate Foam, 1%

Important: For skin use only. Do not use Clindamycin Phosphate Foam, 1% in your eyes, mouth or vagina.

Read the Patient Information that comes with Clindamycin Phosphate Foam, 1% before you start using it and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your doctor about your medical condition or treatment.

What is Clindamycin Phosphate Foam, 1%?

Clindamycin Phosphate Foam, 1% is a prescription medicine used on the skin (topical) to treat acne in people 12 years and older.

It is not known if Clindamycin Phosphate Foam, 1% is safe and effective in children under 12 years of age.

Who should not use Clindamycin Phosphate Foam, 1%?

Do not use Clindamycin Phosphate Foam, 1% if you:

- have Crohn's disease
- have ulcerative colitis
- have had inflammation of the colon (colitis) or severe diarrhea with past antibiotic use

Tell your doctor if you are not sure if you have **any** of the conditions **listed above**.

What should I tell my doctor before using Clindamycin Phosphate Foam, 1%? Before you use Clindamycin Phosphate Foam, 1%, tell your doctor if you:

- have a history of eczema
- are planning to have surgery. Clindamycin Phosphate Foam, 1% may affect how certain medicines work that may be given during surgery.
- have any other medical conditions
- are pregnant or planning to become pregnant. It is not known if Clindamycin Phosphate Foam, 1% may harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if Clindamycin Phosphate Foam, 1% passes through your breast milk.

You and your doctor should decide if you will use Clindamycin Phosphate Foam, 1% or breastfeed. If you use Clindamycin Phosphate Foam, 1% while breastfeeding and Clindamycin Phosphate Foam, 1% is applied on the chest, take care to avoid getting Clindamycin Phosphate Foam, 1% into your baby's mouth.

Tell your doctor about all the medicines you take including prescription and non-prescription medicines, vitamins and herbal supplements. Clindamycin Phosphate Foam, 1% may affect the way other medicines work and other medicines may affect how Clindamycin Phosphate Foam, 1% works.

Especially tell your doctor if you take erythromycin or use products on your skin that contain erythromycin.

Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

How should I use Clindamycin Phosphate Foam, 1%?

- Clindamycin Phosphate Foam, 1% is for use on the skin only. Do not get Clindamycin Phosphate Foam, 1% in your eyes, mouth or vagina.
- Use Clindamycin Phosphate Foam, 1% exactly as your doctor tells you to use it. See the "Instructions for Applying Clindamycin Phosphate Foam, 1%" below.
- Apply Clindamycin Phosphate Foam, 1% 1 time a day.
- Wash your skin with mild soap and water and dry before applying Clindamycin Phosphate Foam, 1%.
- Do not dispense Clindamycin Phosphate Foam, 1% directly onto your hands or face, because the foam will begin to melt on contact with warm skin.

Instructions for Applying Clindamycin Phosphate Foam, 1%.



Figure A: Remove clear cap.

- Remove the clear cap from the Clindamycin Phosphate Foam, 1% can (see Figure A).



Figure B: Hold can upright and press nozzle firmly to dispense.

- Hold the can upright and firmly press the nozzle to dispense Clindamycin Phosphate Foam, 1% (see Figure B).



Figure C: Dispense Clindamycin Phosphate Foam, 1% into the clear cap.

- Dispense Clindamycin Phosphate Foam, 1% into the clear cap or onto a cool surface (see Figure C).



Figure D: Dispense enough Clindamycin Phosphate Foam, 1% to cover affected area.

- Dispense enough Clindamycin Phosphate Foam, 1% to cover the affected area (see Figure D).



Figure E: If the can seems warm or the foam is runny, run the can under cold water.

- If the can seems warm or the foam seems runny, run the can under cold water (see Figure E).



Figure F: Pick up a small amount of Clindamycin Phosphate Foam, 1% on your fingertips and gently rub into the affected area until the foam disappears.

- Pick up small amounts of Clindamycin Phosphate Foam, 1% with your fingertips and gently rub the foam into the affected area until the foam disappears (see Figure F).

- Wash your hands after applying Clindamycin Phosphate Foam, 1%.

Throw away any of the unused medicine that you dispensed out of the can.

What should I avoid while using Clindamycin Phosphate Foam, 1%?

- Clindamycin Phosphate Foam, 1% is flammable. Avoid fire, flames, or smoking during and right after you apply Clindamycin Phosphate Foam, 1% to your skin.
- Avoid getting Clindamycin Phosphate Foam, 1% in or near your eyes, mouth, lips, or broken skin. If you get Clindamycin Phosphate Foam, 1% in your eyes, mouth, on lips or broken skin, rinse well with water.

What are possible side effects with Clindamycin Phosphate Foam, 1%?

Clindamycin Phosphate Foam, 1% can cause serious side effects including:

- Inflammation of the colon (colitis). Clindamycin can cause severe colitis that may lead to death. Stop using Clindamycin Phosphate Foam, 1% and call your doctor right away if you have severe watery diarrhea, or bloody diarrhea.

The most common side effects of Clindamycin Phosphate Foam, 1% include:

- Skin irritation. Clindamycin Phosphate Foam, 1% may cause skin irritation such as burning, itching, or dryness. Stop using Clindamycin Phosphate Foam, 1% and talk with your doctor if you develop excessive skin irritation.
- Headache.

Tell your doctor if you have any side effect that bothers you or that does not go away.

These are not all the side effects of Clindamycin Phosphate Foam, 1%. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may report side effects to Perrigo at 1-866-634-9120.

How should I store Clindamycin Phosphate Foam, 1%?

- Store Clindamycin Phosphate Foam, 1% at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep Clindamycin Phosphate Foam, 1% away from heat. Never throw the can into a fire, even if the can is empty.
- Do not store Clindamycin Phosphate Foam, 1% at temperatures above 120°F (49°C).
- Do not break through (puncture) the Clindamycin Phosphate Foam, 1% can.

Keep Clindamycin Phosphate Foam, 1% and all medicines out of the reach of children.

General information about the safe and effective use of Clindamycin Phosphate Foam, 1%:

Medicines are sometimes prescribed for purposes other than those listed in Patient Information. Do not use Clindamycin Phosphate Foam, 1% for a condition for which it was not prescribed. Do not give Clindamycin Phosphate Foam, 1% to other people, even if they have the same symptoms you have. It may harm them.

This leaflet summarizes the most important information about Clindamycin Phosphate Foam, 1%. If you would like more information, talk with your doctor. You can also ask your pharmacist or doctor for information about Clindamycin Phosphate Foam, 1% that is written for health professionals.

What are the ingredients in Clindamycin Phosphate Foam, 1%?

Active ingredient: clindamycin phosphate

Inactive ingredients: cetyl alcohol, ethanol (58%), polysorbate 60, propylene glycol, purified water, and stearyl alcohol. The can is pressurized with a hydrocarbon (propane/butane) propellant.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Made in Israel

Manufactured by Perrigo

Yeruham, Israel

Distributed by Perrigo

Allegan, MI 49010

www.perrigo.com

Rev 08-19

4T500 RC J4

Clindamycin Phosphate 1% Foam, #100



Each gram contains: Clindamycin Phosphate, at a concentration equivalent to 10 mg clindamycin per gram in a thermolabile hydroethanolic foam vehicle consisting of cetyl alcohol, ethanol (58%), polysorbate 60, propylene glycol, purified water, and stearyl alcohol pressurized with a hydrocarbon (propane/butane) propellant.

This package is not child resistant. Keep out of reach of children. Avoid contact with eyes.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Do not expose to heat or store at temperature above 120°F (49°C).

Warning: Flammable. Avoid fire, flame or smoking during and immediately following application.

For topical use only. Not for ophthalmic, oral or intravaginal use.

NDC 63629-8629-1

Clindamycin Phosphate Foam

1%



Relabeled by:
Bryant Ranch Prepack, Inc.
Burbank, CA 91504 USA

Rx only
100 g

Manufactured by:
Padagis



CLINDAMYCIN PHOSPHATE

clindamycin phosphate aerosol, foam

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:63629-8629(NDC:45802-660)
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CLINDAMYCIN PHOSPHATE (UNII: EH6D7113I8) (CLINDAMYCIN - UNII:3U02EL437C)	CLINDAMYCIN	10 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	

POLYSORBATE 60 (UNII: CAL22UVI4M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
STEARYL ALCOHOL (UNII: 2KR89I4H1Y)	
CETYL ALCOHOL (UNII: 936JST6JCN)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63629-8629-1	1 in 1 CARTON	03/31/2010	
1		100 g in 1 CAN; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090785	03/31/2010	

Labeler - Bryant Ranch Prepack (171714327)

Registrant - Bryant Ranch Prepack (171714327)

Establishment

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	repack(63629-8629) , relabel(63629-8629)

Revised: 7/2023

Bryant Ranch Prepack