

YUNNAN BAIYAO- camphor, menthol, and methyl salicylate spray
YUNNAN BAIYAO GROUP CO., LTD.

Drug Facts

Active ingredients

Camphor 3.5%

Menthol 4.5%

Methyl Salicylate 10%

Purpose

External analgesic

External analgesic

External analgesic

Uses

For the temporary relief of minor aches and pains of muscles and joints due to:
simple backache

arthritis

strains

bruises

sprains

Warnings

For external use only

Flammable: keep away from fire or flame

Do not use

on irritated or damaged skin

on wounds

otherwise than as directed

When using this product

avoid contact with the eyes or mucous membranes

do not bandage tightly

Stop use and ask a doctor if

condition worsens

symptoms persist for more than 7 days

symptoms clear up and occur again within a few days

excessive irritation of the skin develops

nausea, vomiting, abdominal discomfort, diarrhea, or skin rash occurs

when using for pain of arthritis:

pain persists for more than 10 days

redness is present

in conditions affecting children under 12 years of age

Keep out of reach of children to avoid accidental poisoning. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

shake well before using

adults and children 3 years of age and older: remove the attached film from the patch.

Apply to affected area not more than 3 to 4 times daily

children under 3 years of age: do not use, consult your physician refer to the above

warnings; use otherwise than as directed may be dangerous

Other information

keep tightly closed

store at 15 to 30 C (59 to 86 F)

Inactive ingredients

Alcohol, ampelopsis delavayana root, capsicum, Chinese yam rhizome, corydalis yanhusuo rhizome, dong-quai root, fragrant angelica root, Japanese dioscorea rhizome, san-qi ginseng root and rhizome, Sichuan lovage rhizome.

Questions or comments?(888) 221-3496 M-F 9 am to 5 pm

you may also report serious side effects to this phone number

Yunnan Baiyao SPRAY

External Analgesic

NDC 61658-006-01

1 fl oz (30 mL)

YNBY Spray 30mL - Unit Box
雲南白藥氣霧劑
Size: 35 (W) x 35 (D) x 105 (H) mm
03/19/2024



專色



SOLSTICE
MEDICINE COMPANY

YNBY[®]

EXTERNAL ANALGESIC
NDC 61658-006-01

Yunnan Baiyao
Spray

1 fl. oz. 液量盎司
(30 mL 毫升)

雲白藥國際有限公司
YUNNAN INTERNATIONAL LIMITED

Drug Facts (continued)

Other Information

- keep tightly closed
- store at 15° to 30°C (59° to 86°F)

Active Ingredients

Alcohol, ampelopis delavayana root, capsilum, Chinese yam rhizome, corydalis yanhusuo rhizome, dong-quai root, fragrant angelica root, Japanese Dioscorea rhizome, sea-of-ginseng root and rhizome, and Sichuan lovage rhizome

Questions or Comments?

- (888) 221-3496
- M-F, 9 am to 5 pm
- you may also report serious side effects to this phone number

U.S. Distributor 美國總代理:

SOLSTICE MEDICINE COMPANY
美國威氏藥業有限公司
215 W. Ann Street
Los Angeles, CA 90012
Tel: (888) 221-3496
www.solsticepharm.com

Manufactured by 雲南白藥(雲南)

雲南白藥 藥業有限公司
www.yunnanbaiyao.com.cn
No. 3688 Yunnan Baiyao Street,
Chengde District, Kunming, Yunnan
People's Republic of China

MADE IN CHINA
中國製造

SOLSTICE
MEDICINE COMPANY

YNBY[®]

EXTERNAL ANALGESIC
NDC 61658-006-01

Yunnan Baiyao
Spray

1 fl. oz. 液量盎司
(30 mL 毫升)

雲白藥國際有限公司
YUNNAN INTERNATIONAL LIMITED

Drug Facts

Active Ingredients	Purpose
Camphor 5.1%	External analgesic
Methyl Salicylate 4.0%	External analgesic
Methyl Salicylate 12%	External analgesic

Uses For the temporary relief of minor aches and pains of muscles and joints due to:
 ● simple backache ● arthritis ● strain ● bruise ● sprain

Warnings

For external use only

Flammable: keep away from fire or flame.

Do not use

- on irritated or damaged skin
- on wounds
- otherwise than as directed

When using this product:

- avoid contact with the eyes or mucous membranes
- do not bandage tightly

Stop use and ask a doctor if

- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days

When using for pain of arthritis:

- excessive irritation of the skin develops
- nausea, vomiting, abdominal discomfort, diarrhea, or skin rash occurs
- when using for pain of arthritis:
- pain persists for more than 10 days
- redness is present
- in conditions affecting children under 12 years of age

Keep out of reach of children to avoid accidental poisoning.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- shake well before using
- keep nozzle 2 to 4 inches from surface of skin
- adults and children 3 years of age and older: spray onto affected area not more than 3 to 4 times daily
- children under 3 years of age: do not use, consult your physician
- refer to the above warnings. Use otherwise than as directed may be dangerous.

Lot No.:

Exp. Date:




0 49987 01605 5

External Analgesic

NDC 61658-006-02

2.9 fl oz (85 mL)

YNBY Spray 85mL - Unit Box
雲南白藥氣霧劑
Size: 38 (W) x 38 (D) x 170 (H) mm
03/19/2024




YNBY[®]

**Yunnan Baiyao
Spray**

EXTERNAL ANALGESIC
NDC 61658-006-02

2.9 fl. oz. 液量盎司
(85 mL 毫升)



Drug Facts (continued)

Other Information

- keep tightly closed
- store at 15° to 30°C (59° to 85°F)

Inactive Ingredients

Alcohol, ampelopsis delavayana root, capillium, Chinese yam rhizome, corydalis yanhusuo rhizome, dong-quai root, fragrant angelica root, Japanese Dioscorea rhizome, sai-qí ginseng root and rhizome, and Sichuan lovage rhizome

Questions or Comments?

- (888) 221-3496
- M-F, 9 am to 5 pm
- you may also report serious side effects to this phone number




U.S. Distributor 美國總代理：
SOLSTICE MEDICINE COMPANY
 美國蘇沃森藥業有限公司
 215 W. Ann Street
 Los Angeles, CA 90012
 Toll Free: 1-888-221-3496
www.solsticemed.com

Manufactured by 雲南白藥集團股份有限公司
Yunnan Baiyao Group Co., Ltd.
 No. 99# Yunnan Baiyao Street
 Changqing District, Kunming, Yunnan
 People's Republic of China

MADE IN CHINA
中國製造

Lot No.:

Exp. Date:




YNBY[®]

**Yunnan Baiyao
Spray**

EXTERNAL ANALGESIC
NDC 61658-006-02

2.9 fl. oz. 液量盎司
(85 mL 毫升)



Drug Facts

Active Ingredients **Purpose**
 Camphor 3.5% External analgesic
 Menthol 4.5% External analgesic
 Methyl salicylate 10% External analgesic

Uses For the temporary relief of minor aches and pains of muscles and joints due to:
 • simple backache • arthritis
 • strains • bruises • sprains

Warnings

For external use only

Flammable: keep away from fire or flame

Do not use

- on irritated or damaged skin
- on wounds
- otherwise than as directed

When using this product

- avoid contact with the eyes or mucous membranes
- do not bandage tightly

Stop use and seek a doctor if

- condition worsens
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days
- excessive irritation of the skin develops
- nausea, vomiting, abdominal discomfort, diarrhea, or skin rash occurs
- when using for pain of arthritis:
 - pain persists for more than 10 days
 - redness is present
 - in conditions affecting children under 12 years of age

Keep out of reach of children to avoid accidental poisoning. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- shake well before using
- keep nozzle 2 to 4 inches from surface of skin
- adults and children 3 years of age and older: spray onto affected area not more than 3 to 4 times daily
- children under 3 years of age: do not use, consult your physician
- refer to the above warnings; use otherwise than as directed may be dangerous



Yunnan Baiyao SPRAY
 External Analgesic

NDC 61658-006-03

1.7 fl oz (50 mL)

YNBY Spray 50mL - Unit Box
 雲南白藥氣霧劑
 Size: 35 (W) x 35(D) x 105 (H) mm
 05/10/2024



SOLSTICE[®]
 MEDICINE COMPANY

YNBY[®]

EXTERNAL ANALGESIC
 NDC 61658-006-03

**Yunnan Baiyao
 Spray**

1.7 fl. oz. 液量盎司
 (50 mL 毫升)

雲白國際有限公司
 YET INTERNATIONAL LIMITED

Drug Facts (continued)

Other Information

- keep tightly closed
- store at 15° to 30°C (59° to 86°F)

Inactive Ingredients

Alcohol, ampelopsis delavayana root, capaicum, Chinese yim rhizome, corydalis yanhusuo rhizome, dong-quai root, fragrant angelica root, Japanese dioscorea rhizome, san-ci ginseng root and rhizome, and Sichuan lovage rhizome

Questions or Comments?

- (888) 221-3496
 M-F, 9 am to 5 pm
 • you may also report serious side effects to this phone number

U.S. Distributor 美國總代理：
 SOLSTICE MEDICINE COMPANY
 美國蘇氏藥業有限公司
 215 W. Ann Street
 Los Angeles, CA 90012
 Tel: (888) 221-3496
www.solstice.com

Manufactured by 製造商：
 雲南白藥 藥業股份有限公司
 No. 2888 Yunnan Baiyao Street,
 Chengde District, Kunming, Yunnan
 People's Republic of China

MADE IN CHINA
 中國製造

SOLSTICE[®]
 MEDICINE COMPANY

YNBY[®]

EXTERNAL ANALGESIC
 NDC 61658-006-03

**Yunnan Baiyao
 Spray**

1.7 fl. oz. 液量盎司
 (50 mL 毫升)

雲白國際有限公司
 YET INTERNATIONAL LIMITED

Drug Facts

Active Ingredients	Purpose
Camphor 5.1%	External analgesic
Ichthyl 4.1%	External analgesic
Methyl salicylate 10%	External analgesic

Uses For the temporary relief of minor aches and pains of muscles and joints due to:
 • simple backache • arthritis • strain
 • bruise • sprain

Warnings

- For external use only
 Flammable; keep away from fire or flame
 Do not use:
 • on irritated or damaged skin
 • on wounds
 • otherwise than as directed
When using this product:
 • avoid contact with the eyes or mucous membranes
 • do not bandage tightly

Stop use and ask a doctor if:
 • condition worsens
 • symptoms persist for more than 7 days
 • symptoms clear up and occur again within a few days
 • excessive redness of the skin develops
 • nausea, vomiting, abdominal discomfort, diarrhea, or skin rash occurs
 • when using for pain of arthritis:
 • pain persists for more than 10 days
 • redness is present
 • in conditions affecting children under 12 years of age

Keep out of reach of children to avoid accidental poisoning.
 If swallowed get medical help or contact a Poison Control Center right away.

Directions

- shake well before using
- keep nozzle 2 to 4 inches from surface of skin
- adults and children 3 years of age and older: spray onto affected area not more than 2 to 4 times daily
- children under 3 years of age: do not use; consult your physician
- refer to the above warnings; use otherwise than as directed may be dangerous

Lot No:

Exp. Date:



camphor, menthol, and methyl salicylate spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:61658-006
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CAMPHOR (SYNTHETIC) (UNII: 5TJD82A1ET) (CAMPHOR (SYNTHETIC) - UNII:5TJD82A1ET)	CAMPHOR (SYNTHETIC)	3.5 g in 100 mL
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	4.5 g in 100 mL
METHYL SALICYLATE (UNII: LAV5U5022Y) (SALICYLIC ACID - UNII:O414PZ4LPZ)	METHYL SALICYLATE	10 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
AMPELOPSIS DELAVAYANA ROOT (UNII: N93790DMR5)	
CAPSICUM (UNII: 00UK7646FG)	
CHINESE YAM (UNII: 29CIF30B1Z)	
CORYDALIS YANHUSUO WHOLE (UNII: DX4V4VDT5J)	
ANGELICA SINENSIS ROOT (UNII: B66F4574UG)	
ANGELICA DAHURICA ROOT (UNII: 1V63N2S972)	
DIOSCOREA JAPONICA WHOLE (UNII: 1I0396UIN0)	
PANAX NOTOGINSENG WHOLE (UNII: E7XOU43ESD)	
LIGUSTICUM SINENSE SUBSP. CHUANXIONG WHOLE (UNII: CAX256379F)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61658-006-01	1 in 1 BOX	03/19/2024	
1		30 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		
2	NDC:61658-006-02	1 in 1 BOX	03/19/2024	
2		85 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		
3	NDC:61658-006-03	1 in 1 BOX	05/10/2024	
3		50 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
------------------	--	------------------------	----------------------

Category	Citation	Date	Date
OTC Monograph Drug	part348	03/19/2024	

Labeler - YUNNAN BAIYAO GROUP CO., LTD. (654223122)

Establishment

Name	Address	ID/FEI	Business Operations
YUNNAN BAIYAO GROUP CO., LTD.		654223122	manufacture(61658-006)

Revised: 5/2024

YUNNAN BAIYAO GROUP CO., LTD.