

**ACETAMINOPHEN 500 MG- acetaminophen tablet**  
**Pioneer Life Sciences, LLC**

-----  
**Extra Strength Pain Relief ACETAMINOPHEN 500 mg**

***Drug Facts***

***Active ingredient (in each caplet)***

Acetaminophen 500 mg

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- muscular aches
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- If a skin reaction occurs, stop use and seek medical help right away.

***Do not use***

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- If you are allergic to acetaminophen or any of the inactive ingredient in this product.

**Ask a doctor before use if you have**

- liver disease

**Ask a doctor/pharmacist before use if you are**

- taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**KEEP OUT OF REACH OF CHILDREN.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**

**Adults and children 12 years and over:**

- take 2 Caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

**Children under 12 Years**

- ask a doctor

**Other information**

- store at 20° - 25°C (68° -77°F)

**Inactive ingredients**

Pregelatinized starch, Polyvinyl pyrrolidone, stearic acid powder, microcrystalline cellulose, magnesium stearate, Hypromellose-5, Hypromellose-15, polyethylene glycol 400, titanium dioxide, talc

**Questions or comments?**

Call 1 (732) 994-2808 or email [support@gencare.health](mailto:support@gencare.health)

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc.,

McNeil Consumer Healthcare Division., owner of the registered trademark Tyleno® Extra Strength Caplets.

**NDC 72090-038-01 (500ct HDPE Bottle)**

†compare to Tyleno® Extra Strength Caplets active ingredients

# Pain Relief

pain reliever/fever reducer

**EXTRA STRENGTH ACETAMINOPHEN 500mg** each caplet

**500 caplets**

not actual size

NDC 72090-038-01

**TAMPER EVIDENT** Do not use if imprinted safety seal under cap is broken or missing.

Drug Facts
<p><b>Active ingredients (in each caplet)</b></p> <p>Acetaminophen 500 mg</p> <p><b>Purpose</b></p> <p>Pain reliever/fever reducer</p> <p><b>Uses:</b> temporarily relieves minor aches and pains due to: the common cold ■ headache ■ backache ■ minor pain of arthritis ■ muscular aches ■ toothache ■ premenstrual and menstrual cramps ■ temporarily reduces fever</p> <p><b>Warnings:</b> <b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours with other drug containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. <b>Allergy Alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b> ■ with any drug containing acetaminophen (prescription or nonprescription) if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.</p> <p><b>Ask a doctor before use if you have:</b> ■ liver disease</p> <p><b>Ask a doctor/ pharmacist before use if you are:</b> ■ taking the blood thinning drug warfarin. Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur. These could be signs of a serious condition.</p> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use. <b>KEEP OUT OF REACH OF CHILDREN. Overdose Warning:</b> In case of overdose, get medical help or contact Poison Control (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions:</b> ■ do not take more than directed (see overdose warning) ■ Adults and children 12 years and over: ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor ■ Children under 12 years: ask a doctor</p> <p><b>Other Information:</b> ■ store between 20°-25°C (68°-77°F)</p> <p><b>Inactive Ingredients:</b> Pregelatinized starch, polyvinyl pyrrolidone, stearic acid powder, microcrystalline cellulose, magnesium stearate, Hypromellose-5, Hypromellose-15, polyethylene glycol 400, titanium dioxide, talc</p> <p><b>Questions or Comments?</b> Call +1 (732) 994-2808 or email support@gencare.health</p>

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division., owner of the registered trademark Tyleno® Extra Strength. Distributed by: GenCare Consumer Products LLC., East Brunswick, NJ. www.gencare.health P10233-00

Lot No. Exp. Dt.

3 72090 00309 5

**NDC 72090-038-02 (300ct HDPE Bottle)**

†compare to Tylenol® Extra Strength Caplets active ingredients

# Pain Relief

pain reliever/fever reducer

**EXTRA STRENGTH ACETAMINOPHEN 500mg** each caplet

**300 caplets for adults**

not actual size

NDC 72090-038-02

**TAMPER EVIDENT** Do not use if imprinted safety seal under cap is broken or missing.

Drug Facts
<p><b>Active ingredients (in each caplet)</b></p> <p>Acetaminophen 500 mg</p> <p><b>Purpose</b></p> <p>Pain reliever/fever reducer</p> <p><b>Uses:</b> temporarily relieves minor aches and pains due to: the common cold ■ headache ■ backache ■ minor pain of arthritis ■ muscular aches ■ toothache ■ premenstrual and menstrual cramps ■ temporarily reduces fever</p> <p><b>Warnings:</b> <b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours with other drug containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. <b>Allergy Alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b> ■ with any drug containing acetaminophen (prescription or nonprescription) if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.</p> <p><b>Ask a doctor before use if you have:</b> ■ liver disease</p> <p><b>Ask a doctor/ pharmacist before use if you are:</b> ■ taking the blood thinning drug warfarin. Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur. These could be signs of a serious condition.</p> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use. <b>KEEP OUT OF REACH OF CHILDREN. Overdose Warning:</b> In case of overdose, get medical help or contact Poison Control (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions:</b> ■ do not take more than directed (see overdose warning) ■ Adults and children 12 years and over: ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor ■ Children under 12 years: ask a doctor</p> <p><b>Other Information:</b> ■ store between 20°-25°C (68°-77°F)</p> <p><b>Inactive Ingredients:</b> Pregelatinized starch, polyvinyl pyrrolidone, stearic acid powder, microcrystalline cellulose, magnesium stearate, Hypromellose-5, Hypromellose-15, polyethylene glycol 400, titanium dioxide, talc</p> <p><b>Questions or Comments?</b> Call 1-732-994-2808 or visit: www.gencare.health</p>

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division., owner of the registered trademark Tylenol® Extra Strength Caplets. Distributed by: GenCare Consumer Products, LLC 40E Collins Ln, Suite A, East Brunswick, NJ 08816 www.gencare.health P10233-00

Lot No. Exp. Dt.

3 72090 00049 0

**ACETAMINOPHEN 500 MG**

acetaminophen tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72090-038
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	500 mg	
<b>Inactive Ingredients</b>				
Ingredient Name			Strength	
STARCH, CORN (UNII: O8232NY3SJ)				
POVIDONE K30 (UNII: U725QWY32X)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
TALC (UNII: 7SEV7J4R1U)				
<b>Product Characteristics</b>				
Color	white	Score	no score	
Shape	capsule	Size	17mm	
Flavor		Imprint Code	S500	
Contains				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72090-038-01	500 in 1 BOTTLE; Type 0: Not a Combination Product	01/09/2025	
2	NDC:72090-038-02	300 in 1 PACKAGE; Type 0: Not a Combination Product	01/09/2025	
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	01/09/2025		

**Labeler** - Pioneer Life Sciences, LLC (014092742)