

**THOMPSON CRAMP RELIEF- acetaminophen, pamabrom, pyrilamine maleate tablet, film coated**  
**Thompson Safety**

-----  
**Thompson Cramp Relief**

***Drug Facts***

***Active ingredient (in each caplet)***

Acetaminophen 500 mg

Pamabrom 25 mg

Pyrilamine maleate 15 mg

***Purpose***

Pain reliever

Diuretic

Antihistamine

***Uses***

for the temporary relief of these symptoms associated with menstrual periods:

- headache
- bloating
- cramps
- backache
- muscular aches
- irritability
- water-weight gain

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Do not use**

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- liver disease
- glaucoma

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- drowsiness may occur
- avoid alcoholic beverages
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

■ **do not take more than directed**

### **Adults and children: (12 years and over)**

■ take 2 caplets with water every 6 hours as needed

do not exceed 6 caplets in a 24 hour period or as directed by a doctor

**Children under 12 years:** ask a doctor

**Other information**

store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

tamper-evident sealed packets

do not use any opened or torn packets

**Inactive ingredients**

corn starch, croscarmellose sodium, crospovidone, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, shellac wax, silicon dioxide, stearic acid, talc, titanium dioxide

**Questions or comments?** 877.506.4291

Thompson

Cramp Relief

Menstrual Pain Relief

Pain Reliever Acetaminophen 500 mg

Diuretic Pamabrom 25 mg

Antihistamine Pyrilamine Maleate 15 mg

Safety Works | Business Thrives™

Pull to Open

This Package is for Households Without Young Children.

80 Caplets

Tamper Evident Sealed Packets

**CRAMP RELIEF**  
MENSTRUAL PAIN RELIEF  
**THOMPSON**  
SAFETY WORKS | BUSINESS THRIVES

PAIN RELIEVER  
ACETAMINOPHEN 500 MG  
DIURETIC  
PAMABROM 25 MG  
ANTIHISTAMINE  
PYRILAMINE MALEATE 15 MG

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

80 Caplets TAMPER EVIDENT SEALED PACKETS

**Drug Facts**

Active ingredients (in each caplet)	Purpose
Acetaminophen 500 mg	Pain reliever
Pamabrom 25 mg	Diuretic
Pyrilamine maleate 15 mg	Antihistamine

**Uses**  
for the temporary relief of these symptoms associated with menstrual periods:  
 • headache • bloating • cramps • backache • muscular aches • irritability • water-weight gain

**Warnings**  
 • Ever warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours  
 • with other drugs containing acetaminophen  
 • 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
 • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 • if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have** • difficulty in urination due to enlargement of the prostate gland  
 • a breathing problem such as emphysema or chronic bronchitis • liver disease • glaucoma

**Ask a doctor or pharmacist before use if you are** • taking sedatives or tranquilizers  
 • taking the blood thinning drug warfarin (continued on opposite panel)

**CRAMP RELIEF**  
MENSTRUAL PAIN RELIEF  
**THOMPSON**  
SAFETY WORKS | BUSINESS THRIVES

Pull to Open

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

PAIN RELIEVER  
ACETAMINOPHEN 500 MG  
DIURETIC  
PAMABROM 25 MG  
ANTIHISTAMINE  
PYRILAMINE MALEATE 15 MG

80 Caplets TAMPER EVIDENT SEALED PACKETS

**Drug Facts (continued)**

**When using this product** • drowsiness may occur • avoid alcoholic beverages  
 • excitability may occur, especially in children • alcohol, sedatives and tranquilizers may increase drowsiness  
 • use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if** • pain gets worse or lasts more than 10 days  
 • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present  
 These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
 • do not take more than directed  
 • take 2 caplets with water every 6 hours as needed  
 (12 years and over) • do not exceed 6 caplets in a 24 hour period or as directed by a doctor  
 Children under 12 years: ask a doctor

**Other information**  
 • store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
 • tamper-evident sealed packets  
 • do not use any opened or torn packets

**Inactive ingredients** corn starch, croscarmellose sodium, crospovidone, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, shellac wax, silicon dioxide, stearic acid, talc, titanium dioxide

**Questions or comments?** 877.506.4291

**CRAMP RELIEF**  
MENSTRUAL PAIN RELIEF  
**THOMPSON**  
SAFETY WORKS | BUSINESS THRIVES

Reorder C592380

Distributed by  
**Thompson Safety**  
Houston, TX 77073  
877.506.4291  
www.thompson-safety.com

0 73408 92380 6

Retain carton for complete product information

C592380  
Rev. 01/19/22  
Mfg. 24

# THOMPSON CRAMP RELIEF

acetaminophen, pamabrom, pyrilamine maleate tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:73408-923
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
<b>PAMABROM</b> (UNII: UA8U0KJM72) (BROMOTHEOPHYLLINE - UNII:FZ G87K1MQ6)	PAMABROM	25 mg
<b>PYRILAMINE MALEATE</b> (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)	PYRILAMINE MALEATE	15 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CROSPVIDONE</b> (UNII: 2S7830E561)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>FD&amp;C BLUE NO. 2--ALUMINUM LAKE</b> (UNII: 4AQJ3LG584)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

## Product Characteristics

<b>Color</b>	purple	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (Caplet)	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	44;679
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-923-80	40 in 1 BOX	09/26/2022	
1		2 in 1 PACKET; Type 0: Not a Combination		

Product			
<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M027	09/26/2022	

**Labeler** - Thompson Safety (080998015)

Revised: 1/2026

Thompson Safety