

HEAD CONGESTION PLUS FLU SEVERE PE- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated
MEIJER DISTRIBUTION INC

1174-MEI-2022-0923

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purposes</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to the common cold:
 - nasal congestion
 - headache
 - minor aches and pains
 - cough
 - sore throat
 - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 caplets every 4 hours▪ do not take more than 10 caplets in 24 hours
children under 12 years	<ul style="list-style-type: none">▪ ask a doctor

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions or comments?

1-844-705-4384

PRINCIPAL DISPLAY PANEL

meijer®

NDC 41250-574-02

Compare to Sudafed PE® Head Congestion + Flu Severe active ingredients†

head congestion + flu severe PE

Acetaminophen | Dextromethorphan HBr | Guaifenesin | Phenylephrine HCl

Pain Reliever/Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant

Relief of: • Sinus Pressure • Headache

• Sore Throat • Cough • Chest Congestion

24 CAPLETS

actual size



PID 3634002
F117402MEL_R2

Drug Facts (continued)
Other information
 ■ store between 20-25°C (68-77°F) in a dry place
 ■ retain carton for complete product information and warnings

Inactive ingredients
 colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone,

Drug Facts (continued)
 FD&C yellow #6 aluminum lake, magnesium stearate, methocel K100, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions or comments? 1-844-705-4384

Drug Facts (continued)
Warnings
 Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product ■ if a skin reaction occurs, stop use and seek medical help right away. Symptoms may include: ■ skin redness ■ blisters ■ rash ■ severe throat warning: If sore throat is severe, persists for more than 2 days, is accompanied by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
 ■ high blood pressure
 ■ heart disease
 ■ liver disease
 ■ thyroid disease
 ■ diabetes

Drug Facts (continued)
Directions
 ■ do not take more than directed (see overdose warning)
 adults and children ■ take 2 caplets every 4 hours
 12 years and over ■ do not take more than 10 caplets in 24 hours
 children under 12 years ■ ask a doctor

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children.
 If pregnant or breast-feeding, ask a health professional before use. These could be signs of a serious condition.
 ■ cough comes back or occurs with rash or headache that lasts
 ■ redness or swelling is present ■ new symptoms occur
 ■ fever gets worse or lasts more than 3 days
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
 ■ nervousness, dizziness, or sleeplessness occur

Stop use and ask a doctor if
 ■ trouble urinating due to an enlarged prostate gland
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
 ■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Drug Facts
Active ingredients (in each caplet) Purpose
 Acetaminophen 325 mg: Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg: Cough suppressant
 Guaifenesin 100 mg: Expectorant
 Phenylephrine HCl 5 mg: Nasal decongestant

Drug Facts (continued)
Uses ■ temporarily relieves these symptoms due to the common cold: ■ nasal congestion ■ headache ■ minor aches and pains ■ cough ■ sore throat

■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
 ■ temporarily reduces fever

NDC 41250-574-02
Compare to Sudafed PE®
Head Congestion + Flu Severe
active ingredients†

head congestion + flu severe PE

Acetaminophen | Dextromethorphan HBr | Guaifenesin | Phenylephrine HCl
Pain Reliever/Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant

Relief of: • Sinus Pressure • Headache • Sore Throat • Cough • Chest Congestion

24 CAPLETS



actual size

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

†This product is not manufactured or distributed by Johnson & Johnson, owner of the registered trademark, Sudafed PE®.

meijer
 DIST. BY MEIJER DISTRIBUTION, INC.
 GRAND RAPIDS, MI 49514
 www.meijer.com

100% Recycled Content
 100% Recycled Fiber
 100% Recycled Tray

NC

NC

NC

HEAD CONGESTION PLUS FLU SEVERE PE

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-574
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (UNII: 2S7830E561)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	AAA;1134
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:41250-574-02	2 in 1 CARTON	09/23/2022	08/31/2026
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/23/2022	08/31/2026

Labeler - MEIJER DISTRIBUTION INC (006959555)

Revised: 10/2024

MEIJER DISTRIBUTION INC