

**TYLENOL COLD PLUS FLU PLUS COUGH NIGHT- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and phenylephrine hydrochloride solution**  
**Kenvue Brands LLC**

-----

**Drug Facts**

<b>Active ingredients (in each 15 mL)</b>	<b>Purpose</b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine Succinate 6.25 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- temporarily relieves these common cold/flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - runny nose and sneezing
  - cough
  - sinus congestion and pressure
- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

## **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

## **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

## **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

## **When using this product**

- **do not exceed recommended dose**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

## **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see overdose warning)**
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults and children 12 years and over	<ul style="list-style-type: none"><li>▪ take 30 mL in the dosing cup provided every 4 hours while symptoms last</li><li>▪ do not take more than 150 mL in 24 hours, unless directed by a doctor</li></ul>
---------------------------------------	---

children under 12 years	ask a doctor
-------------------------	--------------

### **Other information**

- each 15 mL contains: **sodium 5 mg**
- store between 20-25°C (68-77°F). Do not refrigerate.
- **do not use if neck band imprinted with "TYLENOL COLD+FLU+COUGH" or foil inner seal imprinted with "SAFETY SEAL®" is broken or missing**

### **Inactive ingredients**

acetic acid, alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

### **Questions or comments?**

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

### **PRINCIPAL DISPLAY PANEL**

NDC 50580-369-01

TYLENOL®  
FOR ADULTS  
COLD + FLU + COUGH

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate, Phenylephrine HCl, Pain Reliever-Fever Reducer, Cough Suppressant, Antihistamine, Nasal Decongestant

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- RUNNY NOSE

NIGHT

WILD BERRY  
BURST

Alcohol 0.7%  
8 fl oz (240 mL)

30040371

NDC 50580-369-01

# TYLENOL<sup>®</sup>

FOR ADULTS

## COLD + FLU + COUGH

**Acetaminophen**, Dextromethorphan HBr, Doxylamine Succinate, Phenylephrine HCl,  
Pain Reliever—Fever Reducer, Cough Suppressant, Antihistamine, Nasal Decongestant



- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- RUNNY NOSE



NIGHT



WILD BERRY  
BURST



Alcohol 0.7%

8 fl oz (240 mL)

OPEN TO READ DRUG FACTS  
(Warnings, Directions...)




PEEL



30040372 Do not use if neck band imprinted with  
"TYLENOL COLD+FLU+ COUGH" or foil  
inner seal imprinted with "SAFETY  
SEAL<sup>®</sup>" is broken or missing.

**DO NOT USE WITH OTHER MEDICINES  
CONTAINING ACETAMINOPHEN**



3 0045-0562-081

## Drug Facts

### Active ingredients (in each 15 mL) Purpose

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine Succinate 6.25 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

### Uses

- temporarily relieves these common cold/flu symptoms:
  - minor aches and pains
  - headache
  - nasal congestion
  - runny nose and sneezing
  - cough
  - sinus congestion and pressure
  - helps clear nasal passages
- sore throat

## Drug Facts (continued)

- relieves cough to help you sleep
- temporarily reduces fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

## Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional

## Drug Facts (continued)

conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as

## Drug Facts (continued)

## Drug Facts (continued)

- emphysema or chronic bronchitis
- glaucoma
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- **do not exceed recommended dose**
- excitability may occur, especially in children
- marked drowsiness may occur ▶

- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
  - pain, nasal congestion or cough gets worse or lasts more than 7 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition. ▶

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device. ▶

**Drug Facts (continued)**

adults and children 12 years and over	<ul style="list-style-type: none"> <li>■ take 30 mL in the dosing cup provided every 4 hours while symptoms last</li> <li>■ do not take more than 150 mL in 24 hours, unless directed by a doctor</li> </ul>
children under 12 years	ask a doctor

**Other information**

- each 15 mL contains: **sodium 5 mg**
- store between 20-25°C (68-77°F). Do not refrigerate.
- **do not use if neck band imprinted with "TYLENOL COLD+FLU+COUGH" or foil inner seal imprinted with "SAFETY SEAL®"** ▶

**Drug Facts (continued)**

is broken or missing

**Inactive ingredients**

acetic acid, alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no.

**TYLENOL®**  
**COLD + FLU + COUGH**  
**NIGHT**

*How can we help?*  
**1-877-895-3665**

40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

**Questions or comments?**  
call **1-877-895-3665** (toll-free) or  
**215-273-8755** (collect)

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**SNAP THIS CODE  
OR TEXT TYCOLD7  
TO 87715**

MSG & DATA RATES MAY APPLY  
TEXT HELP FOR HELP.  
TEXT STOP FOR STOP.



Acetaminophen  
(325 mg per 15 mL)  
Dextromethorphan HBr  
(10 mg per 15 mL)  
Doxylamine succinate  
(6.25 mg per 15 mL)  
Phenylephrine HCl  
(5 mg per 15 mL)  
8 fl oz (240 mL)

Distributed by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA

© J&JCI 2017

**30040372**

## TYLENOL COLD PLUS FLU PLUS COUGH NIGHT

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and phenylephrine hydrochloride solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50580-369
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	

<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>ACETIC ACID</b> (UNII: Q40Q9N063P)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Product Characteristics

<b>Color</b>	purple	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY (Berry Mint)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-369-01	240 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package	06/17/2019	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/17/2019	

**Labeler** - Kenvue Brands LLC (118772437)

Revised: 11/2024

Kenvue Brands LLC