

**RASAGILINE - rasagiline tablet**  
**Macleods Pharmaceuticals Limited**

-----  
**HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use RASAGILINE TABLETS safely and effectively. See full prescribing information for RASAGILINE TABLETS.**

**RASAGILINE tablets, for oral use**

**Initial U.S. Approval: 2006**

-----  
**INDICATIONS AND USAGE**

Rasagiline tablets, a monoamine oxidase (MAO)-B inhibitor (MAOI), is indicated for the treatment of Parkinson's disease (1)

-----  
**DOSAGE AND ADMINISTRATION**

- Monotherapy: Rasagiline tablets 1 mg once daily (2.1)
- As adjunct without levodopa: Rasagiline tablets 1 mg once daily (2.1)
- As adjunct to levodopa: Rasagiline tablets 0.5 mg once daily. Increase dose to 1 mg daily as needed for sufficient clinical response (2.1)
- Patients taking ciprofloxacin or other CYP1A2 inhibitors: Rasagiline tablets 0.5 mg once daily (2.2, 5.4)
- Patients with mild hepatic impairment: Rasagiline tablets 0.5 mg once daily. Rasagiline tablets should not be used in patients with moderate or severe hepatic impairment (2.3, 5.5)

-----  
**DOSAGE FORMS AND STRENGTHS**

- Rasagiline Tablets 0.5 mg tablets (3)
- Rasagiline Tablets 1 mg tablets (3)

-----  
**CONTRAINDICATIONS**

Concomitant use of meperidine, tramadol, methadone, propoxyphene dextromethorphan, St. John's wort, cyclobenzaprine, or another (selective or non-selective) MAO inhibitor (4)

-----  
**WARNINGS AND PRECAUTIONS**

- May cause hypertension (including severe hypertensive syndromes) at recommended doses (5.1)
- May cause serotonin syndrome when used with antidepressants (5.2)
- May cause falling asleep during activities of daily living, daytime drowsiness, and somnolence (5.3)
- May cause hypotension, especially orthostatic (5.6)
- May cause or exacerbate dyskinesia. Decreasing the levodopa dose may lessen or eliminate this side effect (5.7)
- May cause hallucinations and psychotic-like behavior (5.8)
- May cause impulse control/compulsive behaviors (5.9)
- May cause withdrawal-emergent hyperpyrexia and confusion (5.10)

-----  
**ADVERSE REACTIONS**

Most common adverse reactions (incidence 3% or greater than placebo):

- Rasagiline tablets monotherapy: flu syndrome, arthralgia, depression, dyspepsia (6.1)
- Rasagiline tablets used as adjunct without levodopa: peripheral edema, fall, arthralgia, cough, and insomnia (6.1)
- Rasagiline tablets used as adjunct to levodopa: dyskinesia, accidental injury, weight loss, postural hypotension, vomiting, anorexia, arthralgia, abdominal pain, nausea, constipation, dry mouth, rash, abnormal dreams, fall, and tenosynovitis (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Macleods Pharma USA, Inc., at 1-888-943-3210 or 1-855-926-3384 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)**

-----  
**DRUG INTERACTIONS**

- Meperidine: Risk of serotonin syndrome (4, 7.1)
- Dextromethorphan: Risk of psychosis or bizarre behavior (4, 7.2)
- MAO inhibitors: Risk of non-selective MAO inhibition and hypertensive crisis (4, 7.3)

-----  
**USE IN SPECIFIC POPULATIONS**

- Pregnancy: Based on animal data, may cause fetal harm. (8.1)

**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 7/2024**

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS & USAGE**

**2 DOSAGE & ADMINISTRATION**

- 2.1 General Dosing Recommendations
- 2.2 Patients Taking Ciprofloxacin or Other CYP1A2 Inhibitors
- 2.3 Patients with Hepatic Impairment

**3 DOSAGE FORMS & STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

- 5.1 Hypertension
- 5.2 Serotonin Syndrome
- 5.3 Falling Asleep During Activities of Daily Living and Somnolence
- 5.4 Ciprofloxacin or Other CYP1A2 Inhibitors
- 5.5 Hepatic Impairment
- 5.6 Hypotension / Orthostatic Hypotension
- 5.7 Dyskinesia
- 5.8 Hallucinations / Psychotic-Like Behavior

- 5.9 Impulse Control / Compulsive Behaviors
- 5.10 Withdrawal-Emergent Hyperpyrexia and Confusion

## **6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

## **7 DRUG INTERACTIONS**

- 7.1 Meperidine
- 7.2 Dextromethorphan
- 7.3 MAO Inhibitors
- 7.4 Sympathomimetic Medications
- 7.5 Antidepressants
- 7.6 Ciprofloxacin or Other CYP1A2 Inhibitors
- 7.7 Tyramine/Rasagiline Interaction
- 7.8 Dopaminergic Antagonists

## **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment
- 8.7 Renal Impairment

## **9 DRUG ABUSE AND DEPENDENCE**

- 9.1 Controlled Substance
- 9.2 Abuse
- 9.3 Dependence

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

## **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis & Mutagenesis & Impairment of Fertility

## **14 CLINICAL STUDIES**

- 14.1 Monotherapy Use of Rasagiline Tablets
- 14.2 Adjunct Use of Rasagiline tablets

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS & USAGE**

Rasagiline tablets are indicated for the treatment of Parkinson's disease (PD).

### **2 DOSAGE & ADMINISTRATION**

#### **2.1 General Dosing Recommendations**

When rasagiline tablets are prescribed as monotherapy or as adjunct therapy in patients not taking levodopa, patients may start rasagiline tablets at the recommended dose of 1 mg administered orally once daily.

In patients taking levodopa, with or without other PD drugs (e.g., dopamine agonist, amantadine, anticholinergics) the recommended initial dose of rasagiline tablets is 0.5 mg once daily. If the patient tolerates the daily 0.5 mg dose, but a sufficient clinical response is not achieved, the dose may be increased to 1 mg once daily.

When rasagiline tablets are used in combination with levodopa, a reduction of the levodopa dose may be considered, based upon individual response.

The recommended doses of rasagiline tablets should not be exceeded because of risk of hypertension [see *Warnings and Precautions (5.1)*].

#### **2.2 Patients Taking Ciprofloxacin or Other CYP1A2 Inhibitors**

Patients taking concomitant ciprofloxacin or other CYP1A2 inhibitors should not exceed a dose of rasagiline tablets 0.5 mg once daily [see *Warnings and Precautions (5.4)*, *Drug Interactions (7.6)*, and *Clinical Pharmacology (12.3)*].

### 2.3 Patients with Hepatic Impairment

Patients with mild hepatic impairment should not exceed a dose of rasagiline tablets 0.5 mg once daily. Rasagiline tablets should not be used in patients with moderate or severe hepatic impairment [see *Warnings and Precautions (5.5)*, *Use in Specific Populations (8.6)*, and *Clinical Pharmacology (12.3)*].

### 3 DOSAGE FORMS & STRENGTHS

Rasagiline 0.5 mg Tablets: White to off-white, round, flat, beveled tablets, debossed with "C12" on one side and plain on the other side.

Rasagiline 1 mg Tablets: White to off-white, round, flat, beveled tablets, debossed with "C13" on one side and plain on the other side.

### 4 CONTRAINDICATIONS

Rasagiline Tablets are contraindicated for use with meperidine, tramadol, methadone, propoxyphene and MAO inhibitors (MAOIs), including other selective MAO-B inhibitors, because of risk of serotonin syndrome [See *Warnings and Precautions (5.2)*]. At least 14 days should elapse between discontinuation of rasagiline tablets and initiation of treatment with these medications.

Rasagiline Tablets are contraindicated for use with St. John's wort and with cyclobenzaprine.

Rasagiline Tablets are contraindicated for use with dextromethorphan because of risk of episode of psychosis or bizarre behavior.

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Hypertension

Exacerbation of hypertension may occur during treatment with rasagiline tablets. Medication adjustment may be necessary if elevation of blood pressure is sustained. Monitor patients for new onset hypertension or hypertension that is not adequately controlled after starting rasagiline tablets.

In Study 3, rasagiline tablets (1 mg/day) given in conjunction with levodopa, produced an increased incidence of significant blood pressure elevation (systolic > 180 or diastolic > 100 mm Hg) of 4% compared to 3% for placebo [see *Adverse Reactions (6.1)*].

When used as an adjunct to levodopa (Studies 3 and 4), the risk for developing post-treatment high blood pressure (e.g., systolic > 180 or diastolic >100 mm Hg) combined with a significant increase from baseline (e.g., systolic > 30 or diastolic > 20 mm Hg) was higher for rasagiline tablets (2%) compared to placebo (1%).

Dietary tyramine restriction is not required during treatment with recommended doses of rasagiline tablets. However, certain foods that may contain very high amounts (i.e., more than 150 mg) of tyramine that could potentially cause severe hypertension because of tyramine interaction (including various clinical syndromes referred to as hypertensive urgency, crisis, or emergency) in patients taking rasagiline tablets, even at the recommended doses, due to increased sensitivity to tyramine. Patients should be advised to avoid foods containing a very large amount of tyramine while taking recommended doses of rasagiline tablets because of the potential for large increases in blood pressure including clinical syndromes referred to as hypertensive urgency, crisis, or emergency. Rasagiline tablets are a selective inhibitor of MAO-B at the recommended doses of 0.5 or 1 mg daily. Selectivity for inhibiting MAO-B diminishes in a dose-related manner as the dose is progressively increased above the recommended daily doses.

#### 5.2 Serotonin Syndrome

Serotonin syndrome has been reported with concomitant use of an antidepressant (e.g., selective serotonin reuptake inhibitors-SSRIs, serotonin-norepinephrine reuptake inhibitors-SNRIs, tricyclic antidepressants, tetracyclic antidepressants, triazolopyridine antidepressants) and a nonselective MAOI (e.g., phenelzine, tranylcypromine) or selective MAO-B inhibitors, such as selegiline (Eldepryl) and rasagiline tablets. Serotonin syndrome has also been reported with concomitant use of rasagiline tablets with meperidine, tramadol, methadone, or propoxyphene. Rasagiline tablets are contraindicated for use with meperidine, tramadol, methadone, propoxyphene and MAO inhibitors (MAOIs), including other selective MAO-B inhibitors [see *Contraindications (4)* and *Drug Interactions (7.1, 7.2, 7.3)*].

In the postmarketing period, potentially life-threatening serotonin syndrome has been

reported in patients treated with antidepressants concomitantly with rasagiline tablets. Concomitant use of rasagiline tablets with one of many classes of antidepressants (e.g., SSRIs, SNRIs, triazolopyridine, tricyclic or tetracyclic antidepressants) is not recommended [see *Drug Interactions (7.5)*].

The symptoms of serotonin syndrome have included behavioral and cognitive/mental status changes (e.g., confusion, hypomania, hallucinations, agitation, delirium, headache, and coma), autonomic effects (e.g., syncope, shivering, sweating, high fever/hyperthermia, hypertension, tachycardia, nausea, diarrhea), and somatic effects (e.g., muscular rigidity, myoclonus, muscle twitching, hyperreflexia manifested by clonus, and tremor). Serotonin syndrome can result in death.

Rasagiline tablets clinical trials did not allow concomitant use of fluoxetine or fluvoxamine with rasagiline tablets, and the potential drug interaction between rasagiline tablets and antidepressants has not been studied systematically. Although a small number of rasagiline tablets-treated patients were concomitantly exposed to antidepressants (tricyclics n=115; SSRIs n=141), the exposure, both in dose and number of subjects, was not adequate to rule out the possibility of an untoward reaction from combining these agents. At least 14 days should elapse between discontinuation of rasagiline tablets and initiation of treatment with a SSRI, SNRI, tricyclic, tetracyclic, or triazolopyridine antidepressant. Because of the long half-lives of certain antidepressants (e.g., fluoxetine and its active metabolite), at least five weeks (perhaps longer, especially if fluoxetine has been prescribed chronically and/or at higher doses) should elapse between discontinuation of fluoxetine and initiation of rasagiline tablets [see *Drug Interactions (7.5)*].

### **5.3 Falling Asleep During Activities of Daily Living and Somnolence**

It has been reported that falling asleep while engaged in activities of daily living always occurs in a setting of preexisting somnolence, although patients may not give such a history. For this reason, prescribers should monitor patients for drowsiness or sleepiness, because some of the events occur well after initiation of treatment with dopaminergic medication. Prescribers should also be aware that patients may not acknowledge drowsiness or sleepiness until directly questioned about drowsiness or sleepiness during specific activities.

Cases of patients treated with rasagiline tablets and other dopaminergic medications have reported falling asleep while engaged in activities of daily living including the operation of motor vehicles, which sometimes resulted in accidents. Although many of these patients reported somnolence while on rasagiline tablets with other dopaminergic medications, some perceived that they had no warning signs, such as excessive drowsiness, and believed that they were alert immediately prior to the event. Some of these events have been reported more than 1-year after initiation of treatment.

In Study 3, somnolence was a common occurrence in patients receiving rasagiline tablets and was more frequent in patients with Parkinson's disease receiving rasagiline tablets than in respective patients receiving placebo (6% rasagiline tablets compared to 4% Placebo)[see *Adverse Reactions (6.1)*].

Before initiating treatment with rasagiline tablets, patients should be advised of the potential to develop drowsiness and specifically asked about factors that may increase the risk with rasagiline tablets such as concomitant sedating medications, the presence of sleep disorders, and concomitant medications that increase rasagiline plasma levels (e.g., ciprofloxacin) [see *Drug Interactions (7.6)*]. If a patient develops significant daytime sleepiness or episodes of falling asleep during activities that require active participation (e.g., driving a motor vehicle, conversations, eating), rasagiline tablets should ordinarily be discontinued. If a decision is made to continue these patients on rasagiline tablets, advise them to avoid driving and other potentially dangerous activities. There is insufficient information to establish that dose reduction will eliminate episodes of falling asleep while engaged in activities of daily living.

### **5.4 Ciprofloxacin or Other CYP1A2 Inhibitors**

Rasagiline plasma concentrations may increase up to 2 fold in patients using concomitant ciprofloxacin and other CYP1A2 inhibitors. Patients taking concomitant ciprofloxacin or other CYP1A2 inhibitors should not exceed a dose of rasagiline tablets 0.5 mg once daily [see *Dosage and Administration (2.2)*, *Drug Interactions (7.6)*, and *Clinical Pharmacology (12.3)*].

### **5.5 Hepatic Impairment**

Rasagiline plasma concentration may increase in patients with hepatic impairment.

Patients with mild hepatic impairment should be given the dose of rasagiline tablets 0.5 mg once daily. Rasagiline tablets should not be used in patients with moderate or severe hepatic impairment [see *Dosage and Administration (2.3) and Clinical Pharmacology (12.3)*].

## 5.6 Hypotension / Orthostatic Hypotension

In Study 3, the incidence of orthostatic hypotension consisting of a systolic blood pressure decrease (> 30 mm Hg) or a diastolic blood pressure decrease (> 20 mm Hg) after standing was 13% with rasagiline tablets (1 mg/day) compared to 9% with placebo [see *Adverse Reactions (6.1)*].

At the 1 mg dose, the frequency of orthostatic hypotension (at any time during the study) was approximately 44% for rasagiline tablets vs 33% for placebo for mild to moderate systolic blood pressure decrements (> 20 mm Hg), 40% for rasagiline tablets vs 33% for placebo for mild to moderate diastolic blood pressure decrements (> 10 mm Hg), 7% for rasagiline tablets vs 3% for placebo for severe systolic blood pressure decrements (> 40 mm Hg), and 9% for rasagiline tablets vs 6% for placebo for severe diastolic blood pressure decrements ( $\geq$  20 mm Hg). There was also an increased risk for some of these abnormalities at the lower 0.5 mg daily dose and for an individual patient having mild to moderate or severe orthostatic hypotension for both systolic and diastolic blood pressure.

In Study 2 where rasagiline tablets were given as an adjunct therapy in patients not taking concomitant levodopa, there were 5 reports of orthostatic hypotension in patients taking rasagiline tablets 1 mg (3.1%) and 1 report in patients taking placebo (0.6%) [see *Adverse Reactions (6.1)*].

Clinical trial data further suggest that orthostatic hypotension occurs most frequently in the first two months of rasagiline tablets treatment and tends to decrease over time. Some patients treated with rasagiline tablets experienced a mildly increased risk for significant decreases in blood pressure unrelated to standing but while supine. The risk for post-treatment hypotension (e.g., systolic < 90 or diastolic < 50 mm Hg) combined with a significant decrease from baseline (e.g., systolic > 30 or diastolic > 20 mm Hg) was higher for rasagiline tablets 1 mg (3.2%) compared to placebo (1.3%). There was no clear increased risk for lowering of blood pressure or postural hypotension associated with rasagiline tablets 1 mg/day as monotherapy. When used as an adjunct to levodopa, postural hypotension was also reported as an adverse reaction in approximately 6% of patients treated with rasagiline tablets 0.5 mg, 9% of patients treated with rasagiline tablets 1 mg and 3% of patients treated with placebo. Postural hypotension led to drug discontinuation and premature withdrawal from clinical trials in one (0.7%) patient treated with rasagiline tablets 1 mg/day, no patients treated with rasagiline tablets 0.5 mg/day and no placebo-treated patients.

## 5.7 Dyskinesia

When used as an adjunct to levodopa, rasagiline tablets may cause dyskinesia or potentiate dopaminergic side effects and exacerbate pre-existing dyskinesia. In Study 3, the incidence of dyskinesia was 18% for patients treated with 0.5 mg or 1 mg rasagiline tablets as an adjunct to levodopa and 10% for patients treated with placebo as an adjunct to levodopa. Decreasing the dose of levodopa may mitigate this side effect [see *Adverse Reactions (6.1)*].

## 5.8 Hallucinations / Psychotic-Like Behavior

In the monotherapy study (Study 1), the incidence of hallucinations reported as an adverse event was 1.3% in patients treated with rasagiline tablets 1 mg and 0.7% in patients treated with placebo. In Study 1, the incidence of hallucinations reported as an adverse reaction and leading to drug discontinuation and premature withdrawal was 1.3% in patients treated with rasagiline tablets 1 mg and 0% in placebo-treated patients.

When studied as an adjunct therapy without levodopa (Study 2), hallucinations were reported as an adverse reaction in 1.2% of patients treated with 1 mg/day rasagiline tablets and 1.8% of patients treated with placebo. Hallucination led to drug discontinuation and premature withdrawal from the clinical trial in 0.6% of patients treated with rasagiline tablets 1 mg/day and in none of the placebo-treated patients.

When studied as an adjunct to levodopa (Study 3), the incidence of hallucinations was approximately 5% in patients treated with rasagiline tablets 0.5 mg/day, 4% in patients treated with rasagiline tablets 1 mg/day and 3% in patients treated with placebo. The

incidence of hallucinations leading to drug discontinuation and premature withdrawal was about 1% in patients treated with 0.5 mg rasagiline tablets and 1 mg rasagiline tablets /day, and 0% in placebo-treated patients [see *Adverse Reactions (6.1)*].

Postmarketing reports indicate that patients may experience new or worsening mental status and behavioral changes, which may be severe, including psychotic-like behavior during treatment with rasagiline tablets or after starting or increasing the dose of rasagiline tablets. Other drugs prescribed to improve the symptoms of Parkinson's disease can have similar effects on thinking and behavior. This abnormal thinking and behavior can consist of one or more of a variety of manifestations including paranoid ideation, delusions, hallucinations, confusion, psychotic-like behavior, disorientation, aggressive behavior, agitation, and delirium.

Patients should be informed of the possibility of developing hallucinations and instructed to report them to their healthcare provider promptly should they develop.

Patients with a major psychotic disorder should ordinarily not be treated with rasagiline tablets because of the risk of exacerbating the psychosis with an increase in central dopaminergic tone. In addition, many treatments for psychosis that decrease central dopaminergic tone may decrease the effectiveness of rasagiline tablets [see *Drug Interactions (7.8)*].

Consider dose reduction or stopping the medication if a patient develops hallucinations or psychotic like behaviors while taking rasagiline tablets.

### **5.9 Impulse Control / Compulsive Behaviors**

Case reports suggest that patients can experience intense urges to gamble, increased sexual urges, intense urges to spend money, binge eating, and/or other intense urges, and the inability to control these urges while taking one or more of the medications, including rasagiline tablets, that increase central dopaminergic tone and that are generally used for the treatment of Parkinson's disease. In some cases, although not all, these urges were reported to have stopped when the dose was reduced or the medication was discontinued. Because patients may not recognize these behaviors as abnormal, it is important for prescribers to specifically ask patients or their caregivers about the development of new or increased gambling urges, sexual urges, uncontrolled spending or other urges while being treated with rasagiline tablets. Consider dose reduction or stopping the medication if a patient develops such urges while taking rasagiline tablets.

### **5.10 Withdrawal-Emergent Hyperpyrexia and Confusion**

A symptom complex resembling neuroleptic malignant syndrome (characterized by elevated temperature, muscular rigidity, altered consciousness, and autonomic instability), with no other obvious etiology, has been reported in association with rapid dose reduction, withdrawal of, or changes in drugs that increase central dopaminergic tone.

## **6 ADVERSE REACTIONS**

The following adverse reactions are described in more detail in the Warnings and Precautions section of the label:

- Hypertension [see *Warnings and Precautions (5.1)*]
- Serotonin Syndrome [see *Warnings and Precautions (5.2)*]
- Falling Asleep During Activities of Daily Living and Somnolence [see *Warnings and Precautions (5.3)*]
- Hypotension / Orthostatic Hypotension [see *Warnings and Precautions (5.6)*]
- Dyskinesia [see *Warnings and Precautions (5.7)*]
- Hallucinations / Psychotic-Like Behavior [see *Warnings and Precautions (5.8)*]
- Impulse Control /Compulsive Behaviors [see *Warnings and Precautions (5.9)*]
- Withdrawal-Emergent Hyperpyrexia and Confusion [see *Warnings and Precautions (5.10)*]

### **6.1 Clinical Trials Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to the incidence of adverse reactions in the clinical trials of another drug and may not reflect the rates of adverse reactions observed in practice.

During the clinical development of rasagiline tablets, Parkinson's disease patients received rasagiline tablets as initial monotherapy (Study 1) and as adjunct therapy (Study 2, Study 3, Study 4). As the populations in these studies differ, not only in the adjunct use of dopamine agonists or levodopa during rasagiline tablets treatment, but

also in the severity and duration of their disease, the adverse reactions are presented separately for each study.

#### Monotherapy Use of Rasagiline Tablets

In Study 1, approximately 5% of the 149 patients treated with rasagiline tablets discontinued treatment due to adverse reactions compared to 2% of the 151 patients who received placebo.

The only adverse reaction that led to the discontinuation of more than one patient was hallucinations.

The most commonly observed adverse reactions in Study 1 (incidence in rasagiline tablets -treated patients 3% or greater than the incidence in placebo-treated patients) included flu syndrome, arthralgia, depression, and dyspepsia. Table 1 lists adverse reactions that occurred in 2% or greater of patients receiving rasagiline tablets as monotherapy and were numerically more frequent than in the placebo group in Study 1.

**Table 1: Adverse Reactions\* in Study 1**

	<b>Rasagiline Tablets 1 mg (N=149)</b>	<b>Placebo (N=151)</b>
	<b>% of Patients</b>	<b>% of Patients</b>
Headache	14	12
Arthralgia	7	4
Dyspepsia	7	4
Depression	5	2
Fall	5	3
Flu syndrome	5	1
Conjunctivitis	3	1
Fever	3	1
Gastroenteritis	3	1
Rhinitis	3	1
Arthritis	2	1
Ecchymosis	2	0
Malaise	2	0
Neck Pain	2	0
Paresthesia	2	1
Vertigo	2	1

\*Incidence 2% or greater in rasagiline tablets 1 mg group and numerically more frequent than in placebo group  
There were no significant differences in the safety profile based on age or gender.

#### Adjunct Use of Rasagiline Tablets

Rasagiline tablets were studied as an adjunct therapy without levodopa (Study 2), or as an adjunct therapy to levodopa, with some patients also taking dopamine agonists, COMT inhibitors, anticholinergics, or amantadine (Study 3 and Study 4).

In Study 2, approximately 8% of the 162 patients treated with rasagiline tablets discontinued treatment due to adverse reactions compared to 4% of the 164 patients who received placebo.

Adverse reactions that led to the discontinuation of more than one patient were nausea and dizziness.

The most commonly observed adverse reactions in Study 2 (incidence in rasagiline

tablets -treated patients 3% or greater than incidence in placebo-treated patients) included peripheral edema, fall, arthralgia, cough, and insomnia. Table 2 lists adverse reactions that occurred in 2% or greater in patients receiving rasagiline tablets as adjunct therapy without levodopa and numerically more frequent than in the placebo group in Study 2.

Table 2: Adverse Reactions\* in Study 2

	<b>Rasagiline Tablets 1 mg (N=162)</b>	<b>Placebo (N=164)</b>
	<b>% of Patients</b>	<b>% of Patients</b>
Dizziness	7	6
Peripheral edema	7	4
Headache	6	4
Nausea	6	4
Fall	6	1
Arthralgia	5	2
Back pain	4	3
Cough	4	1
Insomnia	4	1
Upper respiratory tract infection	4	2
Orthostatic hypotension	3	1

\*Incidence 2% or greater in rasagiline tablets 1 mg group and numerically more frequent than in placebo group

There were no significant differences in the safety profile based on age or gender.

In Study 3, adverse event reporting was considered more reliable than Study 4; therefore, only the adverse event data from Study 3 are presented below.

In Study 3, approximately 9% of the 164 patients treated with rasagiline tablets 0.5 mg/day and 7% of the 149 patients treated with rasagiline tablets 1 mg/day discontinued treatment due to adverse reactions, compared to 6% of the 159 patients who received placebo. The adverse reactions that led to discontinuation of more than one rasagiline - treated patient were diarrhea, weight loss, hallucination, and rash.

The most commonly observed adverse reactions in Study 3 (incidence in rasagiline tablets -treated patients 3% or greater than the incidence in placebo-treated patients) included dyskinesia, accidental injury, weight loss, postural hypotension, vomiting, anorexia, arthralgia, abdominal pain, nausea, constipation, dry mouth, rash, abnormal dreams, fall and tenosynovitis.

Table 3 lists adverse reactions that occurred in 2% or greater of patients treated with rasagiline tablets 1 mg/day and that were numerically more frequent than the placebo group in Study 3.

Table 3: Adverse Reactions\* in Study 3

	<b>Rasagiline Tablets 1 mg (N=149)</b>	<b>Rasagiline Tablets 0.5 mg (N=164)</b>	<b>Placebo (N=159)</b>
	<b>% of Patients</b>	<b>% of Patients</b>	<b>% of Patients</b>
Dyskinesia	18	18	10
Accidental injury	12	8	5



Nausea	12	10	8
Headache	11	8	10
Fall	11	12	8
Weight loss	9	2	3
Constipation	9	4	5
Postural hypotension	9	6	3
Arthralgia	8	6	4
Vomiting	7	4	1
Dry mouth	6	2	3
Rash	6	3	3
Somnolence	6	4	4
Abdominal pain	5	2	1
Anorexia	5	2	1
Diarrhea	5	7	4
Ecchymosis	5	2	3
Dyspepsia	5	4	4
Paresthesia	5	2	3
Abnormal dreams	4	1	1
Hallucinations	4	5	3
Ataxia	3	6	1
Dyspnea	3	5	2
Infection	3	2	2
Neck pain	3	1	1
Sweating	3	2	1
Tenosynovitis	3	1	0
Dystonia	3	2	1
Gingivitis	2	1	1
Hemorrhage	2	1	1
Hernia	2	1	1
Myasthenia	2	2	1

\*Incidence 2% or greater in rasagiline tablets 1 mg group and numerically more frequent than in placebo group. Several of the more common adverse reactions seemed dose-related, including weight loss, postural hypotension, and dry mouth. There were no significant differences in the safety profile based on age or gender. During all Parkinson's disease phase 2/3 clinical trials, the long-term safety profile was similar to that observed with shorter duration exposure.

## 6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of rasagiline tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a

causal relationship to drug exposure.  
*Skin and Subcutaneous Tissue Disorders: Melanoma*

## **7 DRUG INTERACTIONS**

### **7.1 Meperidine**

Serious, sometimes fatal reactions have been precipitated with concomitant use of meperidine (e.g., Demerol and other tradenames) and MAO inhibitors including selective MAO-B inhibitors [see *Contraindications (4)*].

### **7.2 Dextromethorphan**

The concomitant use of rasagiline tablets and dextromethorphan was not allowed in clinical studies. The combination of MAO inhibitors and dextromethorphan has been reported to cause brief episodes of psychosis or bizarre behavior. Therefore, in view of rasagiline's MAO inhibitory activity, dextromethorphan is contraindicated for use with rasagiline tablets [see *Contraindications (4)*].

### **7.3 MAO Inhibitors**

Rasagiline tablets are contraindicated for use with other MAO inhibitors because of the increased risk of nonselective MAO inhibition that may lead to a hypertensive crisis [see *Contraindications (4)*].

### **7.4 Sympathomimetic Medications**

The concomitant use of rasagiline tablets and sympathomimetic medications was not allowed in clinical studies. Severe hypertensive reactions have followed the administration of sympathomimetics and nonselective MAO inhibitors. Hypertensive crisis has been reported in patients taking the recommended dose of rasagiline tablets and sympathomimetic medications. Severe hypertension has been reported in patients taking the recommended dose of rasagiline tablets and ophthalmic drops containing sympathomimetic medications.

Because rasagiline tablets are a selective MAOI, hypertensive reactions are not ordinarily expected with the concomitant use of sympathomimetic medications. Nevertheless, caution should be exercised when concomitantly using recommended doses of rasagiline tablets with any sympathomimetic medications including nasal, oral, and ophthalmic decongestants and cold remedies.

### **7.5 Antidepressants**

Concomitant use of rasagiline tablets with one of many classes of antidepressants (e.g., SSRIs, SNRIs, triazolopyridine, tricyclic or tetracyclic antidepressants) is not recommended [see *Warnings and Precautions (5.2) and Clinical Pharmacology (12.3)*]. Concomitant use of rasagiline tablets and MAO inhibitors is contraindicated [see *Contraindications (4)*].

### **7.6 Ciprofloxacin or Other CYP1A2 Inhibitors**

Rasagiline plasma concentrations may increase up to 2 fold in patients using concomitant ciprofloxacin and other CYP1A2 inhibitors. This could result in increased adverse events. Patients taking concomitant ciprofloxacin or other CYP1A2 inhibitors should not exceed a dose of rasagiline tablets 0.5 mg once daily [see *Warnings and Precautions (5.4) and Clinical Pharmacology (12.3)*].

### **7.7 Tyramine/Rasagiline Interaction**

MAO in the gastrointestinal tract and liver (primarily type A) provides protection from exogenous amines (e.g., tyramine) that have the capacity, if absorbed intact, to cause a tyramine reaction with hypertension including clinical syndromes referred to as hypertensive urgency, crisis, or emergency. Foods and medications containing large amounts of exogenous amines (e.g., from fermented cheese, herring, over-the-counter cough/cold medications) may cause release of norepinephrine resulting in a rise in systemic blood pressure.

Results of a special tyramine challenge study indicate that rasagiline is selective for MAO-B at recommended doses and can be used without dietary tyramine restriction. However, certain foods may contain very high amounts (i.e., 150 mg or greater) of tyramine and could potentially cause a hypertensive reaction in individual patients taking

rasagiline tablets due to increased sensitivity to tyramine. Selectivity for inhibiting MAO-B diminishes in a dose-related manner as the dose is progressively increased above the recommended daily doses.

There were no cases of hypertensive crisis in the clinical development program associated with 1 mg daily rasagiline tablets treatment, in which most patients did not follow dietary tyramine restriction.

There have been postmarketing reports of patients who experienced significantly elevated blood pressure (including rare cases of hypertensive crisis) after ingestion of unknown amounts of tyramine- rich foods while taking recommended doses of rasagiline tablets. Patients should be advised to avoid foods containing a very large amount of tyramine while taking recommended doses of rasagiline tablets [see *Warnings and Precautions (5.1)*].

## 7.8 Dopaminergic Antagonists

It is possible that dopamine antagonists, such as antipsychotics or metoclopramide, could diminish the effectiveness of rasagiline tablets.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

There are no adequate data on the developmental risks associated with the use of rasagiline tablets in pregnant women. In animal studies, oral administration of rasagiline to rats during gestation and lactation resulted in decreased survival and reduced body weight in the offspring at doses similar to those used clinically. When administered to pregnant animals in combination with levodopa/carbidopa, there were increased incidences of fetal skeletal variations in rats and increases in embryofetal death and cardiovascular abnormalities in rabbits [see Data].

In the US general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively. The background risks of major birth defects and miscarriage for the indicated population is unknown.

#### Data

##### Animal Data

In a combined mating/fertility and embryofetal development study in pregnant rats, no effect on embryo-fetal development was observed at oral doses up to 3 mg/kg/day (approximately 30 times the plasma exposure (AUC) in humans at the maximum recommended human dose [MRHD, 1 mg/day]).

In pregnant rabbits administered rasagiline throughout the period of organogenesis at oral doses of up to 36 mg/kg/day, no developmental toxicity was observed. At the highest dose tested, the plasma AUC was approximately 800 times that in humans at the MRHD.

In pregnant rats administered rasagiline (0, 0.1, 0.3, 1 mg/kg/day) orally during gestation and lactation, offspring survival was decreased and offspring body weight was reduced at 0.3 mg/kg/day and 1 mg/kg/day (10 and 16 times the plasma AUC in humans at the MRHD). The no-effect dose (0.1 mg/kg) for adverse developmental effects is similar to the MRHD on a body surface area (mg/m<sup>2</sup>) basis. The effect of rasagiline on physical and behavioral development was not adequately assessed in this study.

Rasagiline may be given as an adjunct therapy to levodopa/carbidopa treatment. In pregnant rats administered rasagiline (0, 0.1, 0.3, 1 mg/kg/day) and levodopa/carbidopa (80/20 mg/kg/day)(alone and in combination orally throughout the period of organogenesis, there was an increased incidence of fetal skeletal variations in fetuses from rats treated with rasagiline in combination with levodopa/carbidopa at 1/80/20 mg/kg/day (approximately 8 times the rasagiline plasma AUC in humans at the MRHD and similar to the MRHD of levodopa/carbidopa [800/200 mg/day] on a mg/m<sup>2</sup> basis). In pregnant rabbits dosed orally throughout the period of organogenesis with rasagiline alone (3 mg/kg) or in combination with levodopa/carbidopa (rasagiline: 0.1, 0.6, 1.2 mg/kg, levodopa/carbidopa: 80/20 mg/kg/day), an increase in embryofetal death was noted at rasagiline doses of 0.6 and 1.2 mg/kg/day when administered in combination with levodopa/carbidopa (approximately 7 and 13 times, respectively, the rasagiline plasma AUC in humans at the MRHD). There was an increase in cardiovascular abnormalities with levodopa/carbidopa alone (similar to the MRHD on a mg/m<sup>2</sup> basis) and to a greater extent when rasagiline(at all doses; 1-13 times the rasagiline plasma AUC in humans at the MRHD)was administered in combination with levodopa/carbidopa.

### 8.2 Lactation

### Risk Summary

There are no data on the presence of rasagiline in human milk or the effects on the breastfed infant. In rats, rasagiline was shown to inhibit prolactin secretion. The clinical relevance in humans is unknown, and there are no data on the effects of rasagiline on prolactin secretion or milk production in humans.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for rasagiline tablets and any potential adverse effects on the breastfed infant from rasagiline tablets or from the underlying maternal condition.

### **8.4 Pediatric Use**

The safety and effectiveness in pediatric patients have not been established.

### **8.5 Geriatric Use**

Approximately half of patients in clinical trials were 65 years and over. There were no significant differences in the safety profile of the geriatric and nongeriatric patients.

### **8.6 Hepatic Impairment**

Rasagiline plasma concentration may be increased in patients with mild (up to 2 fold, Child-Pugh score 5-6), moderate (up to 7 fold, Child-Pugh score 7-9), and severe (Child-Pugh score 10-15) hepatic impairment. Patients with mild hepatic impairment should not exceed a dose of 0.5 mg/day. Rasagiline tablets should not be used in patients with moderate or severe hepatic impairment [see *Dosage and Administration (2.3)*, *Warnings and Precautions (5.5)* and *Clinical Pharmacology (12.3)*].

### **8.7 Renal Impairment**

Dose adjustment of rasagiline tablets are not required for patients with mild or moderate renal impairment because rasagiline tablets plasma concentrations are not increased in patients with moderate renal impairment. Rasagiline has not been studied in patients with severe renal impairment [see *Clinical Pharmacology (12.3)*].

## **9 DRUG ABUSE AND DEPENDENCE**

### **9.1 Controlled Substance**

Rasagiline is not a controlled substance.

### **9.2 Abuse**

Studies conducted in mice and rats did not reveal any potential for drug abuse and dependence. Clinical trials have not revealed any evidence of the potential for abuse, tolerance or physical dependence; however, systematic studies in humans designed to evaluate these effects have not been performed.

### **9.3 Dependence**

Studies conducted in mice and rats did not reveal any potential for drug abuse and dependence. Clinical trials have not revealed any evidence of the potential for abuse, tolerance or physical dependence; however, systematic studies in humans designed to evaluate these effects have not been performed.

## **10 OVERDOSAGE**

In a dose escalation study in patients on chronic levodopa therapy treated with 10 mg of rasagiline tablets there were three reports of cardiovascular side effects (including hypertension and postural hypotension) which resolved following treatment discontinuation.

Although no cases of overdose have been observed with rasagiline tablets during the clinical development program, the following description of presenting symptoms and clinical course is based upon overdose descriptions of nonselective MAOI inhibitors. The signs and symptoms of nonselective MAOI overdose may not appear immediately. Delays of up to 12 hours after ingestion of drug and the appearance of signs may occur. The peak intensity of the syndrome may not be reached until for a day following the overdose. Death has been reported following overdose; therefore, immediate hospitalization, with continuous patient observation and monitoring for at least two days following the ingestion of such drugs in overdose, is strongly recommended.

The severity of the clinical signs and symptoms of MAOI overdose varies and may be related to the amount of drug consumed. The central nervous and cardiovascular systems are prominently involved. Signs and symptoms of MAOI overdose may include:

drowsiness, dizziness, faintness, irritability, hyperactivity, agitation, severe headache, hallucinations, trismus, opisthotonos, convulsions, and coma; rapid and irregular pulse, hypertension, hypotension and vascular collapse; precordial pain, respiratory depression and failure, hyperpyrexia, diaphoresis, and cool, clammy skin.

There is no specific antidote for rasagiline tablets overdose. The following suggestions are offered based upon the assumption that rasagiline tablets overdose may be modeled after nonselective MAO inhibitor poisoning. Treatment of overdose with nonselective MAO inhibitors is symptomatic and supportive. Respiration should be supported by appropriate measures, including management of the airway, use of supplemental oxygen, and mechanical ventilatory assistance, as required. Body temperature should be monitored closely. Intensive management of hyperpyrexia may be required.

Maintenance of fluid and electrolyte balance is essential. For this reason, in cases of overdose with rasagiline tablets, dietary tyramine restriction should be observed for several weeks to reduce the risk of hypertensive tyramine reaction.


A poison control center should be called for the most current treatment guidelines.

A postmarketing report described a single patient who developed a nonfatal serotonin syndrome after ingesting 100 mg of rasagiline tablets in a suicide attempt. Another patient who was treated in error with 4 mg rasagiline tablets daily and tramadol also developed a serotonin syndrome. One patient who was treated in error with 3 mg rasagiline tablets daily experienced alternating episodes of vascular fluctuations consisting of hypertension and orthostatic hypotension.

## 11 DESCRIPTION

Rasagiline tablets contain rasagiline (as the mesylate), a propargylamine-based drug indicated for the treatment of idiopathic Parkinson's disease. Rasagiline mesylate is designated chemically as: 1H-Inden-1-amine, 2, 3-dihydro-N-2-propynyl-, (1R)-, methanesulfonate. The empirical formula of rasagiline mesylate is  $C_{12}H_{13}N$ .  $CH_4SO_3$  and its molecular weight is 267.34.

Its structural formula is:



Rasagiline mesylate is a white to off-white powder, freely soluble in water or ethanol and sparingly soluble in isopropanol. Each rasagiline tablets for oral administration contains 0.5 mg or 1 mg of rasagiline (equivalent to 0.78 mg or 1.56 mg of rasagiline mesylate).

Each rasagiline tablet also contains the following inactive ingredients: microcrystalline cellulose, corn starch, colloidal silicon dioxide, anhydrous citric acid, pregelatinized starch, talc and stearic acid.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Rasagiline tablets are a selective, irreversible MAO-B inhibitor indicated for the treatment of idiopathic Parkinson's disease. The results of a clinical trial designed to examine the effects of rasagiline tablets on blood pressure when it is administered with increasing doses of tyramine indicates the functional selectivity can be incomplete when healthy subjects ingest large amounts of tyramine while receiving recommended doses of rasagiline tablets. The selectivity for inhibiting MAO-B diminishes in a dose-related manner.

MAO, a flavin-containing enzyme, is classified into two major molecular species, A and B, and is localized in mitochondrial membranes throughout the body in nerve terminals, brain, liver and intestinal mucosa. MAO regulates the metabolic degradation of catecholamines and serotonin in the CNS and peripheral tissues. MAO-B is the major form in the human brain. In ex vivo animal studies in brain, liver, and intestinal tissues, rasagiline was shown to be a potent, irreversible monoamine oxidase type B (MAO-B) selective inhibitor. Rasagiline at the recommended therapeutic dose was also shown to be a potent and irreversible inhibitor of MAO-B in platelets. The precise mechanisms of action of rasagiline are unknown. One mechanism is believed to be related to its MAO-B inhibitory activity, which causes an increase in extracellular levels of dopamine in the striatum. The elevated dopamine level and subsequent increased dopaminergic activity

are likely to mediate rasagiline's beneficial effects seen in models of dopaminergic motor dysfunction.

## 12.2 Pharmacodynamics

### Tyramine Challenge Test

Results of a tyramine challenge study indicate that rasagiline at recommended doses is relatively selective for inhibiting MAO-B and can be used without dietary tyramine restriction. However, certain foods (e.g., aged cheeses, such as Stilton cheese) may contain very high amounts of tyramine (i.e., 150 mg or greater) and could potentially cause severe hypertension caused by tyramine interaction in patients taking rasagiline tablets due to mild increased sensitivity to tyramine at recommended doses. Relative selectivity of rasagiline tablets for inhibiting MAO-B diminished in a dose-related manner as the dose progressively increased above the highest recommended daily dose (1 mg) [see *Warnings and Precautions (5.1) and Drug Interactions (7.7)*].

### Platelet MAO Activity in Clinical Studies

Studies in healthy subjects and in Parkinson's disease patients have shown that rasagiline inhibits platelet MAO-B irreversibly. The inhibition lasts at least 1 week after last dose. Almost 25-35% MAO-B inhibition was achieved after a single rasagiline dose of 1 mg/day and more than 55% of MAO-B inhibition was achieved after a single rasagiline dose of 2 mg/day. Over 90% inhibition was achieved 3 days after rasagiline daily dosing at 2 mg/day and this inhibition level was maintained 3 days postdose. Multiple doses of rasagiline of 0.5, 1 and 2 mg per day resulted in complete MAO-B inhibition.

## 12.3 Pharmacokinetics

Rasagiline in the range of 1 mg to 6 mg demonstrated a more than proportional increase in AUC, while  $C_{max}$  was dose proportional. Rasagiline mean steady-state half life is 3 hours but there is no correlation of pharmacokinetics with its pharmacological effect because of its irreversible inhibition of MAO-B.

### Absorption

Rasagiline is rapidly absorbed, reaching peak plasma concentration ( $C_{max}$ ) in approximately 1 hour. The absolute bioavailability of rasagiline is about 36%.

Food does not affect the  $T_{max}$  of rasagiline, although  $C_{max}$  and exposure (AUC) are decreased by approximately 60% and 20%, respectively, when the drug is taken with a high fat meal. Because AUC is not significantly affected, rasagiline tablets can be administered with or without food.

### Distribution

The mean volume of distribution at steady-state is 87 L, indicating that the tissue binding of rasagiline is in excess of plasma protein binding. Plasma protein binding ranges from 88-94% with mean extent of binding of 61-63% to human albumin over the concentration range of 1-100 ng/mL.

### Metabolism and Elimination

Rasagiline undergoes almost complete biotransformation in the liver prior to excretion. The metabolism of rasagiline proceeds through two main pathways: N-dealkylation and/or hydroxylation to yield 1-aminoindan (AI), 3-hydroxy-N-propargyl-1 aminoindan (3-OH-PAI) and 3-hydroxy-1-aminoindan (3-OH-AI). In vitro experiments indicate that both routes of rasagiline metabolism are dependent on the cytochrome P450 (CYP) system, with CYP1A2 being the major isoenzyme involved in rasagiline metabolism. Glucuronide conjugation of rasagiline and its metabolites, with subsequent urinary excretion, is the major elimination pathway.

After oral administration of <sup>14</sup>C-labeled rasagiline, elimination occurred primarily via urine and secondarily via feces (62% of total dose in urine and 7% of total dose in feces over 7 days), with a total calculated recovery of 84% of the dose over a period of 38 days. Less than 1% of rasagiline was excreted as unchanged drug in urine.

### Specific Populations

#### *Hepatic Impairment*

Following repeat dose administration (7 days) of rasagiline (1 mg/day) in subjects with mild hepatic impairment (Child-Pugh score 5-6), AUC and  $C_{max}$  were increased by 2 fold and 1.4 fold, respectively, compared to healthy subjects. In subjects with moderate

hepatic impairment (Child-Pugh score 7-9), AUC and  $C_{max}$  were increased by 7 fold and 2 fold, respectively, compared to healthy subjects [see *Dosage and Administration (2.3)* and *Warnings and Precautions (5.5)*].

#### *Renal Impairment*

Following repeat dose administration (8 days) of rasagiline (1 mg/day) in subjects with moderate renal impairment, rasagiline exposure (AUC) was similar to rasagiline exposure in healthy subjects, while the major metabolite 1-AI exposure (AUC) was increased 1.5-fold in subjects with moderate renal impairment, compared to healthy subjects. Because 1-AI is not an MAO inhibitor, no dose adjustment is needed for patients with mild and moderate renal impairment. Data are not available for patients with severe renal impairment.

#### *Elderly*

Since age has little influence on rasagiline pharmacokinetics, it can be administered at the recommended dose in the elderly (> 65 years).

#### *Pediatric*

Rasagiline tablets have not been investigated in patients below 18 years of age.

#### *Gender*

The pharmacokinetic profile of rasagiline is similar in men and women.

#### Drug-Drug Interactions

##### *Levodopa*

A study in Parkinson's disease patients, in which the effect of levodopa/carbidopa (LD/CD) on rasagiline pharmacokinetics at steady state was investigated, showed that the pharmacokinetics of rasagiline was not affected by concomitant administration of LD/CD.

##### *Effect of Other Drugs on the Metabolism of Rasagiline Tablets*

In vitro metabolism studies showed that CYP1A2 was the major enzyme responsible for the metabolism of rasagiline tablets. There is the potential for inhibitors of this enzyme to alter rasagiline clearance when coadministered [see *Dosage and Administration (2.2)* and *Warnings and Precautions (5.4)*].

Ciprofloxacin: When ciprofloxacin, an inhibitor of CYP1A2, was administered to healthy volunteers (n=12) at 500 mg (BID) with rasagiline at 2 mg/day, the AUC of rasagiline increased by 83% and there was no change in the elimination half life [see *Dosage and Administration (2.2)* and *Warnings and Precautions (5.4)*].

Theophylline: Coadministration of rasagiline 1 mg/day and theophylline, a substrate of CYP1A2, up to 500 mg twice daily to healthy subjects (n=24) did not affect the pharmacokinetics of either drug.

Antidepressants: Severe CNS toxicity (occasionally fatal) associated with hyperpyrexia as part of a serotonin syndrome, has been reported with combined treatment of an antidepressant (e.g., from one of many classes including tricyclic or tetracyclic antidepressants, SSRIs, SNRIs, triazolopyridine antidepressants) and nonselective MAOI or a selective MAO-B inhibitor [see *Warnings and Precautions (5.2)*].

##### *Effect of Rasagiline Tablets on Other Drugs*

No additional in vivo trials have investigated the effect of rasagiline tablets on other drugs metabolized by the cytochrome P450 enzyme system. In vitro studies showed that rasagiline at a concentration of 1 mcg/mL (equivalent to a level that is 160 times the average  $C_{max}$  ~ 5.9-8.5 ng/mL in Parkinson's disease patients after 1 mg rasagiline multiple dosing) did not inhibit cytochrome P450 isoenzymes, CYP1A2, CYP2A6, CYP2C9, CYP2C19, CYP2D6, CYP2E1, CYP3A4 and CYP4A. These results indicate that rasagiline is unlikely to cause any clinically significant interference with substrates of these enzymes.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis & Mutagenesis & Impairment of Fertility**

#### **Carcinogenesis**

Two-year carcinogenicity studies were conducted in mice at oral doses of 0, 1, 15, and 45 mg/kg/day and in rats at oral doses of 0.3, 1, and 3 mg/kg/day (males) or 0, 0.5, 2, 5, and 17 mg/kg/day (females). In rats, there was no increase in tumors at any dose tested. Plasma exposures (AUC) at the highest dose tested were approximately 33 and 260 times, in male and female rats, respectively, that in humans at the maximum recommended human dose (MRHD) of 1 mg/day.

In mice, there was an increase in lung tumors (combined adenomas/carcinomas) at 15 and 45 mg/kg in males and females. At the lowest dose tested, plasma AUCs were approximately 5 times those expected in humans at the MRHD.

The carcinogenic potential of rasagiline administered in combination with levodopa/carbidopa has not been examined.

#### Mutagenesis

Rasagiline was reproducibly clastogenic in in vitro chromosomal aberration assays in human lymphocytes in the presence of metabolic activation and was mutagenic and clastogenic in the in vitro mouse lymphoma tk assay in the absence and presence of metabolic activation. Rasagiline was negative in the in vitro bacterial reverse mutation (Ames) assay and in the in vivo micronucleus assay in mice. Rasagiline was also negative in the in vivo micronucleus assay in mice when administered in combination with levodopa/carbidopa.

#### Impairment of Fertility

Rasagiline had no effect on mating performance or fertility in rats treated prior to and throughout the mating period and continuing in females through gestation day 17 at oral doses of up to 3 mg/kg/day (approximately 30 times the plasma AUC in humans at the MRHD). The effect of rasagiline administered in combination with levodopa/carbidopa on mating and fertility has not been examined.

### 14 CLINICAL STUDIES

The effectiveness of rasagiline tablets for the treatment of Parkinson’s disease was established in four 18-to 26-week, randomized, placebo-controlled trials, as initial monotherapy or adjunct therapy.

#### 14.1 Monotherapy Use of Rasagiline Tablets

Study 1 was a double-blind, randomized, fixed-dose parallel group, 26-week study in early Parkinson’s disease patients not receiving any concomitant dopaminergic therapy at the start of the study. The majority of the patients were not treated with medications for Parkinson’s disease before receiving rasagiline tablets.

In Study 1, 404 patients were randomly assigned to receive placebo (138 patients), rasagiline tablets 1 mg/day (134 patients) or rasagiline tablets 2 mg/day (132 patients). Patients were not allowed to take levodopa, dopamine agonists, selegiline or amantadine, but could take stable doses of anticholinergic medication, if necessary. The average Parkinson’s disease duration was approximately 1 year (range 0 to 11 years).

The primary measure of effectiveness was the change from baseline in the total score of the Unified Parkinson’s Disease Rating Scale (UPDRS), [mentation (Part I) + activities of daily living (ADL) (Part II) + motor function (Part III)]. The UPDRS is a multi-item rating scale that measures the ability of a patient to perform mental and motor tasks as well as activities of daily living. A reduction in the score represents improvement and a beneficial change from baseline appears as a negative number.

Rasagiline tablets (1 or 2 mg once daily) were superior to placebo on the primary measure of effectiveness in patients receiving six months of treatment and not on dopaminergic therapy. The effectiveness of rasagiline tablets 1 mg and 2 mg was comparable. Table 4 shows the results of Study 1. There were no differences in effectiveness based on age or gender between rasagiline tablets 1 mg/day and placebo.

**Table 4: Change in Total UPDRS Score in Study 1**

	<b>Baseline score</b>	<b>Change from baseline to termination score</b>	<b>p-value vs. placebo</b>
Placebo	24.5	3.9	---
Rasagiline tablets 1 mg	24.7	0.1	0.0001
Rasagiline tablets 2 mg	25.9	0.7	0.0001

#### 14.2 Adjunct Use of Rasagiline tablets

Study 2 was a double-blind, randomized, placebo-controlled, parallel group, 18-week study, investigating rasagiline tablets 1 mg as adjunct therapy to dopamine agonists without levodopa. Patients were on a stable dose of dopamine agonist (ropinirole, mean 8 mg/day or pramipexole, mean 1.5 mg/day) therapy for ≥ 30 days, but at doses not



sufficient to control Parkinson's disease symptoms.

In Study 2, 321 patients randomly received placebo (162 patients) or rasagiline tablets 1 mg/day (159 patients) and had a post-baseline assessment. The average Parkinson's disease duration was approximately 2 years (range 0.1 to 14.5 years). The primary measure of effectiveness was the change from baseline in the total score of the Unified Parkinson's Disease Rating Scale (UPDRS) [mentation (Part I) + activities of daily living (ADL) (Part II) + motor function (Part III)]. In Study 2, rasagiline tablets 1 mg was superior to placebo on the primary measure of effectiveness (see Table 5).

**Table 5: Change in Total UPDRS Score in Study 2**

	<b>Baseline score</b>	<b>Change from baseline to termination score*</b>	<b>p-value vs. placebo</b>
Placebo	29.8	-1.2	-----
Rasagiline tablets 1 mg	32.1	-3.6	0.012

\*A negative change from baseline indicates improvement in the UPDRS Secondary outcome assessment of the individual subscales of the UPDRS indicates that the UPDRS Part III motor subscale was primarily responsible for the overall rasagiline tablets effect on the UPDRS score (see Table 6).

**Table 6: Secondary Measures of Effectiveness in Study 2**

	<b>Baseline (score)</b>	<b>Change from baseline to termination score</b>
<b>UPDRS Part II ADL (Activities of Daily Living) subscale score</b>		
Placebo	7.9	0.4
Rasagiline tablets 1 mg	8.6	-0.3
<b>UPDRS Part III Motor subscale score</b>		
Placebo	20.4	-1.2
Rasagiline tablets 1 mg	22.2	-3.7

Study 3 and Study 4 were randomized, multinational trials conducted in more advanced Parkinson's disease patients treated chronically with levodopa and experiencing motor fluctuations (including but not limited to, end of dose "wearing off," sudden or random "off," etc.). Study 3 was conducted in North America (U.S. and Canada) and compared rasagiline tablets 0.5 mg and 1 mg daily to placebo. Study 4 was conducted outside of North America in Europe, Argentina and Israel, and compared rasagiline tablets 1 mg daily to placebo.

Patients had Parkinson's disease for an average of 9 years (range 5 months to 33 years), had taken levodopa for an average of 8 years (range 5 months to 32 years), and had motor fluctuations for approximately 3 to 4 years (range 1 month to 23 years). Patients kept home Parkinson's disease diaries just prior to baseline and at specified intervals during the trial. Diaries recorded one of the following four conditions for each halfhour interval over a 24-hour period: "ON" (period of relatively good function and mobility) as either "ON" with no dyskinesia or without troublesome dyskinesia, or "ON" with troublesome dyskinesia, "OFF" (period of relatively poor function and mobility) or asleep. "Troublesome" dyskinesia is defined as dyskinesia that interferes with the patient's daily activity. All patients had inadequate control of their motor symptoms with motor fluctuations typical of advanced stage disease despite receiving levodopa/decarboxylase inhibitor. The average dose of levodopa taken with a decarboxylase inhibitor was approximately 700 to 800 mg (range 150 to 3000 mg/day). Patients continued their stable doses of additional anti-PD medications at entry into the trials. Approximately 65% of patients in both studies were also taking a dopamine agonist. In the North American study (Study 3), approximately 35% of patients took

entacapone with levodopa/decarboxylase inhibitor. The majority of patients taking entacapone were also taking a dopamine agonist.

In Study 3 and Study 4, the primary measure of effectiveness was the change in the mean number of hours spent in the "OFF" state at baseline compared to the mean number of hours spent in the "OFF" state during the treatment period.

In Study 3, patients were randomly assigned to receive placebo (159 patients), rasagiline tablets 0.5 mg/day (164 patients), or rasagiline tablets 1 mg/day (149 patients) for 26 weeks. Patients averaged 6 hours daily in the "OFF" state at baseline as confirmed by home diaries.

In Study 4, patients were randomly assigned to receive placebo (229 patients), rasagiline tablets 1 mg/day (231 patients) or a COMT inhibitor (active comparator), taken along with scheduled doses of levodopa/decarboxylase inhibitor (227 patients) for 18 weeks. Patients averaged 5.6 hours daily in the "OFF" state at baseline as confirmed by home diaries.

In Study 3 and Study 4, rasagiline tablets 1 mg once daily reduced "OFF" time compared to placebo when added to levodopa in patients experiencing motor fluctuations (Tables 7 and 8). The lower dose (0.5 mg) of rasagiline tablets also significantly reduced "OFF" time (Table 7), but had a numerically smaller effect than the 1 mg dose of rasagiline tablets. In Study 4, the active comparator also reduced "OFF" time when compared to placebo.

**Table 7: Change in mean total daily "OFF" time in Study 3 Table**

	<b>Baseline (hours)</b>	<b>Change from baseline to treatment period (hours)</b>	<b>p-value vs. placebo</b>
Placebo	6.0	-0.9	----
Rasagiline tablets 0.5 mg	6.0	-1.4	0.0199
Rasagiline tablets 1 mg	6.3	-1.9	< 0.0001

**Table 8: Change in mean total daily "OFF" time in Study 4**

	<b>Baseline (hours)</b>	<b>Change from baseline to treatment period (hours)</b>	<b>p-value vs. placebo</b>
Placebo	5.5	-0.40	----
Rasagiline tablets 1 mg	5.6	-1.2	0.0001

In Study 3 and Study 4, dose reduction of levodopa was allowed within the first 6 weeks, if dopaminergic side effects developed including dyskinesia or hallucinations. In Study 3, the levodopa dose was reduced in 8% of patients in the placebo group and in 16% and 17% of patients in the 0.5 mg/day and 1 mg/day rasagiline tablets groups, respectively. When levodopa was reduced, the dose was reduced by 7%, 9%, and 13% in the placebo, 0.5 mg/day, and 1 mg/day groups, respectively. In Study 4, levodopa dose reduction occurred in 6% of patients in the placebo group and in 9% in the rasagiline tablets 1 mg/day groups, respectively. When levodopa was reduced, it was reduced by 13% and 11% in the placebo and the rasagiline tablets groups, respectively. There were no differences in effectiveness based on age or gender between rasagiline tablets 1 mg/day and placebo.

Several secondary outcome assessments in the two studies showed statistically significant improvements with rasagiline. These included effects on the activities of daily living (ADL) subscale of the UPDRS performed during an "OFF" period and the motor subscale of the UPDRS performed during an "ON" period. In both scales, a negative response represents improvement. Tables 9 and 10 show these results for Studies 3 and 4.

**Table 9: Secondary Measures of Effectiveness in Study 3**

	<b>Baseline (score)</b>	<b>Change from baseline to last value</b>
<b>UPDRS ADL (Activities of Daily Living) subscale score while "OFF"</b>		
Placebo	15.5	0.68
Rasagiline tablets 0.5 mg	15.8	-0.60
Rasagiline tablets 1 mg	15.5	-0.68
<b>UPDRS Motor subscale score while "ON"</b>		

Placebo	20.8	1.21
Rasagiline tablets 0.5 mg	21.5	-1.43
Rasagiline tablets 1 mg	20.9	-1.30

**Table 10: Secondary Measures of Effectiveness in Study 4**

	Baseline (score)	Change from baseline to last value
<b>UPDRS ADL (Activities of Daily Living) subscale score while "OFF"</b>		
Placebo	18.7	-0.89
Rasagiline tablets 1 mg	19.0	-2.61
<b>UPDRS Motor subscale score while "ON"</b>		
Placebo	23.5	-0.82
Rasagiline tablets 1 mg	23.8	-3.87

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Rasagiline is supplied as round, flat, beveled tablets, available in bottles and unit dose blister packages as follows:

Each 0.5 mg tablet is white to off-white, round, flat, beveled tablets, debossed with "C12" on one side and plain on the other side and is available as follows:

Bottle of 30 tablets                      NDC 33342-308-07

Each 1 mg tablet is white to off-white, round, flat, beveled tablets, debossed with "C13" on one side and plain on the other side and is available as follows:

Bottle of 30 tablets                      NDC 33342-309-07

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

## 17 PATIENT COUNSELING INFORMATION

### Hypertension

Advise patients that treatment with recommended doses of rasagiline tablets may be associated with elevations of blood pressure. Tell patients who experience elevation of blood pressure while taking rasagiline tablets to contact their healthcare provider. The risk of using higher than recommended daily doses of rasagiline tablets should be explained, and a brief description of the tyramine associated hypertensive reaction provided.

Advise patients to avoid certain foods (e.g., aged cheese) containing a very large amount of tyramine while taking recommended doses of rasagiline tablets because of the potential for large increases in blood pressure. If patients eat foods very rich in tyramine and do not feel well soon after eating, they should contact their healthcare provider [see *Warnings and Precautions (5.1)*].

### Serotonin Syndrome

Tell patients to inform their physician if they are taking, or planning to take, any prescription or over-the-counter drugs, especially antidepressants and over-the-counter cold medications, since there is a potential for interaction with rasagiline tablets. Because patients should not use meperidine or certain other analgesics with rasagiline tablets, they should contact their healthcare provider before taking analgesics [see *Contraindications (4) and Warnings and Precautions (5.2)*].

### Falling Asleep During Activities of Daily Living and Somnolence

Advise and alert patients about the potential for sedating effects associated with rasagiline tablets and other dopaminergic medications, including somnolence and particularly to the possibility of falling asleep while engaged in activities of daily living. Because somnolence can be a frequent adverse reaction with potentially serious consequences, patients should neither drive a car nor engage in other potentially dangerous activities until they have gained sufficient experience with rasagiline tablets and other dopaminergic medications to gauge whether or not it affects their mental and/or motor performance adversely. Advise patients that if increased somnolence or

new episodes of falling asleep during activities of daily living (e.g., watching television, passenger in a car, etc.) are experienced at any time during treatment, they should not drive or participate in potentially dangerous activities until they have contacted their physician. Patients should not drive, operate machinery, or work at heights during treatment if they have previously experienced somnolence and/or have fallen asleep without warning prior to use of rasagiline tablets.

Because of possible additive effects, advise patients to exercise caution when patients are taking other sedating medications, alcohol, or other central nervous system depressants (e.g., benzodiazepines, antipsychotics, antidepressants) in combination with rasagiline tablets or when taking concomitant medications that increase plasma levels of rasagiline (e.g., ciprofloxacin)[see *Warnings and Precautions (5.3)*].

### **Ciprofloxacin or Other CYP1A2 Inhibitors**

Inform patients that they should contact their healthcare provider of rasagiline tablets if they take ciprofloxacin or a similar drug that could increase blood levels of rasagiline because of the need to adjust the dose of rasagiline tablets [see *Dosage and Administration (2.2) and Warnings and Precautions (5.4)*].

### **Hepatic Impairment**

Tell patients who have hepatic problems to contact their healthcare provider regarding possible changes in rasagiline tablets dosing [see *Warnings and Precautions (5.5)*].

### **Dyskinesia**

Advise patients taking rasagiline tablets as adjunct to levodopa that there is a possibility of dyskinesia or increased dyskinesia [see *Warnings and Precautions (5.7)*].

### **Hallucinations / Psychotic-Like Behavior**

Inform patients that hallucinations or other manifestations of psychotic-like behavior can occur when taking rasagiline tablets. Advise patients that, if they have a major psychotic disorder, that rasagiline tablets should not ordinarily be used because of the risk of exacerbating the psychosis. Patients with a major psychotic disorder should also be aware that many treatments for psychosis may decrease the effectiveness of rasagiline tablets [see *Warnings and Precautions (5.8)*].

### **Impulse Control/Compulsive Behaviors**

Advise patients that they may experience intense urges to gamble, increased sexual urges, other intense urges, and the inability to control these urges while taking one or more of the medications that increase central dopaminergic tone and that are generally used for the treatment of Parkinson's disease (including rasagiline tablets). Although it is not proven that the medications caused these events, these urges were reported to have stopped in some cases when the dose was reduced or the medication was stopped. Prescribers should ask patients about the development of new or increased gambling urges, sexual urges, or other urges while being treated with rasagiline tablets. Patients should inform their physician if they experience new or increased gambling urges, increased sexual urges, or other intense urges while taking rasagiline tablets. Physicians should consider dose reduction or stopping the medication if a patient develops such urges while taking rasagiline tablets [see *Warnings and Precautions 5.9*].

### **Withdrawal-Emergent Hyperpyrexia and Confusion**

Tell patients to contact their healthcare provider if they wish to discontinue rasagiline tablets [see *Warnings and Precautions (5.10)*].

### **Missing Dose**

Instruct patients to take rasagiline tablets as prescribed. If a dose is missed, the patient should not double-up the dose of rasagiline tablets. The next dose should be taken at the usual time on the following day.

### **Pregnancy**

Advise patients to notify their healthcare provider if they are pregnant or plan to become pregnant [see *Use in Specific Populations (8.1)*].

Manufactured for:

Macleods Pharma USA, Inc.  
Princeton, NJ 08540.

Manufactured by:

Macleods Pharmaceuticals Ltd.  
At Oxalis Labs  
Baddi, Himachal Pradesh, INDIA

Revised: July 2024

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

Rasagiline Tablets, 0.5 mg  
 NDC 33342-308-07  
 Bottles of 30 tablets

Rasagiline Tablets, 1 mg  
 NDC 33342-309-07  
 Bottles of 30 tablets

**RASAGILINE**

rasagiline tablet

Product Information			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:33342-308
<b>Route of Administration</b>	ORAL		
Active Ingredient/Active Moiety			
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
	RASAGILINE MESYLATE (UNII: LH8C2J1290) (RASAGILINE - UNII:003N66TS6T)	RASAGILINE	0.5 mg
Inactive Ingredients			
	<b>Ingredient Name</b>		<b>Strength</b>
	CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)		
	STARCH, CORN (UNII: O8232NY35J)		
	ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
	TALC (UNII: 7SEV7J4R1U)		
	SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
	STEARIC ACID (UNII: 4ELV7Z65AP)		
Product Characteristics			
<b>Color</b>	WHITE (white to off-white)	<b>Score</b>	no score
<b>Shape</b>	ROUND (Flat faced, Beveled edged,Circular shaped)	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	C;12
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:33342-308-07	30 in 1 BOTTLE; Type 0: Not a Combination Product	06/29/2024	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA208866	06/29/2024	

**RASAGILINE**  
rasagiline tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:33342-309
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
RASAGILINE MESYLATE (UNII: LH8C2J1290) (RASAGILINE - UNII:003N66TS6T)	RASAGILINE	1 mg

Inactive Ingredients	
Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3S)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics			
Color	WHITE (white to off-white)	Score	no score
Shape	ROUND (Flat faced, Beveled edged,Circular shaped)	Size	8mm
Flavor		Imprint Code	C;13
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:33342-309-07	30 in 1 BOTTLE; Type 0: Not a Combination Product	06/29/2024	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA208866	06/29/2024	

**Labeler** - Macleods Pharmaceuticals Limited (862128535)

Establishment			
Name	Address	ID/FEI	Business Operations
Macleods Pharmaceuticals Limited		676369519	ANALYSIS(33342-308, 33342-309) , LABEL(33342-308, 33342-309) , MANUFACTURE(33342-308, 33342-309) , PACK(33342-308, 33342-309)