

# **THOMPSON XS PAIN ACHE- aspirin, caffeine tablet, film coated**

## **Thompson Safety**

-----

### **Thompson XS Pain Relief**

#### ***Drug Facts***

#### ***Active ingredients (in each caplet)***

Aspirin 500 mg (NSAID)\*

Caffeine 32.5 mg

\*nonsteroidal anti-inflammatory drug

#### ***Purpose***

Pain reliever

Pain reliever aid

#### ***Uses***

for the temporary relief of

- backache pain
- muscle aches and pains
- minor aches and pains of arthritis

#### ***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

#### **Allergy alert:**

Aspirin may cause a severe allergic reaction which may include:

- hives
- shock
- facial swelling
- asthma (wheezing)

#### **Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcohol drinks every day while using this product

- have had stomach ulcers or bleeding problems

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer.

- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you

- you have a history of stomach problems, such as heartburn

- you have asthma

- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout

- diabetes

- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.

- you experience any of the following signs of stomach bleeding: ■ vomit blood ■ have bloody or black stools ■ feel faint ■ have stomach pain that does not get better

- ringing in the ears or a loss of hearing occurs

- pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days

- redness or swelling is present

- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- **drink a full glass of water with each dose**

**Adults and children (12 years and over):** take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours

**Children under 12 years:** do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- tamper evident sealed packets
- do not use any opened or torn packets

**Inactive ingredients**

corn starch, hypromellose, maltodextrin, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate

**Questions or comments?** 877.506.4291

Thompson

Extra Strength Pain & Ache Relief

Pain Relief

Aspirin (NSAID) 500 mg

Pain Reliever Aid

Caffeine 32.5 mg

Safety Works | Business Thrives™

Fast Acting Advanced Formula

Pull to Open

This Package is for Households Without Young Children.

80 Caplets

Tamper Evident Sealed Packets

**EXTRA STRENGTH PAIN & ACHE**  
FAST ACTING ADVANCED FORMULA  
THOMPSON SAFETY WORKS | BUSINESS THRIVES  
PAIN RELIEF  
ASPIRIN (NSAID) 500 MG  
PAIN RELIEVER AID  
CAFFEINE 32.5 MG  
THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.  
80 Caplets TAMPER EVIDENT SEALED PACKETS

**Drug Facts**  
**Active ingredients (in each caplet)** Purpose  
Aspirin (NSAID) 500 mg Pain reliever  
Caffeine 32.5 mg Pain reliever and  
nonsteroidal anti-inflammatory drug  
**Uses**  
For the temporary relief of  
• backache pain  
• muscle aches and pains  
• minor aches and pains of arthritis  
**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:  
• hives  
• rash  
• facial swelling  
• asthma (wheezing)  
**Do not use**  
• if you are allergic to aspirin or any other pain reliever/fever reducer.  
• if you have ever had an allergic reaction to this product or any of its ingredients.  
**Ask a doctor before use if**  
• stomach bleeding warning applies to you  
• you have a history of stomach problems, such as heartburn  
• you have asthma  
• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
• you are taking a diuretic (continued on opposite panel)  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  
• take more or for a longer time than directed  
• take a blood thinner (anticoagulant) or steroid drug  
• are age 60 or older  
• have 3 or more alcohol drinks every day while using this product  
• have had stomach ulcers or bleeding problems  
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
**Stop use and ask a doctor if**  
• an allergic reaction occurs. Seek medical help right away.  
• you experience any of the following signs of stomach bleeding:  
• vomit blood  
• have bloody or black stools  
• feel faint  
• have stomach pain that does not get better  
• ringing in the ears or loss of hearing occurs  
• pain gets worse or lasts more than 10 days  
• fever gets worse or lasts more than 3 days  
• redness or swelling is present  
• new symptoms occur  
These could be signs of a serious condition.  
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 30 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.  
**Directions**  
• drink a full glass of water with each dose  
**Adults and children (12 years and over):** take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours  
**Children under 12 years:** do not use unless directed by a doctor  
**Other information**  
• store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
• tamper evident sealed packets  
• do not use any opened or torn packets  
**Inactive ingredients**  
corn starch, hypromellose, maltodextrin, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate  
**Questions or comments?** 877.506.4291  
Reorder CS92180  
Distributed by Thompson Safety Houston, TX 77073 877.506.4291 www.thompson-safety.com  
73408 92180 2



<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (Caplet)	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	44;413
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-821-80	40 in 1 BOX	09/26/2022	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/26/2022	

**Labeler** - Thompson Safety (080998015)

Revised: 9/2025

Thompson Safety