RITUSSIN EXPECTORANT- guaifenes in liquid Rij Pharmaceutical Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each 5 mL)(one teaspoonful)

Guaifenesin, USP 100 mg

Purpose

Expectorant

Use

helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Warnings

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not take more than 6 doses in any 24-hour period

age	dose
adults and children 12 years of	2-4 teaspoons every
age and over	4 hours
children 6 years to under 12	1-2 teaspoonful
years of age	every 4 hours
children 2 years to under 6	1/2-1 teaspoonful
years of age	every 4 hours
children under 2 years of age	ask a doctor

Other information

- store at room temperature 15° 30°C (59° 86°F)
- protect from freezing.

- protect from light.
- Each teaspoon (5 mL) contains: sodium 2 mg.
- TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF BREAKAWAY BAND ON CAP IS BROKEN OR MISSING

Inactive ingredients

caramel, citric acid, FD&C Red #40, , flavors, glucose, glycerin, high fructose corn syrup, menthol, propylene glycol, sodium saccharin, sodium benzoate, water

PRINCIPAL DISPLAY PANEL



(Guaifenesin Syrup)



This is a bulk package. Dispense contents with a child-resistant closure in a tight, light resistant container as defined in the USP.

ALCOHOL FREE

Children & Adults
Cough Formula

16 Fl. oz. (473 mL)

Drug Facts

Uses • helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Warning

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if

 cough lasts more than 7 days, comes back or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • Do not take more than 6 doses in any 24-hour period.

age	dose
adults and children 12 years of age and over	2-4 teaspoonfuls every 4 hours
children 6 years to under 12 years of age	1-2 teaspoonful every 4 hours
children 2 years to under 6 years of age	1/2-1 teaspoonful every 4 hours
under 2 years of age	ask a doctor

Other information - store at room temperature 15° - 30°C (59° - 86°F)

- protect from freezing protect from light
- Each teaspoon (5 mL) contains: sodium 2 mg.

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.

Inactive ingredients: caramel, citric acid, FD&C Red #40, flavors, glucose, glycerin, high fructose corn syrup, menthol, sodium saccharin, sodium benzoate, water.

†This product is not manufactured or distributed by the owner of the registered trademark Robitussin®.

RIJ PHARMACEUTICAL CORPORATION 40 COMMERCIAL AVENUE

MIDDLETOWN, NY 10941



guaifenesin liquid

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53807-408	
Route of Administration	ORAL			

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)			
CARAMEL (UNII: T9D99G2B1R)			
DEXTROSE (UNII: IY9 XDZ35W2)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)			
MENTHOL (UNII: L7T10EIP3A)			
SACCHARIN SO DIUM (UNII: SB8 ZUX40 TY)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
WATER (UNII: 059QF0KO0R)			
GLYCERIN (UNII: PDC6A3C0OX)			

Product Characteristics				
Color	RED	Score		
Shape		Size		
Flavor	CHERRY	Imprint Code		
Contains				

F	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:53807-408-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/1999		
2	NDC:53807-408-08	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/1999		
3	NDC:53807-408-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/1999		
4	NDC:53807-408-28	3785 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/1999		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	03/16/1999	

Labeler - Rij Pharmaceutical Corporation (144679156)

Establishment			
Name	Address	ID/FEI	Business Operations
Rij Pharmaceutical Corporation		144679156	manufacture(53807-408)

Revised: 4/2018 Rij Pharmaceutical Corporation