# CHILDRENS PAIN AND FEVER- acetaminophen 160mg/5ml solution CalmCo LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### **DrKids Children's Pain and Fever**

#### Active Ingredient (in each 5mL)

Acetaminophen, 160mg

#### Purpose

Pain reliever/fever reducer

#### Uses

Temporarily: • reduces fever • relieves minor aches and pains due to: • the common cold • flu • headache • sore throat • toothache

#### Warnings

**Liver Warning**: This product contains acetaminophen. Severe liver damage may occur if your child takes • more than 5 doses in 24 hours, which is the maximum daily amount for this product • With other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

#### Ask a doctor before use if your child has liver disease

**Ask a doctor or pharmacist before use if** your child is taking the blood thinning drug warfarin.

#### Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

#### Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical even if you do not notice any signs or symptoms.

#### Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

• If your child is allergic to acetaminophen or any of the inactive ingredients in this product

#### Directions

# • this product does not contain directions or complete warnings for adult use.

- do not give more than directed (see overdose warning)
- use the chart below to confirm dose. If possible, use weight to confirm, otherwise use age

• give the recommended number of vials below following the instructions under "Easy to Use" on the side panel.

- empty each Single-Use Vial required of all liquid when dosing
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

#### **Dosing Chart**

Weight (lbs)	Age (Years)	Dose (mL)	
under 24 lbs	under 2 years	Talk to a doctor	
24 -35 lbs	2 - 3 years	1 Single-Use Vial (5 mL)	
36 - 47 lbs	4-5 years	Talk to a doctor	
48 - 71 lbs	6-10 years	2 Single-Use Vial (10 mL)	
72 - 95 lbs	11 years	3 Single-Use Vial (15 mL)	

#### Other information

- each Single-Use vial contains 8mg of sodium
- store at 68-77°F (20-25°C)
- tamper evident: do not use if top flap of carton is open. Do not use if any vials are open or broken

#### Inactive ingredients

citric acid, cherry flavor, edetate disodium, glycerin, maltitol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium metabisulfite, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum

#### **Questions or Comments?**

+1 941-202-3270 or visit www.DrKids.com

### Package



#### 04246 In the USA: Distributed by CalmCo LLC , Sarasota FL 34240 DKPF-04P-001-01 Questions or Comments? 941-202-3270 or visit www.Drkids.com ▲ beunitnoJ stati gunG present. These could be signs of a serious condition. .mup nenthan more than 3 days. Mew symptoms occur. Redness or swelling is sodium metabisulfite, sorbitol, sucralose, trisodium citrate dihydrate, Pain gets worse or last more than 5 days. Fever gets worse or last polyethylene glycol, propylene glycol, purified water, sodium benzoate, Stop use and ask doctor it: citric acid, cherry flavor, edetate disodium, glycerin, maltitol, stnaibargni avitosni .ninshew gunb gninnint Ask a doctor or pharmacist before use if your child is taking the blood any vials are open or broken. Tamper evident: do not use if top flap of carton is open. Do not use if Ask a doctor before use if your child has liver disease. (20-25°C). ■ Each Single-Use Vial contains 8mg of sodium. Store at 68-77°F ingredients in this product. Other Information If your child is allergic to acetaminophen or any of the inactive acetaminophen, ask a doctor or pharmacist. 11 years 'Sdl 26-27 nonprescription). If you are not sure whether a drug contains 6-10 years 'Sql 17-84 with any other drug containing acetaminophen (prescription or 4-5 years 36-47 lbs. Do not use 2-3 years 54-32 lps. vomiting, consult a doctor promptly. under 2 years under 24 Ibs. days, is accompanied or followed by fever, headache, rash, nausea, or Sore throat warning: If sore throat is severe, persists for more than 2 (2169Y) 9pA (sdl) theight If a skin reaction occurs, stop use and seek medical help right away. hed) pnizod Symptoms may include: a skin reddening a blisters a rash Do not give for more than 5 days unless directed by a doctor. Allergy alert: Acetaminophen may cause severe skin reactions. Instant ast. Do not give more than 5 times in 24 hours. acetaminophen. liquid when dosing. If needed, repeat dose every 4 hours while which is the maximum daily amount With other drugs containing above "Easy to Use". Empty each Single Use Vial required of all damage may occur if your child takes: More than 5 doses in 24 hours, recommended number of vials below following the instructions Liver Warning: This product contains acetaminophen. Severe liver possible, use weight to confirm, otherwise use age. Give the spnimeW (see overdose warning). Use the chart below to confirm dose. If warnings for adult use. Do not give more than directed dose due to: I the common cold I flu I headache I sore throat I toothache. This product does not contain directions or complete Uses Temporarily: Reduces fever Relieves minor aches and pains Directions Pain reliever/fever reduce Acetaminophen, 160 mg... symptoms. (јш д цэве ці) medical attention is critical even if you do not notice any signs or Active Ingredient esognu contact a Poison Control Center right away (1-800-222-1222). Prompt Keep out of reach of children. In case of overdose, get medical help or Complete Product Information Stor Facts Keep Outer Package for (beunitnos) **21367** (continued) air bubbles eliminate the vial off the vial to the top of

JSIW

dp

Easy to Use

0.17 fl.oz (5 mL) Each

0.68 fl.oz (20 mL) Total



(Jm Gf) slisiv seu slgnis 6

(Jm 0f) slaiv seu slgnis S

child's mouth

the liquid into

azaanbo

Talk to a doctor (Jm č) lisiv szu signis f

Talk to a doctor

(Jm) 9200



🗸 Alcohol ✓ Dyes

🗸 Sugar

HERP



CHILDRENS PAIN AN acetaminophen 160mg/5ml s				
Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Sourc	e) NDC:73282-2164(M	NDC:70302-126)
Route of Administration	ORAL			
Active Ingredient/Active	Moiety			
Ingre	dient Name		<b>Basis of Strength</b>	Strength
ACETAMINOPHEN (UNII: 36209IT	L9D) (ACETAMINOPHE	N - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 ml
Inactive Ingredients				
	Ingredient Na	ame		Strength
SODIUM METABISULFITE (UNII: 4	4VON5FNS3C)			
WATER (UNII: 059QF0K00R)				
TRISODIUM CITRATE DIHYDRAT	<b>E</b> (UNII: B22547B95K)			
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYLENE GLYCOL (UNII: 6DC9	Q167V3)			
POLYETHYLENE GLYCOL, UNSP	ECIFIED (UNII: 3WJQ0	SDW1A)		
SUCRALOSE (UNII: 96K6UQ3ZD4)				
SORBITOL (UNII: 506T60A25R)				
CITRIC ACID MONOHYDRATE (U	NII: 2968PHW8QP)			
MALTITOL (UNII: D65DG142WK)				
EDETATE DISODIUM (UNII: 7FLD9	1C86K)			
SODIUM BENZOATE (UNII: OJ245	FE5EU)			
XANTHAN GUM (UNII: TTV12P4NE	E)			
Product Characteristics				
Color		Score		
Shape		Size		
Flavor	CHERRY	Imprint Code		

Contains

Pa	ackaging					
#	ltem Code		Package Description		Marketing Start Date	Marketing End Date
	NDC:73282- 2164-1	4 in 1	VIAL, SINGLE-DOSE		09/01/2022	
		5 ml i	n 1 VIAL, SINGLE-DOSE; Type 2: Prefilled Drug Deliv	iverv		
1			/System (syringe, patch, etc.)	i i ci j		
	arketin	Device	/System (syringe, patch, etc.)			
1 M	arketin Marketin Categor	Device	/System (syringe, patch, etc.)	Marke	ting Start Date	Marketing End Date

## Labeler - CalmCo LLC (079308308)

## Registrant - CalmCo LLC (079308308)

Revised: 7/2023

CalmCo LLC