

BUPIVACAINE HYDROCHLORIDE- bupivacaine hydrochloride injection
Brookfield Pharmaceuticals, LLC.

Bupivacaine Hydrochloride in Dextrose Injection, USP

STERILE HYPERBARIC SOLUTION
FOR SPINAL ANESTHESIA

Rx only

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Bupivacaine Hydrochloride in Dextrose Injection, USP safely and effectively. See full prescribing information for Bupivacaine Hydrochloride in Dextrose Injection, USP.

Bupivacaine Hydrochloride in Dextrose Injection, USP for subarachnoid injection

Initial U.S. Approval: 2019

----- **INDICATIONS AND USAGE** -----

Bupivacaine Hydrochloride in Dextrose Injection, USP is an amide-local anesthetic indicated in adults for subarachnoid injection for the production of subarachnoid block (spinal anesthesia). (1)

----- **DOSAGE AND ADMINISTRATION** ---

The dosage of Bupivacaine Hydrochloride in Dextrose Injection, USP administered varies with the anesthetic procedure, the vascularity of the tissues, the number of neuronal segments to be blocked, the depth of anesthesia and degree of muscle relaxation required, the duration of anesthesia desired, individual tolerance, and the physical condition of the patient. Administer the smallest dosage and concentration required to produce the desired result. The following are general dosage guidelines:

- Bupivacaine Hydrochloride in Dextrose Injection, USP 6 mg is generally adequate for vaginal delivery. (2.2)
 - Bupivacaine Hydrochloride in Dextrose Injection, USP 7.5 mg is generally adequate for spinal anesthesia for lower extremity and perineal procedures. (2.2)
- Bupivacaine Hydrochloride in Dextrose Injection, USP 12 mg is generally adequate for lower abdominal procedures. (2.2)
- Bupivacaine Hydrochloride in Dextrose Injection, USP 7.5 mg to 10.5 mg is generally adequate for Cesarean section. (2.2)

----- **DOSAGE FORMS AND STRENGTHS** ---

Injection: 15 mg/2 mL (7.5 mg/mL) in single-dose glass ampules. (3)

----- **CONTRAINDICATIONS** -----

- Intravenous regional anesthesia (Bier Block). (4)
- Septicemia. (4)
- Severe hemorrhage, severe hypotension or shock, and arrhythmias, such as complete heartblock, which severely reduce cardiac output. (4)
- Known hypersensitivity to bupivacaine or to any local anesthetic agent of the amide-type or to other components of Bupivacaine Hydrochloride in Dextrose Injection, USP. (4)
- Local infection at the site of proposed lumbar puncture. (4)

----- WARNINGS AND PRECAUTIONS-----

- Use of Spinal Anesthetics During Uterine Contractions: Spinal anesthetics, including Bupivacaine Hydrochloride in Dextrose Injection, USP, should not be injected during uterine contractions because cerebrospinal fluid current may carry the drug further cephalad than desired, resulting in a high motor block. (5.1)
- Patients with Hypertension: Sympathetic blockade due to spinal anesthesia may result in peripheral vasodilation and hypotension. Monitor blood pressure frequently. Hypotension may be controlled by administration of vasoconstrictor agents in titrated dosages depending on the severity of hypotension and response to treatment. Monitor the onset of adequate spinal anesthesia frequently. (5.2)
- Dose-Related Toxicity: Monitor cardiovascular and respiratory vital signs and patient's state of consciousness after injection of Bupivacaine Hydrochloride in Dextrose Injection, USP. (5.3)
- Risk of Systemic Toxicities with Unintended Intravascular Injection: Unintended intravascular injection may be associated with systemic toxicities, including CNS or cardiorespiratory depression and coma, progressing ultimately to respiratory arrest. Aspirate for blood and cerebrospinal fluid (where applicable) prior to each dose. (5.4)
- Methemoglobinemia: Cases of methemoglobinemia have been reported in association with local anesthetic use. See full prescribing information for more detail on managing these risks. (5.5)
- Risk of Cardiac Arrest with Use of Epidural Bupivacaine in Obstetrical Anesthesia: There have been reports of cardiac arrest during use of Bupivacaine Hydrochloride Injection, USP 0.75% solution for epidural anesthesia in obstetrical patients. Bupivacaine Hydrochloride in Dextrose Injection, USP is recommended for spinal anesthesia in obstetrical patients. (5.6)
- Chondrolysis with Intra-Articular Infusion: Intra-articular infusions of local anesthetics including bupivacaine following arthroscopic and other surgical procedures is an unapproved use, and there have been post-marketing reports of chondrolysis in patients receiving such infusions. (5.7)
- Risk of Cardiac Arrest with Intravenous Regional Anesthesia Use (Bier Block): There have been reports of cardiac arrest and death during the use of bupivacaine for intravenous regional anesthesia (Bier Block). (5.8)

----- ADVERSE REACTIONS -----

Most common adverse reactions are hypotension due to loss of sympathetic tone, and diaphragmatic paralysis or hypoventilation due to cephalad spread and high motor block. (6)

To report SUSPECTED ADVERSE REACTIONS, contact FDA at 1-800-FDA-1088

or www.fda.gov/medwatch.

----- DRUG INTERACTIONS-----

- Local Anesthetics: The toxic effects of local anesthetics are additive. Monitor for neurologic and cardiovascular effects when additional local anesthetics are administered. (7.1)
- Drugs Associated with Methemoglobinemia: Patients are at increased risk of developing methemoglobinemia when concurrently exposed to nitrates, nitrites, local anesthetics, antineoplastic agents, antibiotics, antimalarials, anticonvulsants, and other drugs. (7.2)

----- USE IN SPECIFIC POPULATIONS -----

- Pediatric Use: Administration of Bupivacaine Hydrochloride in Dextrose Injection, USP in patients younger than 18 years is not recommended. (8.4)
- Geriatric Use: Patients 65 years and over, particularly those with hypertension, may be at increased risk for developing hypotension while undergoing spinal anesthesia with Bupivacaine Hydrochloride in Dextrose Injection, USP. (8.5)
- Moderate to Severe Hepatic Impairment: Consider increased monitoring for bupivacaine systemic toxicity. (8.6)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 08/2023

TABLE OF CONTENTS

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Information 2.2 Recommended Dosages of Bupivacaine Hydrochloride in Dextrose Injection, USP

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS 5 WARNINGS AND PRECAUTIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Use of Spinal Anesthetics During Uterine Contractions
- 5.2 Patients with Hypertension
- 5.3 Dose-Related Toxicity
- 5.4 Risk of Systemic Toxicities with Unintended Intravascular Injection
- 5.5 Methemoglobinemia
- 5.6 Risk of Cardiac Arrest with Use of Epidural Bupivacaine in Obstetrical Anesthesia
- 5.7 Chondrolysis with Intra-Articular Infusion
- 5.8 Risk of Cardiac Arrest with Intravenous Regional Anesthesia Use (Bier Block)
- 5.9 Patients with Severe Disturbances of Cardiac Rhythm, Shock, or Heart Block
- 5.10 Risk of Toxicity in Patients with Hepatic Impairment 5.11 Risk of Use in Patients with Impaired Cardiovascular Function

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

7.1 Local Anesthetics 7.2 Drugs Associated with Methemoglobinemia

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment 8.7 Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

1 INDICATIONS AND USAGE

Bupivacaine Hydrochloride in Dextrose Injection, USP is indicated for subarachnoid injection in adults for the production of subarachnoid block (spinal anesthesia).

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Information

- Visually inspect this product for particulate matter and discoloration prior to administration. Bupivacaine Hydrochloride in Dextrose Injection, USP is a clear, colorless solution. Do not administer solutions which are discolored or contain particulate matter.
- Mixing or the prior or intercurrent use of any other local anesthetic with Bupivacaine Hydrochloride in Dextrose Injection, USP is not recommended because of insufficient data on the clinical use of such mixtures.
- Discard unused portions of Bupivacaine Hydrochloride in Dextrose Injection, USP following initial use.

Administration Precautions

- Bupivacaine Hydrochloride in Dextrose Injection, USP is to be administered in carefully adjusted dosages by or under the supervision of experienced clinicians who are well versed in the diagnosis and management of dose-related toxicity and other

acute emergencies which might arise from the block to be employed.

- Use Bupivacaine Hydrochloride in Dextrose Injection, USP only if the following are immediately available: oxygen, cardiopulmonary resuscitative equipment and drugs, and the personnel resources needed for proper management of toxic reactions and related emergencies [see *Warnings and Precautions (5.3)*, *Adverse Reactions (6)*, *Overdosage (10)*].
- The toxic effects of local anesthetics are additive. Monitor for neurologic and cardiovascular effects related to local anesthetic systemic toxicity when additional local anesthetics are administered with Bupivacaine Hydrochloride in Dextrose Injection, USP [see *Warnings and Precautions (5.3)*, *Drug Interactions (7.1)*, *Overdosage (10)*].
- Aspirate for blood and cerebrospinal fluid prior to injecting Bupivacaine Hydrochloride in Dextrose Injection, USP, for both the initial dose and all subsequent doses (where applicable), to avoid intravascular injection and to confirm entry into the subarachnoid space. Aspiration of cerebrospinal fluid into a Bupivacaine Hydrochloride in Dextrose Injection, USP-filled syringe will result in an identifiable swirl in the solution. A negative aspiration for blood does not ensure against an intravascular injection [see *Warnings and Precautions (5.4)*].
- Avoid rapid injection of Bupivacaine Hydrochloride in Dextrose Injection, USP.
- The patient should have an indwelling intravenous catheter to assure adequate intravenous access. The lowest dosage of Bupivacaine Hydrochloride in Dextrose Injection, USP that results in effective spinal anesthesia should be used to avoid a high motor block and serious adverse reactions.
- Perform careful and constant monitoring of cardiovascular and respiratory (adequacy of oxygenation and ventilation) vital signs and the patient's level of consciousness during spinal anesthesia.

Conditions Which May Preclude Use of Spinal Anesthesia

The following conditions may preclude the use of spinal anesthesia, depending upon the physician's evaluation of the patient:

- Pre-existing diseases of the central nervous system (CNS), such as those attributable to pernicious anemia, poliomyelitis, syphilis, or tumor.
- Hematological disorders predisposing to coagulopathies or patients on anticoagulant therapy. Trauma to a blood vessel during the conduct of spinal anesthesia may, in some instances, result in uncontrollable CNS hemorrhage, soft tissue hemorrhage, or development of a hematoma.
- Chronic backache and preoperative headache.
- Hypotension and hypertension.
- Technical problems (persistent paresthesias, persistent bloody tap).
- Arthritis or spinal deformity.
- Extremes of age.
- Psychosis, dementia, or other illnesses resulting in poor patient cooperation.

2.2 Recommended Dosages of Bupivacaine Hydrochloride in Dextrose Injection, USP

The dosage of Bupivacaine Hydrochloride in Dextrose Injection, USP administered varies with the anesthetic procedure, the vascularity of the tissues, the number of neuronal segments to be blocked, the depth of anesthesia and degree of muscle relaxation

required, the duration of anesthesia desired, individual tolerance, and the physical condition of the patient. Administer the smallest dosage and concentration required to produce the desired result.

The extent and degree of spinal anesthesia depend upon several factors including dosage, baricity of the anesthetic solution, volume of solution, force of injection, level of puncture, and position of the patient during and immediately after injection.

In recommended doses, Bupivacaine Hydrochloride in Dextrose Injection, USP produces complete motor and sensory block.

The following table summarizes general dosage guidelines for adult patients for the procedures described:

PROCEDURES	DOSAGE GUIDELINES
Vaginal Delivery	Starting dose, 6 mg (0.8 mL)
Lower Extremity and Perineal Procedures, such as:	
• transurethral resection of the prostate (TURP)	7.5 mg (1 mL)
• vaginal hysterectomy	
Lower Abdominal Procedures, such as:	
• abdominal hysterectomy	12 mg (1.6 mL)
• tubal ligation	
• appendectomy	
Cesarean Section	7.5 mg to 10.5 mg (1 mL to 1.4 mL)

3 DOSAGE FORMS AND STRENGTHS

Bupivacaine Hydrochloride in Dextrose Injection, USP is a clear, colorless solution available as:

- 15 mg/2 mL (7.5 mg/mL) in single-dose glass ampules.

4 CONTRAINDICATIONS

Bupivacaine Hydrochloride in Dextrose Injection, USP is contraindicated in:

- intravenous regional anesthesia (Bier Block) [see *Warnings and Precautions* (5.8)].
- patients with septicemia.
- patients with severe hemorrhage, severe hypotension or shock, due to a reduced cardiac output.
- patients with clinically significant arrhythmias, such as complete heartblock, due a reduced cardiac output.
- patients with a known hypersensitivity to bupivacaine or to any local anesthetic agent of the amide-type or to other components of Bupivacaine Hydrochloride in Dextrose Injection, USP.
- patients with local infection at the site of proposed lumbar puncture.

5 WARNINGS AND PRECAUTIONS

5.1 Use of Spinal Anesthetics During Uterine Contractions

Spinal anesthetics including Bupivacaine Hydrochloride in Dextrose Injection, USP should not be injected during uterine contractions because cerebrospinal fluid current may carry the drug further cephalad than desired, resulting in a high motor block.

5.2 Patients with Hypertension

Sympathetic blockade due to spinal anesthesia may result in peripheral vasodilation and hypotension, the extent of which depends on the number of dermatomes blocked. Patients over 65 years, particularly those with hypertension, may be at increased risk for experiencing the hypotensive effects of Bupivacaine Hydrochloride in Dextrose Injection, USP. Monitor blood pressure frequently, especially in the early phases of anesthesia. Hypotension may be controlled by administration of vasoconstrictor agents in titrated dosages depending on the severity of hypotension and response to treatment. Monitor the onset of adequate spinal anesthesia because it is not always possible to control the level of anesthesia after subarachnoid injection of Bupivacaine Hydrochloride in Dextrose Injection, USP.

5.3 Dose-Related Toxicity

The safety and effectiveness of Bupivacaine Hydrochloride in Dextrose Injection, USP depend on proper dosage, correct technique, adequate precautions, and readiness for emergencies. Careful and constant monitoring of cardiovascular and respiratory (adequacy of oxygenation and ventilation) vital signs and the patient's state of consciousness should be performed after injection of Bupivacaine Hydrochloride in Dextrose Injection, USP solutions.

Possible early warning signs of central nervous system (CNS) toxicity are restlessness, anxiety, incoherent speech, lightheadedness, numbness and tingling of the mouth and lips, metallic taste, tinnitus, dizziness, blurred vision, tremors, twitching, CNS depression, or drowsiness. Delay in proper management of dose-related toxicity, hypoventilation from any cause, and/or altered sensitivity may lead to the development of acidosis, cardiac arrest, and possibly death.

The patient should have an indwelling intravenous catheter to assure adequate intravenous access. Use the lowest dosage of Bupivacaine Hydrochloride in Dextrose Injection, USP that results in effective anesthesia to avoid serious adverse reactions. Avoid rapid injection of a large volume of Bupivacaine Hydrochloride in Dextrose Injection, USP.

Injection of repeated doses of Bupivacaine Hydrochloride in Dextrose Injection, USP may cause significant increases in plasma bupivacaine levels with each repeated dose due to slow accumulation of the drug or its metabolites, or to slow metabolic degradation. Tolerance to elevated blood levels varies with the status of the patient. Debilitated, elderly patients and acutely ill patients should be given reduced doses commensurate with their age and physical status. Reduced doses may be indicated in patients with increased intra-abdominal pressure (including obstetrical patients), if otherwise suitable for spinal anesthesia.

5.4 Risk of Systemic Toxicities with Unintended Intravascular Injection

Unintended intravascular injection of Bupivacaine Hydrochloride in Dextrose Injection,

USP may be associated with systemic toxicities, including CNS or cardiorespiratory depression and coma, progressing ultimately to respiratory arrest [*see Adverse Reactions (6)*].

Aspirate for blood and cerebrospinal fluid before injecting Bupivacaine Hydrochloride in Dextrose Injection, USP, for both the initial dose and all subsequent doses (where applicable), to confirm entry into the subarachnoid space and to avoid intravascular injection. Aspiration of cerebrospinal fluid into a Bupivacaine Hydrochloride in Dextrose Injection, USP-filled syringe will result in an identifiable swirl in the solution. A negative aspiration for blood does not ensure against an intravascular injection.

5.5 Methemoglobinemia

Cases of methemoglobinemia have been reported in association with local anesthetic use. Although all patients are at risk for methemoglobinemia, patients with glucose 6 phosphate dehydrogenase deficiency, congenital or idiopathic methemoglobinemia, cardiac or pulmonary compromise, infants under 6 months of age, and concurrent exposure to oxidizing agents or their metabolites are more susceptible to developing clinical manifestations of the condition [*see Drug Interactions (7.2)*]. If local anesthetics must be used in these patients, close monitoring for symptoms and signs of methemoglobinemia is recommended.

Signs of methemoglobinemia may occur immediately or may be delayed some hours after exposure, and are characterized by a cyanotic skin discoloration and/or abnormal coloration of the blood. Methemoglobin levels may continue to rise; therefore, immediate treatment is required to avert more serious CNS and cardiovascular adverse effects, including seizures, coma, arrhythmias, and death. Discontinue Bupivacaine Hydrochloride in Dextrose Injection, USP and any other oxidizing agents. Depending on the severity of the signs and symptoms, patients may respond to supportive care, i.e., oxygen therapy, hydration. A more severe clinical presentation may require treatment with methylene blue, exchange transfusion, or hyperbaric oxygen.

5.6 Risk of Cardiac Arrest with Use of Epidural Bupivacaine in Obstetrical Anesthesia

There have been reports of cardiac arrest with difficult resuscitation or death during use of Bupivacaine Hydrochloride Injection, USP for epidural anesthesia in obstetrical patients. In most cases, this has followed use of Bupivacaine Hydrochloride Injection, USP 0.75%, not Bupivacaine Hydrochloride in Dextrose Injection, USP. The package insert for Bupivacaine Hydrochloride Injection, USP for epidural, nerve block, etc., has a more complete discussion of preparation for, and management of cardiac arrest following epidural administration. Bupivacaine Hydrochloride in Dextrose Injection, USP is recommended for spinal anesthesia in obstetrical patients.

5.7 Chondrolysis with Intra-Articular Infusion

Intra-articular infusions of local anesthetics including bupivacaine following arthroscopic and other surgical procedures is an unapproved use, and there have been post-marketing reports of chondrolysis in patients receiving such infusions. The majority of reported cases of chondrolysis have involved the shoulder joint; cases of gleno-humeral chondrolysis have been described in pediatric and adult patients following intra-articular infusions of local anesthetics with and without epinephrine for periods of 48 to 72

hours. There is insufficient information to determine whether shorter infusion periods are not associated with chondrolysis. The time of onset of symptoms, such as joint pain, stiffness, and loss of motion can be variable, but may begin as early as the 2nd month after surgery. Currently, there is no effective treatment for chondrolysis; patients who experienced chondrolysis have required additional diagnostic and therapeutic procedures and some required arthroplasty or shoulder replacement.

5.8 Risk of Cardiac Arrest with Intravenous Regional Anesthesia Use (Bier Block)

There have been reports of cardiac arrest and death during the use of bupivacaine for intravenous regional anesthesia (Bier Block). Information on safe dosages and techniques of administration of Bupivacaine Hydrochloride in Dextrose Injection, USP in this procedure is lacking. Therefore, Bupivacaine Hydrochloride in Dextrose Injection, USP is contraindicated for use with this technique [*see Contraindications (4)*].

5.9 Patients with Severe Disturbances of Cardiac Rhythm, Shock, or Heart Block

Consider alternate anesthetic techniques in patients with severe disturbances of cardiac rhythm, shock, or heart block [*see Contraindications (4)*].

5.10 Risk of Toxicity in Patients with Hepatic Impairment

Because amide-type local anesthetics such as bupivacaine are metabolized by the liver, consider reduced dosing and increased monitoring for bupivacaine systemic toxicity in patients with moderate to severe hepatic impairment who are treated with Bupivacaine Hydrochloride in Dextrose Injection, USP [*see Use in Specific Populations (8.6)*]. Most information regarding dose-related hepatic impairment is based on larger dosages of bupivacaine administered for other neuraxial, peripheral nerve, or fascial plane blocks.

5.11 Risk of Use in Patients with Impaired Cardiovascular Function

Bupivacaine Hydrochloride in Dextrose Injection, USP should be given in reduced doses in patients with impaired cardiovascular function (e.g., hypotension, heartblock, valvular abnormalities) because they may be less able to compensate for functional changes associated with the sympathetic blockade observed after subarachnoid administration of Bupivacaine Hydrochloride in Dextrose Injection, USP and the prolongation of AV conduction produced by the drug. Monitor patients closely for blood pressure, heart rate, and ECG changes.

6 ADVERSE REACTIONS

The following clinically significant adverse reactions have been reported and described in other sections of the labeling:

- Allergic-Type Reactions [*see Contraindications (4)*]
- Dose-Related Toxicity [*see Warnings and Precautions (5.3)*]
- Systemic Toxicities with Unintended Intravascular Injection [*see Warnings and Precautions (5.4)*]
- Methemoglobinemia [*see Warnings and Precautions (5.5)*]
- Cardiac Arrest in Obstetrical Anesthesia [*see Warnings and Precautions (5.6)*]

- Chondrolysis with Intra-Articular Infusion [see *Warnings and Precautions (5.7)*]
- Cardiac Arrest with Intravenous Regional Anesthesia Use [see *Contraindications (4), Warnings and Precautions (5.8)*]

The following adverse reactions from voluntary reports or clinical studies have been reported with bupivacaine. Because many of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Adverse reactions to Bupivacaine Hydrochloride in Dextrose Injection, USP are characteristic of those associated with other amide-type local anesthetics. A major cause of adverse reactions to Bupivacaine Hydrochloride in Dextrose Injection, USP is due to cephalad extension of the motor level of anesthesia and/or excessive plasma levels, which may be due to overdosage, unintentional intravascular injection, or slow metabolic degradation.

The most commonly encountered acute adverse reactions that demand immediate counter-measures following the administration of spinal anesthesia were hypotension due to loss of sympathetic tone and respiratory paralysis or underventilation due to cephalad extension of the motor level of anesthesia. These have led to cardiac arrest if untreated. In addition, dose-related convulsions and cardiovascular collapse have resulted from diminished tolerance, rapid absorption from the injection site, or from unintentional intravascular injection of a local anesthetic solution.

Respiratory System: Respiratory paralysis or underventilation have been noted as a result of cephalad spread of spinal anesthesia and has led to secondary hypoxic cardiac arrest when untreated. Preanesthetic medication, intraoperative anesthetics, analgesics, and sedatives, as well as surgical manipulation, may contribute to underventilation. This has usually been noted within minutes of the injection of spinal anesthetic solution, but because of differing maximal onset times, differing intercurrent drug usage, and differing surgical manipulation, it may occur at any time during surgery or the immediate recovery period.

Cardiac Disorders: Hypotension due to loss of sympathetic tone has been commonly encountered following spinal anesthesia. This has been more commonly observed in elderly patients, particularly those with hypertension, and patients with reduced blood volume, reduced interstitial fluid volume, cephalad spread of the local anesthetic, and/or mechanical obstruction of venous return. Nausea and vomiting have been frequently associated with hypotensive episodes following the administration of spinal anesthesia. High doses, or inadvertent intravascular injection, have led to high plasma levels and related depression of the myocardium, decreased cardiac output, bradycardia, heart block, ventricular arrhythmias, and cardiac arrest [see *Warnings and Precautions (5.4)*].

Nervous System Disorders: Respiratory paralysis or underventilation secondary to cephalad spread of the level of spinal anesthesia (see *Respiratory System*) and hypotension for the same reason (see *Cardiac Disorders*) have been the two most commonly encountered CNS-related adverse observations which demand immediate counter-measures.

High doses or inadvertent intravascular injection have led to high plasma levels and related CNS toxicity. Adverse reactions were characterized by excitation and/or depression of the CNS and included restlessness, anxiety, dizziness, tinnitus, blurred vision, tremors, convulsions, drowsiness, unconsciousness, and respiratory arrest.

The incidences of adverse neurologic reactions associated with the use of local anesthetics may be related to the total dose of local anesthetic administered and are also dependent upon the particular drug used, the route of administration, and the physical status of the patient.

Convulsions: Incidence varied with the procedure used and the total dose administered. In a survey of studies of epidural anesthesia, overt toxicity progressing to convulsions occurred in approximately 0.1% of local anesthetic administrations. The incidences of adverse neurologic reactions associated with the use of local anesthetics may be related to the total dose of local anesthetic administered and are also dependent upon the particular drug used, the route of administration, and the physical status of the patient.

Neurologic effects following spinal anesthesia have included loss of perineal sensation and sexual function; persistent anesthesia, paresthesia, weakness and paralysis of the lower extremities, and loss of sphincter control with slow, incomplete, or no recovery; hypotension, high or total spinal block; urinary retention; headache; backache; septic meningitis, meningismus; arachnoiditis; slowing of labor; increased incidence of forceps delivery; shivering; cranial nerve palsies due to traction on nerves from loss of cerebrospinal fluid; and fecal and urinary incontinence.

Immune System Disorders: Allergic-type reactions have occurred as a result of sensitivity to bupivacaine. These reactions were characterized by signs such as urticaria, pruritus, erythema, angioneurotic edema (including laryngeal edema), tachycardia, sneezing, nausea, vomiting, dizziness, syncope, excessive sweating, elevated temperature, and severe hypotension. Cross sensitivity among members of the amide-type local anesthetic group has been reported.

7 DRUG INTERACTIONS

7.1 Local Anesthetics

The toxic effects of local anesthetics are additive. If coadministration of other local anesthetics with Bupivacaine Hydrochloride in Dextrose Injection, USP cannot be avoided, monitor patients for neurologic and cardiovascular effects related to local anesthetic systemic toxicity [*see Dosage and Administration (2.1), Warnings and Precautions (5.3)*].

7.2 Drugs Associated with Methemoglobinemia

Patients who are administered Bupivacaine Hydrochloride in Dextrose Injection, USP are at increased risk of developing methemoglobinemia when concurrently exposed to the following drugs, which could include other local anesthetics [*see Warnings and Precautions (5.5)*].

Examples of Drugs Associated with Methemoglobinemia:

Class	Examples
Nitrates/Nitrites	nitric oxide, nitroglycerin, nitroprusside, nitrous oxide
Local anesthetics	articaine, benzocaine, bupivacaine, lidocaine, mepivacaine, prilocaine, procaine, ropivacaine, tetracaine

Antineoplastic agents	cyclophosphamide, flutamide, hydroxyurea, isofamide, rasburicase
Antibiotics	dapsone, nitrofurantoin, para-aminosalicylic acid, sulfonamides
Antimalarials	chloroquine, primaquine
Anticonvulsants	phenobarbital, phenytoin, sodium valproate
Other drugs	acetaminophen, metoclopramide, quinine, sulfasalazine

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

The available data on the use of Bupivacaine Hydrochloride in Dextrose Injection, USP in pregnant women do not establish the presence or absence of developmental toxicity related to the use of Bupivacaine Hydrochloride in Dextrose Injection, USP.

In animal studies, embryo-fetal lethality was noted when bupivacaine was administered subcutaneously to pregnant rabbits during organogenesis and decreased pup survival was observed in a rat pre-and post-natal developmental study (dosing from implantation through weaning). These effects were observed at dose levels approximately 30 times the daily maximum recommended human dose (MRHD) on a body surface area (BSA) basis. Based on animal data, advise pregnant women of the potential risk to a fetus (see *Data*).

Local anesthetics rapidly cross the placenta, and when used for epidural, caudal, or pudendal block anesthesia, can cause varying degrees of maternal, fetal, and neonatal toxicity [see *Clinical Pharmacology (12.3)*]. The incidence and degree of toxicity depend upon the procedure performed, the type, and amount of drug used, and the technique of drug administration. Adverse reactions in the parturient, fetus, and neonate involve alterations of the CNS, peripheral vascular tone, and cardiac function.

If this drug is used during pregnancy or if the patient becomes pregnant while taking this drug, inform the patient of the potential hazard to the fetus. The estimated background risk of major birth defects and miscarriage for the indicated populations are unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

Clinical Considerations

Maternal Adverse Reactions

Maternal hypotension has resulted from regional and neuraxial anesthesia. Local anesthetics produce vasodilation by blocking sympathetic nerves. The supine position is dangerous in pregnant women at term because of aortocaval compression by the gravid uterus. Therefore, during treatment of systemic toxicity, maternal hypotension, or fetal bradycardia following regional or neuraxial block, the parturient should be maintained in the left lateral decubitus position if possible, or manual displacement of the uterus off the great vessels be accomplished. Elevating the patient's legs and right-side-up positioning will help prevent decreases in blood pressure. The fetal heart rate also should be monitored continuously and electronic fetal monitoring is highly advisable.

Labor or Delivery

Spinal anesthesia is commonly used during labor and delivery. Bupivacaine hydrochloride, when administered properly, via the epidural route in doses 10 to 12 times the amount used in spinal anesthesia has been used for obstetrical analgesia and anesthesia without evidence of adverse effects on the fetus.

Spinal anesthesia may alter the forces of parturition through changes in uterine contractility or maternal expulsive efforts. Spinal anesthesia has also been reported to prolong the second stage of labor by removing the parturient's reflex urge to bear down or by interfering with motor function. The use of obstetrical anesthesia may increase the need for forceps assistance.

The use of some local anesthetic drug products during labor and delivery may be followed by diminished muscle strength and tone for the first day or two of life. This has not been reported with bupivacaine.

It is extremely important to avoid aortocaval compression by the gravid uterus during administrations of regional or neuraxial block to parturients. To do this, the patient must be maintained in the left lateral decubitus position or a blanket roll or sandbag may be placed beneath the right hip and the gravid uterus displaced to the left.

Data

Animal Data

Bupivacaine hydrochloride produced developmental toxicity when administered subcutaneously to pregnant rats and rabbits at doses 30-times the MHRD.

Bupivacaine hydrochloride was administered subcutaneously to rats at doses of 4.4, 13.3, and 40 mg/kg and to rabbits at doses of 1.3, 5.8, and 22.2 mg/kg during the period of organogenesis (implantation to closure of the hard palate). The high doses are approximately 30-times the daily MRHD of 12 mg/day on a mg dose/m² BSA basis. No embryo-fetal effects were observed in rats at the high dose which caused increased maternal lethality. An increase in embryo-fetal deaths was observed in rabbits at the high dose in the absence of maternal toxicity with the fetal No Observed Adverse Effect Level being approximately 8-times the MRHD on a BSA basis.

In a rat pre-and post-natal development study (dosing from implantation through weaning) conducted at subcutaneous doses of 4.4, 13.3, and 40 mg/kg, decreased pup survival was observed at the high dose. The high dose is approximately 30-times the daily MRHD of 12 mg/day on a BSA basis.

8.2 Lactation

Risk Summary

Lactation studies have not been conducted with bupivacaine. Bupivacaine has been reported to be excreted in human milk suggesting that the nursing infant could be theoretically exposed to a dose of the drug. Bupivacaine Hydrochloride in Dextrose Injection, USP should be administered to lactating women only if clearly indicated. Studies assessing the effects of Bupivacaine Hydrochloride in Dextrose Injection, USP in breastfed children have not been performed. Studies to assess the effect of Bupivacaine Hydrochloride in Dextrose Injection, USP on milk production or excretion have not been performed. The developmental and health benefits of breastfeeding should be

considered along with the mother's clinical need for bupivacaine and any potential adverse effects on the breastfed child from bupivacaine or from the underlying maternal condition.

8.4 Pediatric Use

Bupivacaine Hydrochloride in Dextrose Injection, USP is approved for use in adults only. Administration of Bupivacaine Hydrochloride in Dextrose Injection, USP in patients younger than 18 is not recommended.

8.5 Geriatric Use

Patients 65 years and over, particularly those with hypertension, may be at increased risk for developing hypotension while undergoing spinal anesthesia with Bupivacaine Hydrochloride in Dextrose Injection, USP.

In clinical studies of bupivacaine, elderly patients exhibited a greater spread and higher maximal level of anesthesia than younger patients. Elderly patients also reached the maximal level of anesthesia more rapidly than younger patients, and exhibited a faster onset of motor blockade.

Differences in various pharmacokinetic parameters have been observed between elderly and younger patients [*see Clinical Pharmacology (12.3)*].

This product is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function. Elderly patients may require lower doses of Bupivacaine Hydrochloride in Dextrose Injection, USP.

8.6 Hepatic Impairment

Amide-type local anesthetics, such as bupivacaine, are metabolized by the liver. Patients with severe hepatic impairment, because of their inability to metabolize local anesthetics normally, are at a greater risk of developing toxic plasma concentrations, and potentially local anesthetic systemic toxicity. Therefore, consider reduced dosing and increased monitoring for local anesthetic systemic toxicity in patients with moderate to severe hepatic impairment treated with Bupivacaine Hydrochloride in Dextrose Injection, USP [*see Warnings and Precautions (5.10)*].

8.7 Renal Impairment

Bupivacaine is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with renal impairment. This should be considered when selecting the Bupivacaine Hydrochloride in Dextrose Injection, USP dosage [*see Use in Specific Populations (8.5)*].

10 OVERDOSAGE

Clinical Presentation

Acute emergencies from Bupivacaine Hydrochloride in Dextrose Injection, USP are generally related to hypoventilation (and perhaps apnea) secondary to upward extension

of spinal anesthesia or high plasma levels encountered during therapeutic use [see *Warnings and Precautions (5.3)*, *Adverse Reactions (6)*]. Hypotension is commonly encountered during the conduct of spinal anesthesia due to loss of sympathetic tone, and sometimes, contributory mechanical obstruction of venous return due to the gravid uterus exerting pressure on the great vessels [see *Warnings and Precautions (5.2)*, *Adverse Reactions (6)*].

If not treated immediately, convulsions with simultaneous hypoxia, hypercarbia, and acidosis plus myocardial depression from the direct effects of the local anesthetic may result in cardiac arrhythmias, bradycardia, asystole, ventricular fibrillation, or cardiac arrest. Respiratory abnormalities, including apnea, may occur. Hypoventilation or apnea due to a high or total spinal may produce these same signs and also lead to cardiac arrest if ventilatory support is not instituted. If cardiac arrest should occur, successful outcome may require prolonged resuscitative efforts.

Management

The first step in the management of systemic toxic reactions, as well as hypoventilation or apnea due to a high or total spinal, consists of immediate attention to the establishment and maintenance of a patent airway and effective assisted or controlled ventilation with 100% oxygen with a delivery system capable of permitting immediate positive airway pressure by mask. Endotracheal intubation, using drugs and techniques familiar to the clinician, may be indicated after initial administration of oxygen by mask if difficulty is encountered in the maintenance of a patent airway, or if prolonged ventilatory support (assisted or controlled) is indicated.


If necessary, use drugs to manage the convulsions. A bolus intravenous dose of benzodiazepine will counteract the CNS stimulation related to Bupivacaine Hydrochloride in Dextrose Injection, USP. Immediately after the institution of ventilatory measures, evaluate the adequacy of the circulation. Supportive treatment of circulatory depression may require Advanced Cardiac Life Support measures.

Hypotension due to sympathetic relaxation may be managed by giving intravenous fluids (such as isotonic saline or lactated Ringer's solution), in an attempt to relieve mechanical obstruction of venous return, or by using vasopressor agents (such as ephedrine which increases myocardial contractility) and, if indicated, by giving plasma expanders or blood products.


11 DESCRIPTION

Bupivacaine Hydrochloride in Dextrose Injection, USP is an amide-local anesthetic and sterile hyperbaric aqueous solution. The route of administration for Bupivacaine Hydrochloride in Dextrose Injection, USP is by subarachnoid injection. Bupivacaine Hydrochloride in Dextrose Injection, USP contains bupivacaine hydrochloride, as the active pharmaceutical ingredient and also contains Dextrose, as baricity agent.

Bupivacaine Hydrochloride (monohydrate) chemical name is 2-piperidinecarboxamide, 1-butyl-N-(2,6dimethylphenyl)-, monohydrochloride, monohydrate, a white crystalline powder that is freely soluble in 95 percent ethanol, soluble in water, and slightly soluble in chloroform or acetone. Bupivacaine Hydrochloride (monohydrate) has a molecular formula of $C_{18}H_{28}N_2O \cdot HCl \cdot H_2O$ and molecular weight of 342.90 g/mol and has the following structural formula:



Dextrose chemical name is D-glucopyranose. Dextrose (anhydrous) has a molecular formula of C₆H₁₂O₆, molecular weight of 180.16 g/mol and has the following structural formula:



Bupivacaine Hydrochloride in Dextrose Injection, USP is a clear and colorless sterile hyperbaric solution.

Each mL of Bupivacaine Hydrochloride in Dextrose Injection, USP contains 7.5 mg bupivacaine hydrochloride (anhydrous) (equivalent to 7.9 mg of bupivacaine hydrochloride monohydrate), 82.5 mg dextrose (anhydrous) as baricity agent, and sodium hydroxide and hydrochloric acid as pH adjusters in water for injection.

Bupivacaine Hydrochloride in Dextrose Injection, USP pH is between 4.0 and 6.5.

The specific gravity of Bupivacaine Hydrochloride in Dextrose Injection, USP is between 1.030 and 1.035 at 25°C and 1.03 at 37°C.

Bupivacaine Hydrochloride in Dextrose Injection, USP does not contain any preservatives.

12 CLINICAL PHARMACOLOGY

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Bupivacaine blocks the generation and the conduction of nerve impulses, presumably by increasing the threshold for electrical excitation in the nerve, by slowing the propagation of the nerve impulse, and by reducing the rate of rise of the action potential. In general, the progression of anesthesia is related to the diameter, myelination, and conduction velocity of affected nerve fibers. Clinically, the order of loss of nerve function is as follows: (1) pain, (2) temperature, (3) touch, (4) proprioception, and (5) skeletal muscle tone.

12.2 Pharmacodynamics

Systemic absorption of bupivacaine produces effects on the cardiovascular system and CNS. At blood concentrations achieved with normal therapeutic doses, changes in cardiac conduction, excitability, refractoriness, contractility, and peripheral vascular resistance are minimal. However, toxic blood concentrations depress cardiac conduction and excitability, which may lead to atrioventricular block, ventricular arrhythmias, and cardiac arrest, sometimes resulting in fatalities. In addition, myocardial contractility is depressed and peripheral vasodilation occurs, leading to decreased cardiac output and arterial blood pressure. These cardiovascular changes are more likely to occur after unintended intravascular injection of bupivacaine [see *Warnings and Precautions* (5.4)].

Following systemic absorption, bupivacaine can produce CNS stimulation, CNS depression, or both. Apparent central stimulation is manifested as restlessness, tremors and shivering, progressing to convulsions, followed by CNS depression and coma progressing ultimately to respiratory arrest. However, bupivacaine has a primary depressant effect on the medulla and on higher centers. The depressed stage may occur without a prior excited stage.

The duration of local anesthesia after administration of Bupivacaine Hydrochloride in Dextrose Injection, USP is longer than that observed after administration of other commonly used short-acting local anesthetic. There appears to be a period of analgesia that persists after the resolution of the block and the return of sensation.

The onset of sensory blockade following spinal block with Bupivacaine Hydrochloride in Dextrose Injection, USP is rapid (generally within one minute); maximum motor blockade and maximum dermatome level are achieved within 15 minutes in most cases. Duration of sensory blockade (time to return of complete sensation in the operative site or regression of two dermatomes) following Bupivacaine Hydrochloride in Dextrose Injection, USP 12 mg averages 2 hours with or without 0.2 mg epinephrine. The time to return of complete motor ability with Bupivacaine Hydrochloride in Dextrose Injection, USP 12 mg averages 3.5 hours without the addition of epinephrine and 4.5 hours if 0.2 mg epinephrine is added. When compared to equal milligram doses of hyperbaric tetracaine, the duration of sensory blockade was the same, but the time to complete motor recovery was longer for tetracaine. Addition of 0.2 mg epinephrine prolongs the motor blockade and time to first postoperative opioid with Bupivacaine Hydrochloride in Dextrose Injection, USP.

12.3 Pharmacokinetics

Systemic plasma levels of bupivacaine following administration of Bupivacaine Hydrochloride in Dextrose Injection, USP do not correlate with local efficacy.

Absorption

The rate of systemic absorption of bupivacaine is dependent upon the total dose and concentration of drug administered, the route of administration, the vascularity of the administration site, and the presence or absence of epinephrine in the anesthetic solution. A dilute concentration of epinephrine (1:200,000) usually reduces the rate of absorption and peak plasma concentration of bupivacaine, permitting the use of moderately larger total doses and sometimes prolonging the duration of action.

Distribution

Bupivacaine appears to cross the placenta by passive diffusion. The rate and degree of

diffusion is governed by

(1) the degree of plasma protein binding, (2) the degree of ionization, and (3) the degree of lipid solubility. Fetal/maternal ratios of bupivacaine appear to be inversely related to the degree of plasma protein binding, because only the free, unbound drug is available for placental transfer. Bupivacaine with a high protein binding capacity (95%) has a low fetal/maternal ratio (0.2 to 0.4). The extent of placental transfer is also determined by the degree of ionization and lipid solubility of the drug. Lipid soluble, nonionized drugs readily enter the fetal blood from the maternal circulation.

Depending upon the route of administration, bupivacaine is distributed to some extent to all body tissues, with high concentrations found in highly perfused organs such as the liver, lungs, heart, and brain.

Pharmacokinetic studies on the plasma profiles of bupivacaine after direct intravenous injection (Bupivacaine Hydrochloride in Dextrose Injection, USP is not approved for intravenous use, and contraindicated for intravenous regional block, i.e. Bier Block) suggest a three-compartment open model. The first compartment is represented by the rapid intravascular distribution of the drug. The second compartment represents the equilibration of the drug throughout the highly perfused organs such as the brain, myocardium, lungs, kidneys, and liver. The third compartment represents an equilibration of the drug with poorly perfused tissues, such as muscle and fat.

Elimination

The half-life of bupivacaine in adults is 2.7 hours.

Metabolism

Amide-type local anesthetics such as bupivacaine are metabolized primarily in the liver via conjugation with glucuronic acid. Pipecoloxylidine is the major metabolite of bupivacaine. The elimination of drug from tissue distribution depends largely upon the ability of binding sites in the circulation to carry it to the liver where it is metabolized.

Excretion

The kidney is the main excretory organ for most local anesthetics and their metabolites. Urinary excretion is affected by urinary perfusion and factors affecting urinary pH. Only 6% of bupivacaine is excreted unchanged in the urine.

Specific Populations

Geriatric Patients

The total plasma clearance was decreased and the terminal half-life was lengthened in these patients.

Patients with Hepatic Impairment

Various pharmacokinetic parameters of the local anesthetics can be significantly altered by the presence of hepatic disease. Patients with hepatic disease, especially those with severe hepatic disease, may be more susceptible to the potential toxicities of the amide-type local anesthetics [see *Use in Specific Populations* (8.6)].

Patients with Renal Impairment

Various pharmacokinetic parameters of the local anesthetics can be significantly altered by the presence of renal disease, factors affecting urinary pH, and renal blood flow [see

Use in Specific Populations (8.5, 8.7)].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term studies in animals to evaluate the carcinogenic potential of bupivacaine hydrochloride have not been conducted.

Mutagenesis

The mutagenic potential of bupivacaine hydrochloride has not been determined.

Impairment of Fertility

The effect of bupivacaine on fertility has not been determined.

16 HOW SUPPLIED/STORAGE AND HANDLING

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F). [See USP Controlled Room Temperature.]

Bupivacaine Hydrochloride in Dextrose Injection, USP solution may be autoclaved once at 15 pound pressure, 121°C (250°F) for 15 minutes. This product is clear and colorless. Do not use the solution if it is discolored or contains particulate matter.

Single-dose ampules of 2 mL Bupivacaine Hydrochloride in Dextrose Injection, USP (15 mg bupivacaine hydrochloride with 165 mg dextrose) are supplied as follows:

NDC	Container	Concentration	Size	Total (mg)
71351-022-10	Single Dose Ampule (Package of 10)	0.75% (7.5 mg/mL)	2mL	15
71351-022-50	Single Dose Ampule (Package of 50)	0.75% (7.5 mg/mL)	2mL	15

Discard the unused portion.

17 PATIENT COUNSELING INFORMATION

Allergic-Type Reactions

Assess if the patient has had allergic-type reactions to amide-type local anesthetics or to other formulation ingredients [see *Contraindications (4)*, *Adverse Reactions (6)*].

Temporary Loss of Sensation and Motor Activity After Spinal Anesthesia

When appropriate, patients should be informed in advance that they may experience temporary loss of sensation and motor activity, usually in the lower half of the body, including the inability to move the legs, following proper administration of spinal anesthesia.

Methemoglobinemia

Inform patients that use of local anesthetics may cause methemoglobinemia, a serious condition that must be treated promptly. Advise patients or caregivers to seek immediate medical attention if they or someone in their care experience the following signs or symptoms: pale, gray, or blue colored skin (cyanosis), headache, rapid heart rate, shortness of breath, lightheadedness, or fatigue [see Warnings and Precautions (5.5)].

This product's labeling may have been updated.

Distributed by:

Brookfield Pharmaceuticals, LLC

Brookfield, WI 53005

Principal Display Panel

2 mL (10 Ampules Unit Dose) Carton Label

10 x 2 mL Single-dose Ampules

NDC 71351-022-10

STERILE HYPERBARIC SOLUTION FOR SPINAL ANESTHESIA


Bupivacaine Hydrochloride in 8.25% Dextrose Injection, USP SPINAL 0.75%

(15 mg/2 mL)

(7.5 mg/mL)

Distributed by:

Brookfield Pharmaceuticals, LLC



2 mL (50 Ampules Unit Dose) Carton Label

50 x 2 mL Single-dose Ampules

NDC 71351-022-50


STERILE HYPERBARIC SOLUTION FOR SPINAL ANESTHESIA

Bupivacaine Hydrochloride in 8.25%
Dextrose Injection, USP
SPINAL 0.75%

(15 mg/2 mL)
(7.5 mg/mL)

Distributed by:

Brookfield Pharmaceuticals, LLC



2 mL Ampule Label

2 mL Ampule NDC 71351-022-02

EL6461-02

Rx only

**Bupivacaine HCl 0.75%
in Dextrose 8.25%**
Injection, USP
SPINAL


Contains 7.5 mg bupivacaine HCl (anhydrous) and 82.5 mg dextrose (anhydrous) per mL. pH adjusted between 4.0 and 6.5 with NaOH or HCl.

Distributed by:

Brookfield Pharmaceuticals, LLC

LOT NO. : XXXXXX
EXP. DATE : YYYY/MM/DD

33



2 mL Ampule

Bupivacaine HCl
in Dextrose
Injection, USP
SPINAL

Contains 7.5 mg bupivacaine HCl (anhydrous) and 82.5 mg dextrose (anhydrous) per mL. pH adjusted between 4.0 and 6.5 with NaOH or HCl.
Distributed by:
Brookfield Pharmaceuticals, LLC
Brookfield, WI 53005

NDC 71351-022-02

Rx only

LOT NO. : XXXXXX
EXP. DATE : YYYY/MM/DD

EL6461-02

22

BUPIVACAINE HYDROCHLORIDE			
bupivacaine hydrochloride injection			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71351-022
Route of Administration	SUBARACHNOID		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)		BUPIVACAINE HYDROCHLORIDE ANHYDROUS	7.5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
DEXTROSE (UNII: IY9XDZ35W2)	82.5 mg in 1 mL
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71351-022-50	50 in 1 PACKAGE	10/31/2022	
1	NDC:71351-022-02	2 mL in 1 AMPULE; Type 0: Not a Combination Product		
2	NDC:71351-022-10	10 in 1 PACKAGE	02/23/2023	
2	NDC:71351-022-02	2 mL in 1 AMPULE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA212822	10/31/2022	

Labeler - Brookfield Pharmaceuticals, LLC. (080592685)

Revised: 1/2026

Brookfield Pharmaceuticals, LLC.