

**LAXATIVE PILLS MAXIMUM STRENGTH- sennosides tablet
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

Drug Facts

Active ingredient (in each tablet)

Sennosides USP, 25 mg

Purpose

Stimulant laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Ask a doctor before use if you have

- noticed a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist before use if you

- are taking a prescription drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.

When using this product

- do not use for a period longer than 1 week

Stop use and ask a doctor if

- rectal bleeding or failure to have a bowel movement occur after use of a laxative

These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- swallow tablet(s) with a glass of water
- swallow tablet(s) whole, do not crush, break, or chew

| | |
|---|------------------------------------|
| adults and children 12 years of age and older | take 2 tablets once or twice daily |
| children 6 to under 12 years of age | take 1 tablet once or twice daily |
| children under 6 years of age | ask a doctor |

Other information

- each tablet contains: **calcium 40 mg**
- each tablet contains: **magnesium 10 mg**
- each tablet contains: **sodium 3 mg**
- very low sodium
- store between 20-25°C (68-77°F)

Inactive ingredients

acacia, ammonium hydroxide*, anhydrous lactose*, calcium phosphate dihydrate*, calcium sulfate anhydrous*, carnauba wax, corn starch*, croscarmellose sodium, dicalcium phosphate*, FD&C blue #1 aluminum lake, hypromellose*, iron oxide*, iron oxide black, iron oxide yellow*, isopropyl alcohol*, kaolin*, magnesium stearate, microcrystalline cellulose, n-butyl alcohol*, polyethylene glycol, povidone*, propylene glycol*, shellac, silicon dioxide*, sodium lauryl sulfate*, stearic acid*, sucrose, talc, tartaric acid*, titanium dioxide, white wax*

*contains one or more of these ingredients

Questions or comments?

Call toll free **1-866-4-REXALL**

Principal Display Panel

MAXIMUM STRENGTH

Laxative Pills

Sennosides 25 mg

Stimulant Laxative

- For gentle, dependable overnight relief

Visit us at: Rexall.com or call 1-866-4-REXALL

PACKAGED FOR DOLGENCORP, LLC

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072 USA

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT
INFORMATION**

Product Label



MAXIMUM STRENGTH

Laxative Pills

Senosides 25 mg
Stimulant Laxative

- For gentle, dependable overnight relief



48
Pills

Visit us at: Rexall.com or call 1-866-4-REXALL

PACKAGED FOR DOLGENCORP, LLC
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072 USA

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Lot No.:
Exp. Date:

Drug Facts (continued)

Directions
 ■ swallow tablet(s) with a glass of water
 ■ swallow tablet(s) whole, do not crush, break, or chew
 adults and children 12 years of age and older take 2 tablets once or twice daily
 children 6 to under 12 years of age take 1 tablet once or twice daily
 children under 6 years of age ask a doctor

Other information
 ■ each tablet contains: calcium 40 mg
 ■ each tablet contains: magnesium 10 mg
 ■ each tablet contains: sodium 3 mg
 ■ very low sodium
 ■ store between 20-25°C (68-77°F)

Drug Facts (continued)

Drug Facts (continued)
 ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
 ■ ask a doctor or pharmacist before use if you are taking a prescription drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.
 ■ do not use for a period longer than 1 week
 ■ stop use and ask a doctor if rectal bleeding or failure to have a bowel movement occur after use of a laxative
 ■ these may be signs of a serious condition.

Drug Facts (continued)
 ■ pregnant or breast-feeding, ask a health professional before use.
 ■ keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ swallow tablet(s) with a glass of water
 ■ swallow tablet(s) whole, do not crush, break, or chew
 adults and children 12 years of age and older take 2 tablets once or twice daily
 children 6 to under 12 years of age take 1 tablet once or twice daily
 children under 6 years of age ask a doctor

Questions or comments?
 Call toll free 1-866-4-REXALL

Inactive ingredients (continued)
 acacia, ammonium hydroxide, anhydrous lactose, calcium phosphate dihydrate, calcium sulfate anhydrous, carnauba wax, corn starch, croscarmellose sodium, dicalcium phosphate, FD&C blue #1 aluminum lake, hypromellose, iron oxide black, iron oxide yellow, isopropyl alcohol, kaolin, magnesium stearate, microcrystalline cellulose, n-butyl alcohol, polyethylene glycol, povidone, propylene glycol, shellac, silicon dioxide, sodium lauryl sulfate, stearic acid, sucrose, talc, tartaric acid, titanium dioxide, white wax

Drug Facts (continued)

Active ingredient
 (in each tablet)
 Senosides USP, 25 mg.....Stimulant laxative

Purpose
 Stimulant laxative

Uses
 ■ relieves occasional constipation (irregularity)
 ■ generally produces bowel movement in 6 to 12 hours

Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.
 Ask a doctor before use if you have

Drug Facts (continued)

Drug Facts (continued)
 ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks
 ■ ask a doctor or pharmacist before use if you are taking a prescription drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.
 ■ do not use for a period longer than 1 week
 ■ stop use and ask a doctor if rectal bleeding or failure to have a bowel movement occur after use of a laxative
 ■ these may be signs of a serious condition.

PLD-B
FC001666



Rexall Laxative Pills

LAXATIVE PILLS MAXIMUM STRENGTH

sennosides tablet

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:55910-245 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| SENNOSIDES (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX) | SENNOSIDES | 25 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| ACACIA (UNII: 5C5403N26O) | |
| AMMONIA (UNII: 5138Q19F1X) | |
| ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK) | |
| CALCIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: O7TSZ97GEP) | |
| CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| CROSCARMELLOSE SODIUM (UNII: M28OL1HH48) | |
| CALCIUM PHOSPHATE, DIBASIC, ANHYDROUS (UNII: L11K75P92J) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| ALUMINUM OXIDE (UNII: LMI26O6933) | |
| HYPROMELLOSES (UNII: 3NXW29V3WO) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| FERROSFERRIC OXIDE (UNII: XM0M87F357) | |
| FERRIC OXIDE YELLOW (UNII: EX438O2MRT) | |
| ISOPROPYL ALCOHOL (UNII: ND2M416302) | |
| KAOLIN (UNII: 24H4NWX5CO) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) | |
| BUTYL ALCOHOL (UNII: 8PJ61P6TS3) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POVIDONE (UNII: FZ989GH94E) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |

| | |
|---|--|
| SHELLAC (UNII: 46N107B71O) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| SODIUM LAURYL SULFATE (UNII: 368GB5141J) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| SUCROSE (UNII: C151H8M554) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TARTARIC ACID (UNII: W4888I119H) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| WHITE WAX (UNII: 7G1J5DA97F) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|------------|
| Color | blue | Score | no score |
| Shape | ROUND | Size | 11mm |
| Flavor | | Imprint Code | TCL083;S25 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:55910-245-48 | 4 in 1 CARTON | 12/30/2013 | 03/30/2025 |
| 1 | | 12 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M007 | 12/30/2013 | 03/30/2025 |

Labeler - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 3/2024

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)