

CLONIDINE HYDROCHLORIDE- clonidine hydrochloride tablet
Asclemed USA, Inc.

Clonidine hydrochloride

Rx only


CLONIDINE HYDROCHLORIDE TABLETS USP

DESCRIPTION

Clonidine hydrochloride is a centrally acting alpha-agonist hypotensive agent available as tablets for oral administration in three dosage strengths: 0.1 mg, 0.2 mg and 0.3 mg. The 0.1 mg tablet is equivalent to 0.087 mg of the free base.

Clonidine hydrochloride tablets USP contain the following inactive ingredients: lactose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, and sodium starch glycolate. The 0.1 mg also contains D&C yellow #10 aluminum lake, and the 0.3 mg contains D&C yellow #10 aluminum lake and FD&C blue #1 aluminum lake.

Clonidine hydrochloride is an imidazoline derivative and exists as a mesomeric compound. The chemical name is 2-(2,6-dichlorophenylamino)-2-imidazoline hydrochloride. The following is the structural formula:



$C_9H_9Cl_2N_3 \cdot HCl$ Mol. Wt. 266.56

Clonidine hydrochloride is an odorless, bitter, white, crystalline substance soluble in water and alcohol.

CLINICAL PHARMACOLOGY

Clonidine stimulates alpha-adrenoreceptors in the brain stem. This action results in reduced sympathetic outflow from the central nervous system and in decreases in peripheral resistance, renal vascular resistance, heart rate, and blood pressure. Clonidine hydrochloride tablets USP act relatively rapidly. The patient's blood pressure declines within 30 to 60 minutes after an oral dose, the maximum decrease occurring within 2 to 4 hours. Renal blood flow and glomerular filtration rate remain essentially unchanged. Normal postural reflexes are intact; therefore, orthostatic symptoms are mild and infrequent.

Acute studies with clonidine hydrochloride in humans have demonstrated a moderate reduction (15% to 20%) of cardiac output in the supine position with no change in the peripheral resistance: at a 45° tilt there is a smaller reduction in cardiac output and a decrease of peripheral resistance. During long-term therapy, cardiac output tends to return to control values, while peripheral resistance remains decreased. Slowing of the pulse rate has been observed in most patients given clonidine, but the drug does not alter normal hemodynamic response to exercise.

Tolerance to the antihypertensive effect may develop in some patients, necessitating a reevaluation of therapy.

Other studies in patients have provided evidence of a reduction in plasma renin activity and in the excretion of aldosterone and catecholamines. The exact relationship of these pharmacologic actions to the antihypertensive effect of clonidine has not been fully elucidated.

Clonidine acutely stimulates growth hormone release in both children and adults, but does not produce a chronic elevation of growth hormone with long-term use.

Pharmacokinetics

The pharmacokinetics of clonidine is dose-proportional in the range of 100 to 600 mcg. The absolute bioavailability of clonidine on oral administration is 70% to 80%. Peak plasma clonidine levels are attained in approximately 1 to 3 hours.

Following intravenous administration, clonidine displays biphasic disposition with a distribution half-life of about 20 minutes and an elimination half-life ranging from 12 to 16 hours. The half-life increases up to 41 hours in patients with severe impairment of renal function. Clonidine crosses the placental barrier. It has been shown to cross the blood-brain barrier in rats.

Following oral administration about 40% to 60% of the absorbed dose is recovered in the urine as unchanged drug in 24 hours. About 50% of the absorbed dose is metabolized in the liver. Neither food nor the race of the patient influences the pharmacokinetics of clonidine.

The antihypertensive effect is reached at plasma concentrations between about 0.2 and 2.0 ng/mL in patients with normal excretory function. A further rise in the plasma levels will not enhance the antihypertensive effect.

INDICATIONS AND USAGE

Clonidine hydrochloride tablets USP are indicated in the treatment of hypertension. Clonidine hydrochloride tablets USP may be employed alone or concomitantly with other antihypertensive agents.

CONTRAINDICATIONS

Clonidine hydrochloride tablets USP should not be used in patients with known hypersensitivity to clonidine (see **PRECAUTIONS**).

WARNINGS

Withdrawal

Patients should be instructed not to discontinue therapy without consulting their physician. Sudden cessation of clonidine treatment has, in some cases, resulted in symptoms such as nervousness, agitation, headache, and tremor accompanied or followed by a rapid rise in blood pressure and elevated catecholamine concentrations in the plasma. The likelihood of such reactions to discontinuation of clonidine therapy appears to be greater after administration of higher doses or continuation of concomitant beta-blocker treatment and special caution is therefore advised in these situations. Rare instances of hypertensive encephalopathy, cerebrovascular accidents and death have been reported after clonidine withdrawal. When discontinuing therapy with clonidine hydrochloride tablets USP, the physician should reduce the dose gradually over 2 to 4 days to avoid withdrawal symptomatology.

An excessive rise in blood pressure following discontinuation of clonidine hydrochloride tablets USP therapy can be reversed by administration of oral clonidine hydrochloride or by intravenous phentolamine. If therapy is to be discontinued in patients receiving a beta-blocker and clonidine concurrently, the beta-blocker should be withdrawn several days before the gradual discontinuation of clonidine hydrochloride tablets USP.

Because children commonly have gastrointestinal illnesses that lead to vomiting, they may be particularly susceptible to hypertensive episodes resulting from abrupt inability to take medication.

PRECAUTIONS

General

In patients who have developed localized contact sensitization to clonidine-TTS, continuation of clonidine-TTS or substitution of oral clonidine hydrochloride therapy may be associated with the development of a generalized skin rash.

In patients who develop an allergic reaction to clonidine-TTS, substitution of oral clonidine hydrochloride may also elicit an allergic reaction (including generalized rash, urticaria, or angioedema).

The sympatholytic action of clonidine may worsen sinus node dysfunction and atrioventricular (AV) block, especially in patients taking other sympatholytic drugs. There are post-marketing reports of patients with conduction abnormalities and/or taking other sympatholytic drugs who developed severe bradycardia requiring IV atropine, IV isoproterenol and temporary cardiac pacing while taking clonidine.

In hypertension caused by pheochromocytoma, no therapeutic effect of clonidine hydrochloride tablets USP can be expected.

Perioperative Use

Administration of clonidine hydrochloride tablets USP should be continued to within

4 hours of surgery and resumed as soon as possible thereafter. Blood pressure should be carefully monitored during surgery and additional measures to control blood pressure should be available if required.

Information for Patients

Patients should be cautioned against interruption of clonidine hydrochloride tablets USP therapy without their physician's advice.

Since patients may experience a possible sedative effect, dizziness, or accommodation disorder with use of clonidine, caution patients about engaging in activities such as driving a vehicle or operating appliances or machinery. Also, inform patients that this sedative effect may be increased by concomitant use of alcohol, barbiturates, or other sedating drugs.

Patients who wear contact lenses should be cautioned that treatment with clonidine hydrochloride tablets USP may cause dryness of eyes.

Drug Interactions

Clonidine may potentiate the CNS-depressive effects of alcohol, barbiturates or other sedating drugs. If a patient receiving clonidine hydrochloride is also taking tricyclic antidepressants, the hypotensive effect of clonidine may be reduced, necessitating an increase in the clonidine dose. If a patient receiving clonidine is also taking neuroleptics, orthostatic regulation disturbances (e.g., orthostatic hypotension, dizziness, fatigue) may be induced or exacerbated.

Monitor heart rate in patients receiving clonidine concomitantly with agents known to affect sinus node function or AV nodal conduction, e.g., digitalis, calcium channel blockers and beta-blockers. Sinus bradycardia resulting in hospitalization and pacemaker insertion has been reported in association with the use of clonidine concomitantly with diltiazem or verapamil.

Amitriptyline in combination with clonidine enhances the manifestation of corneal lesions in rats (see **Toxicology**).

Based on observations in patients in a state of alcoholic delirium it has been suggested that high intravenous doses of clonidine may increase the arrhythmogenic potential (QT-prolongation, ventricular fibrillation) of high intravenous doses of haloperidol. Causal relationship and relevance for clonidine oral tablets have not been established.

Toxicology

In several studies with oral clonidine hydrochloride, a dose-dependent increase in the incidence and severity of spontaneous retinal degeneration was seen in albino rats treated for six months or longer. Tissue distribution studies in dogs and monkeys showed a concentration of clonidine in the choroid.

In view of the retinal degeneration seen in rats, eye examinations were performed during clinical trials in 908 patients before, and periodically after, the start of clonidine therapy. In 353 of these 908 patients, the eye examinations were carried out over periods of 24 months or longer. Except for some dryness of the eyes, no drug-related abnormal ophthalmological findings were recorded and, according to specialized tests such as electroretinography and macular dazzle, retinal function was unchanged.

In combination with amitriptyline, clonidine hydrochloride administration led to the development of corneal lesions in rats within 5 days.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Chronic dietary administration of clonidine was not carcinogenic to rats (132 weeks) or mice (78 weeks) dosed, respectively, at up to 46 or 70 times the maximum recommended daily human dose as mg/kg (9 or 6 times the MRDHD on a mg/m² basis). There was no evidence of genotoxicity in the Ames test for mutagenicity or mouse micronucleus test for clastogenicity.

Fertility of male or female rats was unaffected by clonidine doses as high as 150 mcg/kg (approximately 3 times MRDHD). In a separate experiment, fertility of female rats appeared to be affected at dose levels of 500 to 2000 mcg/kg (10 to 40 times the oral MRDHD on a mg/kg basis; 2 to 8 times the MRDHD on a mg/m² basis).

Pregnancy

Teratogenic Effects: Pregnancy Category C.

Reproduction studies performed in rabbits at doses up to approximately 3 times the oral maximum recommended daily human dose (MRDHD) of clonidine hydrochloride tablets USP produced no evidence of a teratogenic or embryotoxic potential in rabbits. In rats, however, doses as low as 1/3 the oral MRDHD (1/15 the MRDHD on a mg/m² basis) of clonidine were associated with increased resorptions in a study in which dams were treated continuously from 2 months prior to mating. Increased resorptions were not associated with treatment at the same time or at higher dose levels (up to 3 times the oral MRDHD) when the dams were treated on gestation days 6 to 15. Increases in resorption were observed at much higher dose levels (40 times the oral MRDHD on a mg/kg basis;

4 to 8 times the MRDHD on a mg/m² basis) in mice and rats treated on gestation days 1 to 14 (lowest dose employed in the study was 500 mcg/kg).

No adequate, well-controlled studies have been conducted in pregnant women. Clonidine crosses the placental barrier (see **CLINICAL PHARMACOLOGY, Pharmacokinetics**). Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers

As clonidine hydrochloride is excreted in human milk, caution should be exercised when clonidine hydrochloride tablets USP are administered to a nursing woman.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established in adequate and well-controlled trials (see **WARNINGS, Withdrawal**).

ADVERSE REACTIONS

Most adverse effects are mild and tend to diminish with continued therapy. The most

frequent (which appear to be dose-related) are dry mouth, occurring in about 40 of 100 patients; drowsiness, about 33 in 100; dizziness, about 16 in 100; constipation and sedation, each about 10 in 100.

The following less frequent adverse experiences have also been reported in patients receiving clonidine hydrochloride tablets USP, but in many cases patients were receiving concomitant medication and a causal relationship has not been established.

Body as a Whole: Fatigue, fever, headache, pallor, weakness, and withdrawal syndrome. Also reported were a weakly positive Coombs' test and increased sensitivity to alcohol.

Cardiovascular: Bradycardia, congestive heart failure, electrocardiographic abnormalities (i.e., sinus node arrest, junctional bradycardia, high degree AV block and arrhythmias), orthostatic symptoms, palpitations, Raynaud's phenomenon, syncope, and tachycardia. Cases of sinus bradycardia and atrioventricular block have been reported, both with and without the use of concomitant digitalis.

Central Nervous System: Agitation, anxiety, delirium, delusional perception, hallucinations (including visual and auditory), insomnia, mental depression, nervousness, other behavioral changes, paresthesia, restlessness, sleep disorder, and vivid dreams or nightmares.

Dermatological: Alopecia, angioneurotic edema, hives, pruritus, rash, and urticaria.

Gastrointestinal: Abdominal pain, anorexia, constipation, hepatitis, malaise, mild transient abnormalities in liver function tests, nausea, parotitis, pseudo-obstruction (including colonic pseudo-obstruction), salivary gland pain, and vomiting.

Genitourinary: Decreased sexual activity, difficulty in micturition, erectile dysfunction, loss of libido, nocturia, and urinary retention.

Hematologic: Thrombocytopenia.

Metabolic: Gynecomastia, transient elevation of blood glucose or serum creatine phosphokinase, and weight gain.

Musculoskeletal: Leg cramps and muscle or joint pain.

Oro-otolaryngeal: Dryness of the nasal mucosa.

Ophthalmological: Accommodation disorder, blurred vision, burning of the eyes, decreased lacrimation, and dryness of eyes.

OVERDOSAGE

Hypertension may develop early and may be followed by hypotension, bradycardia, respiratory depression, hypothermia, drowsiness, decreased or absent reflexes, weakness, irritability and miosis. The frequency of CNS depression may be higher in children than adults. Large overdoses may result in reversible cardiac conduction defects or dysrhythmias, apnea, coma and seizures. Signs and symptoms of overdose generally occur within 30 minutes to two hours after exposure. As little as 0.1 mg of clonidine has produced signs of toxicity in children.

There is no specific antidote for clonidine overdosage. Clonidine overdosage may result

in the rapid development of CNS depression; therefore, induction of vomiting with ipecac syrup is not recommended. Gastric lavage may be indicated following recent and/or large ingestions. Administration of activated charcoal and/or a cathartic may be beneficial. Supportive care may include atropine sulfate for bradycardia, intravenous fluids and/or vasopressor agents for hypotension and vasodilators for hypertension. Naloxone may be a useful adjunct for the management of clonidine-induced respiratory depression, hypotension and/or coma; blood pressure should be monitored since the administration of naloxone has occasionally resulted in paradoxical hypertension. Tolazoline administration has yielded inconsistent results and is not recommended as first-line therapy. Dialysis is not likely to significantly enhance the elimination of clonidine.

The largest overdose reported to date involved a 28-year old male who ingested 100 mg of clonidine hydrochloride powder. This patient developed hypertension followed by hypotension, bradycardia, apnea, hallucinations, semicoma, and premature ventricular contractions. The patient fully recovered after intensive treatment. Plasma clonidine levels were 60 ng/ml after 1 hour, 190 ng/ml after 1.5 hours, 370 ng/ml after 2 hours, and 120 ng/ml after 5.5 and 6.5 hours. In mice and rats, the oral LD₅₀ of clonidine is 206 and 465 mg/kg, respectively.

DOSAGE AND ADMINISTRATION

Adults

The dose of clonidine hydrochloride tablets USP must be adjusted according to the patient's individual blood pressure response. The following is a general guide to its administration.

Initial Dose

0.1 mg tablet twice daily (morning and bedtime). Elderly patients may benefit from a lower initial dose.

Maintenance Dose

Further increments of 0.1 mg per day may be made at weekly intervals if necessary until the desired response is achieved. Taking the larger portion of the oral daily dose at bedtime may minimize transient adjustment effects of dry mouth and drowsiness. The therapeutic doses most commonly employed have ranged from 0.2 mg to 0.6 mg per day given in divided doses. Studies have indicated that 2.4 mg is the maximum effective daily dose, but doses as high as this have rarely been employed.

Renal Impairment

Patients with renal impairment may benefit from a lower initial dose. Patients should be carefully monitored. Since only a minimal amount of clonidine is removed during routine hemodialysis, there is no need to give supplemental clonidine following dialysis.

HOW SUPPLIED

Clonidine hydrochloride tablets USP are supplied as follows:

Clonidine hydrochloride tablets USP 0.1 mg, yellow, round, debossed MP 657 on one

side and plain on the other side.

Bottles of 20	NDC 76420-283-20 (repackaged from NDC 52817-180-XX)
Bottles of 30	NDC 76420-283-30 (repackaged from NDC 52817-180-XX)
Bottles of 60	NDC 76420-283-60 (repackaged from NDC 52817-180-XX)
Bottles of 90	NDC 76420-283-90 (repackaged from NDC 52817-180-XX)
Bottles of 100	NDC 76420-283-01 (relabeled from NDC 52817-180-10)

Clonidine hydrochloride tablets USP 0.2 mg, white, round, debossed MP 658 on one side and plain on the other side.

Bottles of 20	NDC 76420-285-20 (repackaged from NDC 52817-181-XX)
Bottles of 30	NDC 76420-285-30 (repackaged from NDC 52817-181-XX)
Bottles of 60	NDC 76420-285-60 (repackaged from NDC 52817-181-XX)
Bottles of 90	NDC 76420-285-90 (repackaged from NDC 52817-181-XX)
Bottles of 100	NDC 76420-285-01 (relabeled from NDC 52817-181-10)

Clonidine hydrochloride tablets USP 0.3 mg, green, round, debossed MP 659 on one side and plain on the other side.

Bottles of 20	NDC 76420-286-20 (repackaged from NDC 52817-182-XX)
Bottles of 30	NDC 76420-286-30 (repackaged from NDC 52817-182-XX)
Bottles of 60	NDC 76420-286-60 (repackaged from NDC 52817-182-XX)
Bottles of 90	NDC 76420-286-90 (repackaged from NDC 52817-182-XX)
Bottles of 100	NDC 76420-286-01 (relabeled from NDC 52817-182-10)

Store at 20° to 25°C (68° to 77°F).

[See USP Controlled Room Temperature]

DISPENSE IN TIGHT, LIGHT-RESISTANT CONTAINER.


Address medical inquiries to: 1-877-541-5504.

Relabeled and Repackaged by:

Enovachem PHARMACEUTICALS

Torrance, CA 90501

PRINCIPAL DISPLAY PANEL

Relabeled By:
 **Enovachem**™ 379 Van Ness Ave.
PHARMACEUTICALS Suite 1403-1406
Torrance, CA 90501
Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-20
Qty: 20
Distributed By: TruPharma, LLC
Source NDC: 52817-180-XX
Description: yellow/round/debossed; MP 657



(01)0 0376420 28320 1
(17)

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-20
S/N:
Qty: 20

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-20
S/N:
Qty: 20

Lot #: 00000000 Exp: (10) 00000000
Batch #: 00000000 (21)
Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-20
S/N:
Qty: 20

Relabeled By:

 **Enovachem**TM 379 Van Ness Ave.
PHARMACEUTICALS Suite 1403-1406
Torrance, CA 90501
Clonidine Hydrochloride Tablets, USP 0.1 mg

NDC: 76420-283-30

Qty: 30

Distributed By: TruPharma, LLC
Source NDC: 52817-180-XX
Description: yellow/round/debossed; MP 657
Lot #: 00000000 Exp:
Batch #: 00000000
Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01)0 0376420 28330 0
(17)
(10) 00000000
(21)

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-30
S/N:
Qty: 30

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-30
S/N:
Qty: 30

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-30
S/N:
Qty: 30

Relabeled By:

 **Enovachem**TM 379 Van Ness Ave.
PHARMACEUTICALS Suite 1403-1406
Torrance, CA 90501
Clonidine Hydrochloride Tablets, USP 0.1 mg

NDC: 76420-283-60

Qty: 60

Distributed By: TruPharma, LLC
Source NDC: 52817-180-XX
Description: yellow/round/debossed; MP 657
Lot #: 00000000 Exp:
Batch #: 00000000
Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01)0 0376420 28360 7
(17)
(10) 00000000
(21)

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-60
S/N:
Qty: 60

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-60
S/N:
Qty: 60

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-60
S/N:
Qty: 60

Relabeled By:

 **Enovachem**TM 379 Van Ness Ave.
PHARMACEUTICALS Suite 1403-1406
Torrance, CA 90501
Clonidine Hydrochloride Tablets, USP 0.1 mg

NDC: 76420-283-90

Qty: 90

Distributed By: TruPharma, LLC
Source NDC: 52817-180-XX
Description: yellow/round/debossed; MP 657
Lot #: 00000000 Exp:
Batch #: 00000000
Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01)0 0376420 28390 4
(17)
(10) 00000000
(21)

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-90
S/N:
Qty: 90

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-90
S/N:
Qty: 90

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-90
S/N:
Qty: 90

Relabeled By:

 **Enovachem**TM 379 Van Ness Ave.
PHARMACEUTICALS Suite 1403-1406
Torrance, CA 90501
Clonidine Hydrochloride Tablets, USP 0.1 mg

NDC: 76420-283-01

Qty: 100

Distributed By: TruPharma, LLC
Source NDC: 52817-180-10
Description: yellow/round/debossed; MP 657
Lot #: 00000000 Exp:
Batch #: 00000000
Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01)0 0376420 28301 0
(17)
(10) 00000000
(21)

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-01
S/N:
Qty: 100

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-01
S/N:
Qty: 100

Clonidine Hydrochloride Tablets, USP 0.1 mg
NDC: 76420-283-01
S/N:
Qty: 100

PRINCIPAL DISPLAY PANEL

Packaged By:



Enovachem
PHARMACEUTICALS

379 Van Ness Ave.
Suite 1403-1406
Torrance, CA 90501

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-20

Qty: 20

Distributed By: TruPharma, LLC

Source NDC: 52817-181-XX

Description: white/round/debossed;MP 658

Lot #: 00000000

Exp:

Batch #: 00000000

Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01) 0 0376420 28520 5
(17)
(10) 00000000
(21)

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-20

S/N:

Qty: 20

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-20

S/N:

Qty: 20

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-20

S/N:

Qty: 20

Packaged By:



Enovachem
PHARMACEUTICALS

379 Van Ness Ave.
Suite 1403-1406
Torrance, CA 90501

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-30

Qty: 30

Distributed By: TruPharma, LLC

Source NDC: 52817-181-XX

Description: white/round/debossed;MP 658

Lot #: 00000000

Exp:

Batch #: 00000000

Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01) 0 0376420 28530 4
(17)
(10) 00000000
(21)

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-30

S/N:

Qty: 30

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-30

S/N:

Qty: 30

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-30

S/N:

Qty: 30

Packaged By:



Enovachem
PHARMACEUTICALS

379 Van Ness Ave.
Suite 1403-1406
Torrance, CA 90501

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-60

Qty: 60

Distributed By: TruPharma, LLC

Source NDC: 52817-181-XX

Description: white/round/debossed;MP 658

Lot #: 00000000

Exp:

Batch #: 00000000

Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01) 0 0376420 28560 1
(17)
(10) 00000000
(21)

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-60

S/N:

Qty: 60

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-60

S/N:

Qty: 60

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-60

S/N:

Qty: 60

Packaged By:



Enovachem
PHARMACEUTICALS

379 Van Ness Ave.
Suite 1403-1406
Torrance, CA 90501

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-90

Qty: 90

Distributed By: TruPharma, LLC



Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-90

S/N:

Qty: 90

Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-90

S/N:

Qty: 90

Source NDC: 52817-181-XX
Description: white/round/debossed;MP 658
Lot #: 00000000 Exp:
Batch #: 00000000
Drug Status: RX

(01) 0 0376420 28590 8
(17)
(10) 00000000
(21)

Qty: 90

Clonidine Hydrochloride Tablets, USP 0.2 mg
NDC: 76420-285-90
S/N:
Qty: 90

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

Relabeled By:

 **Enovachem**[™] 379 Van Ness Ave.
PHARMACEUTICALS Suite 1403-1406
Torrance, CA 90501
Clonidine Hydrochloride Tablets, USP 0.2 mg

NDC: 76420-285-01
Qty: 100



(01) 0 0376420 28501 4
(17)
(10) 00000000
(21)

Distributed By: TruPharma, LLC
Source NDC: 52817-181-10
Description: white/round/debossed;MP 658
Lot #: 00000000 Exp:
Batch #: 00000000
Drug Status: RX

Clonidine Hydrochloride Tablets, USP 0.2 mg
NDC: 76420-285-01
S/N:
Qty: 100

Clonidine Hydrochloride Tablets, USP 0.2 mg
NDC: 76420-285-01
S/N:
Qty: 100

Clonidine Hydrochloride Tablets, USP 0.2 mg
NDC: 76420-285-01
S/N:
Qty: 100

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

PRINCIPAL DISPLAY PANEL

Packaged By:

 **Enovachem**[™] 379 Van Ness Ave.
PHARMACEUTICALS Suite 1403-1406
Torrance, CA 90501
Clonidine Hydrochloride Tablets, USP 0.3 mg

NDC: 76420-286-20
Qty: 20



(01) 0 0376420 28620 2
(17)
(10) 00000000
(21)

Distributed By: TruPharma, LLC
Source NDC: 52817-182-XX
Description: green/round/debossed;MP 659
Lot #: 00000000 Exp:
Batch #: 00000000
Drug Status: RX

Clonidine Hydrochloride Tablets, USP 0.3 mg
NDC: 76420-286-20
S/N:
Qty: 20

Clonidine Hydrochloride Tablets, USP 0.3 mg
NDC: 76420-286-20
S/N:
Qty: 20

Clonidine Hydrochloride Tablets, USP 0.3 mg
NDC: 76420-286-20
S/N:
Qty: 20

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

Packaged By:

 **Enovachem**[™] 379 Van Ness Ave.
PHARMACEUTICALS Suite 1403-1406
Torrance, CA 90501
Clonidine Hydrochloride Tablets, USP 0.3 mg

NDC: 76420-286-30
Qty: 30



(01) 0 0376420 28630 1
(17)
(10) 00000000
(21)

Distributed By: TruPharma, LLC
Source NDC: 52817-182-XX
Description: green/round/debossed;MP 659
Lot #: 00000000 Exp:
Batch #: 00000000
Drug Status: RX

Clonidine Hydrochloride Tablets, USP 0.3 mg
NDC: 76420-286-30
S/N:
Qty: 30

Clonidine Hydrochloride Tablets, USP 0.3 mg
NDC: 76420-286-30
S/N:
Qty: 30

Clonidine Hydrochloride Tablets, USP 0.3 mg
NDC: 76420-286-30
S/N:
Qty: 30

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

Packaged By:

 **Enovachem**[™]

Clonidine Hydrochloride Tablets, USP 0.3 mg
NDC: 76420-286-30

Enovachem[™] 379 Van Ness Ave.
 PHARMACEUTICALS Suite 1403-1406
 Torrance, CA 90501
 Clonidine Hydrochloride Tablets, USP 0.3 mg

NDC: 76420-286-60

Qty: 60

Distributed By: TruPharma, LLC
 Source NDC: 52817-182-XX
 Description: green/round/debossed;MP 659
 Lot #: 00000000 Exp:
 Batch #: 00000000
 Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
 KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01) 0 0376420 28660 8
 (17)
 (10) 00000000
 (21)

NDC: 76420-286-60

S/N:
 Qty: 60

Clonidine Hydrochloride Tablets, USP 0.3 mg

NDC: 76420-286-60
 S/N:
 Qty: 60

Clonidine Hydrochloride Tablets, USP 0.3 mg

NDC: 76420-286-60
 S/N:
 Qty: 60

Packaged By:

Enovachem[™] 379 Van Ness Ave.
 PHARMACEUTICALS Suite 1403-1406
 Torrance, CA 90501
 Clonidine Hydrochloride Tablets, USP 0.3 mg

NDC: 76420-286-90

Qty: 90

Distributed By: TruPharma, LLC
 Source NDC: 52817-182-XX
 Description: green/round/debossed;MP 659
 Lot #: 00000000 Exp:
 Batch #: 00000000
 Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
 KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01) 0 0376420 28690 5
 (17)
 (10) 00000000
 (21)

Clonidine Hydrochloride Tablets, USP 0.3 mg

NDC: 76420-286-90
 S/N:
 Qty: 90

Clonidine Hydrochloride Tablets, USP 0.3 mg

NDC: 76420-286-90
 S/N:
 Qty: 90

Clonidine Hydrochloride Tablets, USP 0.3 mg

NDC: 76420-286-90
 S/N:
 Qty: 90

Relabeled By:

Enovachem[™] 379 Van Ness Ave.
 PHARMACEUTICALS Suite 1403-1406
 Torrance, CA 90501
 Clonidine Hydrochloride Tablets, USP 0.3 mg

NDC: 76420-286-01

Qty: 100

Distributed By: TruPharma, LLC
 Source NDC: 52817-182-10
 Description: green/round/debossed;MP 659
 Lot #: 00000000 Exp:
 Batch #: 00000000
 Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
 KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01) 0 0376420 28601 1
 (17) -----
 (10) 00000000
 (21)

Clonidine Hydrochloride Tablets, USP 0.3 mg

NDC: 76420-286-01
 S/N:
 Qty: 100

Clonidine Hydrochloride Tablets, USP 0.3 mg

NDC: 76420-286-01
 S/N:
 Qty: 100

Clonidine Hydrochloride Tablets, USP 0.3 mg

NDC: 76420-286-01
 S/N:
 Qty: 100

CLONIDINE HYDROCHLORIDE

clonidine hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:76420-283(NDC:52817-180)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
CLONIDINE HYDROCHLORIDE (UNII: W7616XXF06) (CLONIDINE - UNII:MN3L5RMN02)		CLONIDINE HYDROCHLORIDE	0.1 mg	
Inactive Ingredients				
Ingredient Name			Strength	
MAGNESIUM STEARATE (UNII: 70097M6I30)				
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
STARCH, CORN (UNII: O8232NY3SJ)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
Product Characteristics				
Color	yellow	Score	no score	
Shape	ROUND	Size	7mm	
Flavor		Imprint Code	MP;657	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76420-283-20	20 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	
2	NDC:76420-283-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	
3	NDC:76420-283-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	
4	NDC:76420-283-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	09/09/2022	
5	NDC:76420-283-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA070925	06/01/2017		

CLONIDINE HYDROCHLORIDE

clonidine hydrochloride tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:76420-285(NDC:52817-181)

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CLONIDINE HYDROCHLORIDE (UNII: W7616XXF06) (CLONIDINE - UNII:MN3L5RMN02)	CLONIDINE HYDROCHLORIDE	0.2 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	MP;658
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76420-285-20	20 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	
2	NDC:76420-285-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	
3	NDC:76420-285-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	
4	NDC:76420-285-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	
5	NDC:76420-285-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA070924	06/01/2017	

CLONIDINE HYDROCHLORIDE

clonidine hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:76420-286(NDC:52817-182)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CLONIDINE HYDROCHLORIDE (UNII: W7616XXF06) (CLONIDINE - UNII:MN3L5RMN02)	CLONIDINE HYDROCHLORIDE	0.3 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
ALUMINUM OXIDE (UNII: LMI26O6933)	

Product Characteristics

Color	green	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	MP;659
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76420-286-20	20 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	
2	NDC:76420-286-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	
3	NDC:76420-286-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	
4	NDC:76420-286-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	
5	NDC:76420-286-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/16/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA070923	06/01/2017	

Labeler - Asclemed USA, Inc. (059888437)

Establishment

Name	Address	ID/FEI	Business Operations
ASCLEMED USA INC. DBA ENOVACHEM		059888437	relabel(76420-283, 76420-285, 76420-286) , repack(76420-283, 76420-285, 76420-286)

Revised: 9/2022

Asclemed USA, Inc.