

**DAYTIME NIGHTTIME COLD AND FLU RELIEF- acetaminophen,
dextromethorphan hbr, doxylamine succinate, phenylephrine hcl
CVS Pharmacy**

Daytime Nighttime Cold & Flu Relief Drug Facts

Nighttime Cold & Flu

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL dose cup contains:** sodium 39 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments?

1-800-719-9260

Daytime Cold & Flu

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure

- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:** sodium 7 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

COMBO PACK

CVS Health®

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu

CVS Health®

Non-Drowsy

Daytime

ACETAMINOPHEN - Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr - Cough suppressant

PHENYLEPHRINE HCl - Nasal decongestant

Relieves:

Aches, Fever, Sore throat, Cough, Nasal congestion

COLD & FLU

ALCOHOL & ANTIHISTAMINE FREE

TOTAL 24 FL OZ (1.5 PT) (710 mL)

COMBO PACK

CVS Health®

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu

CVS Health®

Multi-Symptom

Nighttime

ACETAMINOPHEN - Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr - Cough suppressant

DOXYLAMINE SUCCINATE - Antihistamine

Relieves:

Aches, Fever, Sore throat, Cough, Sneezing, Runny nose

COLD & FLU

Cherry Flavor

ALCOHOL 10%

TWO 12 FL OZ (355 mL) UNITS

CVS Health.

Non-Drowsy
Daytime

Multi-Symptom
Nighttime

CVS Health.

COMBO PACK

CVS Health.

CVS Health.

Non-Drowsy
Daytime

ACETAMINOPHEN-
Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr-
Cough suppressant
PHENYLEPHRINE HCl-
Nasal decongestant



NDC 69842-733-02

Multi-Symptom
Nighttime

ACETAMINOPHEN-
Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr-
Cough suppressant
DOXYLAMINE SUCCINATE-
Antihistamine

Relieves: Aches, Fever, Sore throat, Cough,
Nasal congestion

Relieves: Aches, Fever, Sore throat, Cough,
Sneezing, Runny nose

COLD & FLU

ALCOHOL &
ANTIHISTAMINE FREE

TOTAL 24 FL OZ
(1.5 pt) (710 mL)

COLD & FLU

Cherry Flavor
ALCOHOL 10%

TWO 12 FL OZ
(355 mL) UNITS

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2022 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16430

100% money back
guaranteed.
CVS.com/returnpolicy

*These products are not manufactured or distributed
by Procter & Gamble, distributor of Vicks® DayQuil®
Cold & Flu and Vicks® NyQuil® Cold & Flu.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

DO NOT USE IF PRINTED NECKBAND
IS BROKEN OR MISSING



#180153

73302 17 C4

Nighttime Cold & Flu

Drug Facts

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

Uses temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat ■ headache ■ minor aches and pains
- fever ■ runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease ■ glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur ■ avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed — see Overdose warning
 - only use the dose cup provided ■ do not exceed 4 doses per 24 hrs
- | | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 6 hrs |
| children 4 to under 12 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- each 30 mL contains: sodium 39 mg
- store at 20-25°C (68-77°F)

Inactive ingredients alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-800-719-9260

Daytime Cold & Flu

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion ■ cough due to minor throat and bronchial irritation
- sore throat ■ headache ■ minor aches and pains ■ fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen

■ adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease ■ heart disease ■ high blood pressure
- thyroid disease ■ diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions


- take only as directed — see Overdose warning
 - only use the dose cup provided ■ do not exceed 4 doses per 24 hrs
- | | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs | 15 mL every 4 hrs |
| children 4 to under 6 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- each 15 mL contains: sodium 7 mg ■ store at 20-25°C (68-77°F)

Inactive ingredients butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments? 1-800-719-9260



DAYTIME NIGHTTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-733
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-733-02	1 in 1 CARTON; Type 0: Not a Combination Product	06/23/2016	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	355 mL
Part 2	1 BOTTLE	355 mL

Part 1 of 2

NIGHTTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Product Information

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL

DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	

Product Characteristics

Color	RED (Clear/Dark Red)	Score	
Shape		Size	
Flavor	CHERRY (Menthol Aroma)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/23/2016	

Part 2 of 2

DAYTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, phenylephrine hcl solution

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A)	

Product Characteristics

Color	ORANGE (clear)	Score	
Shape		Size	
Flavor	MENTHOL (with fruit)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/23/2016	

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
OTC Monograph Drug	M012	06/23/2016	

Labeler - CVS Pharmacy (062312574)

Revised: 11/2023

CVS Pharmacy