

**TUSSIN DM COUGH AND CHEST CONGESTION- dextromethorphan hydrobromide,
guaifenesin liquid
DeMoulas Market Basket**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Tussin DM 503

Drug Facts

**Active ingredients
(in each 5 mL tsp)**

Dextromethorphan HBr, USP 10 mg

Guaifenesin, USP 100 mg

Purpose

Cough Suppressant

Expectorant

Keep out of reach of children

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use

- in a child under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.

These could be signs of a serious condition.

If pregnant or breast-feeding

ask a health professional before use.

Directions

- do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years & over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

Other information

- Keep carton for full Direction for use.
- store at 20-25 ° C (68-77 ° F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

Inactive ingredients

anhydrous citric acid, dextrose, FD and C red no. 40, flavor, glycerin, high fructose corn syrup, purified water, saccharin sodium, sodium benzoate

Questions?

Call weekdays from 9:30 AM to 4:30 PM EST at

1-877-798-5944

Product Label

"MORE FOR YOUR DOLLAR"

MARKET BASKET®

***COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH and CHEST CONGESTION DM**

Tussin DM

Cough and Chest Congestion

Dextromethorphan HBr
Guaifenesin

COUGH SUPPRESSANT /
EXPECTORANT

Helps to Loosen Chest Congestion

NON-DROWSY

ADULT
Cough Formula

for ages 12 and over

8 FL OZ (237 mL)

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN

*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

Distributed by: DEMOULAS SUPERMARKETS INC.
875 EAST STREET - TEWKSBURY, MA 01876

BX-012 REV 01

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Drug Facts

Active ingredients (in each 5mL tsp)
 Dextromethorphan HBr, USP 10mg Cough Suppressant
 Guaifenesin, USP 100mg Expectorant

Uses
 ■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings
 Do not use ■ in a child under 12 years of age
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before using this product.

Ask a doctor before use if you have
 ■ cough that occurs with too much phlegm (mucus)
 ■ cough that lasts or is chronic, such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not take more than 6 doses in any 24-hour period.

Age	Dose
Adults and children 12 years and over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

Other information
 ■ store at 20-25°C (68-77°F) ■ do not refrigerate
 ■ dosage cup provided ■ sodium 3 mg per teaspoonful
 ■ Keep carton for full directions for use.

Inactive ingredients Anhydrous citric acid, dextrose, FD&C red #40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold. DISTRIBUTED BY DEMOULAS SUPERMARKETS INC. 875 EAST STREET TEWKSBURY, MA 01876

BX-012
REV 01



Tussin DM
Cough & Chest Congestion

Dextromethorphan HBr
Guaifenesin

Cough Suppressant/
Expectorant

Helps Loosen Chest
Congestion

NON-DROWSY

ADULT
Cough Formula
for ages 12
& over

8 FL OZ (237 mL)



Tussin DM
Cough & Chest Congestion

Dextromethorphan HBr
Guaifenesin

Cough Suppressant/
Expectorant

Helps Loosen Chest
Congestion

NON-DROWSY

ADULT
Cough Formula
for ages 12
& over

8 FL OZ (237 mL)

*COMPARE TO the active ingredients in Robitussin® Peak Cold Cough & Chest Congestion DM



ADULT Cough Formula for ages 12 & over

NON-DROWSY

Tussin DM
Cough & Chest Congestion

Dextromethorphan HBr / Guaifenesin
Cough Suppressant/Expectorant
Helps Loosen Chest Congestion

8 FL OZ (237 mL)

MARKET BASKET
MORE FOR YOUR DOLLAR

*COMPARE TO the active ingredients in Robitussin® Peak Cold Cough & Chest Congestion DM

Drug Facts

Active ingredients (in each 5 mL tsp)
 Dextromethorphan HBr, USP 10 mg
 Guaifenesin, USP 100 mg

Purpose
 Cough Suppressant
 Expectorant

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Guaifenesin, USP 100 mg Cough Suppressant
 Expectorant

Uses ■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use ■ in a child under 12 years of age
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

■ cough that occurs with too much phlegm (mucus)
 ■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years & over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

Other information

■ store at 20-25°C (68-77°F) ■ do not refrigerate.
 ■ dosage cup provided ■ sodium 3 mg per teaspoonful
 ■ See carton for full labeling

Inactive ingredients anhydrous citric acid, dextrose, FD&C red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944.

*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

DISTRIBUTED BY
DEMOULAS SUPERMARKETS INC.
 875 East Street
 TEWKSBURY, MA 01876

LOT:

LR-031

EXP:

Rev 01

TUSSIN DM COUGH AND CHEST CONGESTION

dextromethorphan hydrobromide, guaifenesin liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53942-503
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL

GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)

GUAIFENESIN

100 mg
in 5 mL**Inactive Ingredients**

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
DEXTROSE (UNII: IY9XDZ35W2)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
MENTHOL (UNII: L7T10EIP3A)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53942-503-26	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2012	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/01/2012	

Labeler - DeMoulas Market Basket (007869647)**Registrant** - AptaPharma Inc. (790523323)**Establishment**

Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(53942-503)

Revised: 4/2017

DeMoulas Market Basket