

**FOSTER AND THRIVE ASPIRIN- aspirin tablet, delayed release
McKesson Medical-Surgical, Inc.**

Adult Low Dose Aspirin

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- temporary relief of minor aches and pains or as recommended by your doctor.
- ask your doctor about other uses for this product.

Because of its delayed release action, this product will not provide fast relief of headache or symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives • shock • facial swelling • asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Do not use

- if you are allergic to aspirin, or any other pain reliever/fever reducer • if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if

- You have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
 - an allergic reaction occurs. Seek medical help right away.
 - ringing in the ears or a loss of hearing occurs
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - new symptoms appear. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- drink a full glass of water with each dose
- **adults and children 12 years and over:** take 4 to 8 tablets every 4 hours while symptoms last • do not take more than 48 tablets in 24 hours unless directed by doctor • Swallow whole. Do not chew or crush tablets.
- **children under 12 years:** ask a doctor

Other information

store at room temperature 15-30°C (59-86°F)

Inactive ingredients

Colloidal silicon dioxide, croscarmellose sodium, D&C yellow 10, iron oxide yellow 6, lactose, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, sodium hydroxide, talc, titanium dioxide, triethyl citrate

Questions and comments?

Call 833-358-6431

This product is not manufactured or distributed by Bayer, owner of the registered trademark Aspirin Regimen BAYER®

© 2023 Northstar Healthcare UC

Distributed by:

McKesson Corp., via SSSL., Memphis, TN 38141

Money Back Guarantee

www.fosterandthrive.com

Product of India.

Mfg. Lic. No.: G/25/1362

Issued: 12/2025

Principal Display Panel

COMPARE TO BAYER® LOW DOSE ASPIRIN ACTIVE INGREDIENT†

Foster & Thrive®

ADULT LOW DOSE

Aspirin 81 mg

Pain Reliever (NSAID)

Adult aspirin regimen# use | enteric-coated

ACTUAL SIZE

Talk to your doctor or other healthcare provider before using this product for your heart.

Aspirin is not right for everyone.

This product is not manufactured or distributed by Bayer, owner of registered trademark Aspirin Regimen BAYER®

NDC 70677-1310-1 COMPARE TO BAYER® LOW DOSE ASPIRIN ACTIVE INGREDIENT†

Foster & Thrive®

ADULT LOW DOSE

Aspirin 81 mg
PAIN RELIEVER (NSAID)

Adult aspirin regimen# use | enteric-coated

ACTUAL SIZE

1,000 TABLETS

DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Aspirin 81 mg (NSAID)*</p> <p>Purpose Pain reliever *nonsteroidal anti-inflammatory drug</p> <p>Uses ■ temporary relief of minor aches and pains or as recommended by your doctor. ■ ask your doctor about other uses for this product</p> <p>Because of its delayed release action, this product will not provide fast relief of headache or symptoms needing immediate relief.</p> <p>Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ shock ■ facial swelling ■ asthma (wheezing)</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed</p> <p>Do not use ■ if you are allergic to aspirin, or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients.</p> <p>Ask a doctor before use if ■ You have asthma ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis or kidney disease ■ you are taking a diuretic</p>	<p>Drug Facts (continue)</p> <p>Ask a doctor or pharmacist before use if you are ■ taking a prescription drug for diabetes, gout, or arthritis</p> <p>Stop use and ask a doctor if</p> <p>Uses ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ have bloody or black stools ■ vomit blood ■ have stomach pain that does not get better ■ an allergic reaction occurs. Seek medical help right away.</p> <p>■ ringing in the ears or loss of hearing occurs ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ redness or swelling is present in the painful area ■ new symptoms appear. These could be sign of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions ■ drink a full glass of water with each dose ■ adults and children 12 years and over: ■ take 4 to 8 tablets every 4 hours while symptoms last ■ do not take more than 48 tablets in 24 hours unless directed by doctor ■ swallow whole. Do not chew or crush tablets.</p> <p>children under 12 years: ask a doctor</p> <p>Other information store at room temperature 15-30°C (59-86°F)</p> <p>Inactive ingredients: Colloidal silicon dioxide, croscarmellose sodium, D&C yellow 10, iron oxide yellow 6, lactose, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, poly sorbate 80, sodium hydroxide, talc, titanium dioxide, triethyl citrate</p> <p>Questions and comments? Call 833-358-6431</p>
---	--

† Talk to your doctor or other healthcare provider before using this product for your heart.

Aspirin is not right for everyone.
This product is not manufactured or distributed by Bayer, owner of the registered trademark Aspirin Regimen BAYER®
© 2023 Northstar Healthcare UC
Distributed by:
McKesson Corp., via SSSL., Memphis, TN 38141
Money Back Guarantee
www.fosterandthrive.com
Product of India.
Mfg. Lic. No.: G/25/1362
Issued: 12/2025

0 110939 196200 3

Unvarnish Area
50x30 mm

FOSTER AND THRIVE ASPIRIN

aspirin tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1310
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
LACTOSE (UNII: J2B2A4N98G)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	81
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1310-1	1000 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
---------------------------	---	-----------------------------	---------------------------

OTC Monograph Drug	M013	04/01/2026	
--------------------	------	------------	--

Labeler - McKesson Medical-Surgical, Inc. (116956644)

Registrant - Elysium Pharmaceuticals Ltd (915664486)

Establishment

Name	Address	ID/FEI	Business Operations
Elysium Pharmaceuticals Ltd		915664486	manufacture(70677-1310) , analysis(70677-1310)

Establishment

Name	Address	ID/FEI	Business Operations
Elysium Pharmaceuticals Ltd		863182240	label(70677-1310) , pack(70677-1310)

Revised: 10/2024

McKesson Medical-Surgical, Inc.