

**DOUBLE DINE PLUS- iodine liquid
IBA**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Double Dine Plus NPE Free

ACTIVE INGREDIENTS

ACTIVE INGREDIENT:

CHLORHEXIDINE GLUCONATE.....0.50%
IODINE.....0.50%
(PROVIDES A Full .5% AVAILABLE IODINE AS USED)

Inactive Ingredients

INERT INGREDIENTS:

GLYCERIN(AS A HUMECTANT AND SKIN CONDITIONER).....10.00%
OTHER INERTS.....89.00%
(INCLUDES BUFFERING AGENTS AND SURFACTANTS)
TOTAL:.....100.00%

THIS PRODUCT CONTAINS HIGHLY EFFECTIVE LEVELS OF
FREE IODINE FOR FAST KILL OF MASTITIS CAUSING ORGANISMS.

INDICATIONS AND USAGE

GENERAL RECOMMENDATIONS

- 1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
- 2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
- 3. Do not use this product for cleaning or sanitizing equipment
- 4. Wash Teat Dip cups after each milking.

RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

- 1. REMOVE VISIBLE SOILS FROM THE TEATS.
- 2. OBSERVE FOREMILK BY STRIPPING TWO OR THREE STREAMS OF MILK INTO A STRIP CUP.
- 3. DIP OR SPRAY THE COW'S TEATS FULL LENGTH WITH DOUBLE DINE PLUS TEAT DIP.
- 4 WAIT APPROXIMATELY 30 SECONDS.
- 5. REMOVE ALL TEAT DIP BY THOROUGHLY DRYING THE TEATS AND TEAT ORIFICES WITH A CLEAN, SINGLE SERVICE PAPER TOWEL(S).
- 6. ATTACH MILKING UNIT.

B. POST-MILKING (POST-DIPPING)

Immediately after milking dip all teats full length with DOUBLE DINE PLUS Teat Dip.

Ensure good coverage of all sides of the teats and teat orifices.

WARNINGS

DO NOT DILUTE THIS PRODUCT.

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.

ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS. PROTECT THIS PRODUCT FROM FREEZING.

CAUTION: KEEP OUT OF REACH OF CHILDREN. AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR READY TO USE PRODUCTS, WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

**ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300**

Double Dine Plus

Double Dine Plus

NDC 29983-1058-9



DOUBLE DINE PLUS

SANITIZING TEAT DIP

Contains Two Powerful And Time Tested Germicides For Quick And Effective Kill Of Contagious And Environmental Bacteria
 A Revolutionary Advance In Teat Dip Technology By IBA That Allows The Combination Of Chlorhexidine
 And Iodine To Remain Stable For The Shelf Life Of The Product
ESPECIALLY FORMULATED FOR DAIRIES WHERE NPE FREE PRODUCTS ARE PREFERRED OR REQUIRED

ACTIVE INGREDIENTS:

Chlorhexidine Gluconate 0.50%
 Iodine 0.50%
 (Provides A Full .5% Available Iodine As Used)

INERT INGREDIENTS:

Glycerin (As A Humectant And Skin Conditioner) 10.00%
 Other Inerts 89.00%
 (Includes Buffering Agents, Stabilizing Agents and Surfactants)
TOTAL..... 100.00%

This product contains highly effective levels of free iodine for fast kill of mastitis causing organisms.

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. If solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.


FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS, KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION FOR MEDICAL EMERGENCIES CALL 1-800-424-9300



U.S. Patent Numbers
 9,913,859 and 11,766,452

RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

1. Remove visible soils from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip, foam or spray the cow's teats full length with DOUBLE DINE PLUS Teat Dip.
4. Wait approximately 90 seconds.
5. Remove all teat dip by thoroughly drying the teats, teat orifices and udder (if foaming or spraying) with a clean, single service paper towel(s) to avoid contamination of milk.
6. Attach milking unit.

B. POST-MILKING (POST-DIPPING)

Immediately after milking, dip or spray all teats full length with DOUBLE DINE PLUS Teat Dip. Ensure good coverage of all sides of the teats and teat orifices.

CAUTION: KEEP OUT OF REACH OF CHILDREN. AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

WARNING

- DO NOT DILUTE THIS PRODUCT.
- DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.
- ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.
- PROTECT THIS PRODUCT FROM FREEZING.

A PART OF THE COMPLETE IBA SANITATION PROGRAM

EXP. DATE: LOT NO.: NET CONTENTS: 275 GALLONS (1,040.875 LITERS) IBA STOCK # 260167 1223W

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 103 GILMORE DRIVE, SUTTON, MA 01590

DOUBLE DINE PLUS

iodine liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-1058
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	0.5 kg in 100 kg
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	0.5 kg in 100 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-1058-9	1081.47 kg in 1 CONTAINER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		11/22/2024	

Labeler - IBA (019494160)

Registrant - IBA (019494160)

Establishment

Name	Address	ID/FEI	Business Operations
Morgan Gallacher Inc. DBA, Custom Chemical Formulators Inc.		028311595	manufacture, api manufacture, pack

Revised: 11/2024

IBA