

**LOXICOM- meloxicam suspension**  
**Norbrook Laboratories Limited**

-----  
**Professional Insert**

**Loxicom<sup>®</sup>**  
**(meloxicam oral suspension)**  
**0.5 mg/mL**


Non-steroidal anti-inflammatory drug for oral use in dogs only

**Caution:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**Warning:** Repeated use of meloxicam in cats has been associated with acute renal failure and death. Do not administer additional injectable or oral meloxicam to cats. See Contraindications, Warnings, and Precautions for detailed information.

**Description:**

Meloxicam is a non-steroidal anti-inflammatory drug (NSAID) of the oxicam class. Each milliliter of Loxicom<sup>®</sup> oral suspension contains meloxicam equivalent to 0.5 milligrams and sodium benzoate (1.5 milligrams) as a preservative. The chemical name for Meloxicam is 4-Hydroxy-2-methyl-N- (5-methyl-2-thiazolyl)-2H-1,2-benzothiazine-3-carboxamide-1, 1-dioxide. The formulation is a yellowish viscous suspension.



**Indications:**

Loxicom oral suspension is indicated for the control of pain and inflammation associated with osteoarthritis in dogs.

**Dosage and Administration:**

Always provide client information sheet with prescription. Carefully consider the potential benefits and risks of Loxicom and other treatment options before deciding to use Loxicom. Use the lowest effective dose for the shortest duration consistent with individual response. Loxicom oral suspension should be administered initially at 0.09 mg/lb (0.2 mg/kg) body weight only on the first day of treatment.

For all treatments after day 1, Loxicom oral suspension should be administered once daily at a dose of 0.045 mg/lb (0.1 mg/kg). The provided syringes are calibrated to

deliver the daily maintenance dose in pounds.

### Directions for Administration:

Loxicom® oral suspension 0.5 mg/mL is packaged with 2 sizes of dosing syringes. The small syringe is calibrated for use in dogs 1 lb. to 10 lbs. The large syringe is calibrated for use in dogs 2 lbs. to 30 lbs. **Only administer Loxicom® oral suspension with the provided syringes. The container should never be used as a dropper bottle for administration of Loxicom® oral suspension.**

#### Dogs 1 to 10 pounds (0.45 kg to 4.5 kg)

Shake well before use, then remove cap. Particular care should be given with regard to the accuracy of dosing. **To prevent accidental overdosing of small dogs, administer product on food only, never directly into the mouth.** Carefully measure suspension onto food to assure that the correct dose is given before presentation of the food to the dog. The 1 mL syringe provided with the meloxicam concentration of 0.5 mg/mL cannot be used to measure doses for dogs weighing less than 1 lb (0.45kg).

For dogs 1-2 pounds (0.45 kg to 0.90 kg), Loxicom oral suspension 0.5 mg/mL can be placed directly onto the food using the 1 lb graduation mark on the provided 1 mL syringe.


For dogs between 2 - 10 pounds, Loxicom oral suspension can be given using the 1 mL or 3 mL syringe provided in the package (see dosing procedure in the next column). The syringes fit on to the bottle and have a scale in pounds designed to deliver the daily maintenance dose (0.05 mg/lb or 0.1 mg/kg). When using the syringes, the dog's weight should be rounded down to the nearest 1 pound increment. Replace and tighten cap after use.

#### Dogs over 10 pounds (4.5 kg)

Shake well before use then remove cap. Loxicom oral suspension may be either mixed with food or placed directly into the mouth. Particular care should be given with regard to the accuracy of dosing. Loxicom oral suspension can be given using the 3 mL measuring syringe provided in the package (see dosing procedure in the next column). The syringe fits on to the bottle and has a scale in pounds designed to deliver the daily maintenance dose (0.05 mg/lb or 0.1 mg/kg). When using the syringe, the dog's weight should be rounded down to the nearest 1 pound increment. Replace and tighten cap after use.



*Shake bottle well.  
Push down and  
unscrew bottle top.*



*Turn the  
bottle/syringe upside  
down. Pull the*



*Turn the bottle right  
way up and with a  
twisting movement*



*Push the plunger to  
empty the contents  
of the syringe.*

*Attach the dosing syringe to the bottle by gently pushing onto the top of the bottle. plunger out until the medication level corresponds to the dog's body weight in pounds. separate the dosing syringe from the bottle.*

### **Contraindications:**

Dogs with known hypersensitivity to meloxicam should not receive Loxicom oral suspension. **Do not use Loxicom oral suspension in cats. Acute renal failure and death have been associated with the use of meloxicam in cats.**

### **Warnings:**

Not for use in humans. Keep this and all medications out of reach of children. Consult a physician in case of accidental ingestion by humans. **For oral use in dogs only.**

As with any NSAID all dogs should undergo a thorough history and physical examination before the initiation of NSAID therapy. Appropriate laboratory testing to establish hematological and serum biochemical baseline data is recommended prior to and periodically during administration. Owner should be advised to observe their dog for signs of potential drug toxicity and be given a client information sheet about Loxicom.

Keep Loxicom in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

### **Precautions:**

The safe use of Loxicom oral suspension in dogs younger than 6 months of age, dogs used for breeding, or in pregnant or lactating dogs has not been evaluated. Meloxicam is not recommended for use in dogs with bleeding disorders, as safety has not been established in dogs with these disorders.

As a class, cyclo-oxygenase inhibitory NSAIDs may be associated with gastrointestinal, renal and hepatic toxicity. Sensitivity to drug-associated adverse events varies with the individual patient. Dogs that have experienced adverse reactions from one NSAID may experience adverse reactions from another NSAID. Patients at greatest risk for renal toxicity are those that are dehydrated, on concomitant diuretic therapy, or those with existing renal, cardiovascular, and/or hepatic dysfunction. Concurrent administration of potentially nephrotoxic drugs should be carefully approached. NSAIDs may inhibit the prostaglandins that maintain normal homeostatic function. Such anti-prostaglandin effects may result in clinically significant disease in patients with underlying or pre-existing disease that has not been previously diagnosed. Since NSAIDs possess the potential to induce gastrointestinal ulcerations and/or perforations, concomitant use with other anti-inflammatory drugs, such as NSAIDs or corticosteroids, should be avoided.

If additional pain medication is needed after administration of the total daily dose of Loxicom oral suspension, a non-NSAID or non-corticosteroid class of analgesia should be considered. The use of another NSAID is not recommended. Consider appropriate washout times when switching from corticosteroid use or from one NSAID to another in dogs. The use of concomitantly protein-bound drugs with Loxicom oral suspension has

not been studied in dogs. Commonly used protein-bound drugs include cardiac, anticonvulsant and behavioral medications. The influence of concomitant drugs that may inhibit metabolism of Loxicom oral suspension has not been evaluated. Drug compatibility should be monitored in patients requiring adjunctive therapy.

### **Adverse Reactions:**

Field safety was evaluated in 306 dogs. Based on the results of two studies, GI abnormalities (vomiting, soft stools, diarrhea, and inappetance) were the most common adverse reactions associated with the administration of meloxicam. The following table lists adverse reactions and the numbers of dogs that experienced them during the studies. Dogs may have experienced more than one episode of the adverse reaction during the study.

<b>Adverse Reactions Observed During Two Field Studies</b>		
Clinical Observation	Meloxicam (n=157)	Placebo (n=149)
Vomiting	40	23
Diarrhea/Soft Stool	19	11
Bloody Stool	1	0
Inappetance	5	1
Bleeding gums after dental procedure	1	0
Lethargy/Swollen Carpus	1	0
Epiphora	1	0

In foreign suspected adverse drug reaction (SADR) reporting over a 9 year period, incidences of adverse reactions related to meloxicam administration included: auto-immune hemolytic anemia (1 dog), thrombocytopenia (1 dog), polyarthritis (1 dog), nursing puppy lethargy (1 dog), and pyoderma (1 dog).

### **Post-Approval Experience (Rev. 2010):**

The following adverse events are based on post-approval adverse drug experience reporting. Not all adverse reactions are reported to FDA/CVM. It is not always possible to reliably estimate the adverse event frequency or establish a causal relationship to product exposure using these data. The following adverse events are listed in decreasing order of frequency by body system.

*Gastrointestinal:* vomiting, anorexia, diarrhea, melena, gastrointestinal ulceration

*Urinary:* azotemia, elevated creatinine, renal failure

*Neurological/Behavioral:* lethargy, depression

*Hepatic:* elevated liver enzymes

*Dermatologic:* pruritus

Death has been reported as an outcome of the adverse events listed above. **Acute renal failure and death have been associated with use of meloxicam in cats.**

## CONTACT INFORMATION

To report suspected adverse drug experiences, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Norbrook at 1-866-591-5777. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at [www.fda.gov/reportanimalae](http://www.fda.gov/reportanimalae).

## Information for Dog Owners

Loxicom, like other drugs of its class, is not free from adverse reactions. Owners should be advised of the potential for adverse reactions and be informed of the clinical signs associated with drug intolerance. Adverse reactions may include vomiting, diarrhea, decreased appetite, dark or tarry stools, increased water consumption, increased urination, pale gums due to anemia, yellowing of gums, skin or white of the eye due to jaundice, lethargy, incoordination, seizure, or behavioral changes. **Serious adverse reactions associated with this drug class can occur without warning and in rare situations result in death (see Adverse Reactions). Owners should be advised to discontinue Loxicom and contact their veterinarian immediately if signs of intolerance are observed.**

The vast majority of patients with drug related adverse reactions have recovered when the signs are recognized, the drug is withdrawn, and veterinary care, if appropriate, is initiated. Owners should be advised of the importance of periodic follow up for all dogs during administration of any NSAID.

## Clinical Pharmacology:

Meloxicam has nearly 100% bioavailability when administered orally with food. The terminal elimination half-life after a single dose is estimated to be approximately 24 hrs (+/-30%) regardless of route of administration. There is no evidence of statistically significant gender differences in drug pharmacokinetics. Drug bioavailability, volume of distribution, and total systemic clearance remain constant up to 5 times the recommended dose for use in dogs. However, there is some evidence of enhanced drug accumulation and terminal elimination half-life prolongation when dogs are dosed for 45 days or longer. Peak drug concentrations can be expected to occur within about 7.5 hrs after oral administration. Corresponding peak concentration is approximately 0.464 mcg/mL following a 0.2 mg/kg oral dose. The drug is 97% bound to canine plasma proteins.

## Effectiveness:

The effectiveness of meloxicam was demonstrated in two field studies involving a total of 277 dogs representing various breeds, between six months and sixteen years of age, all diagnosed with osteoarthritis. Both of the placebo-controlled, masked studies were conducted for 14 days. All dogs received 0.2 mg/kg on day 1. All dogs were maintained on 0.1 mg/kg oral meloxicam from days 2 through 14 of both studies. Parameters evaluated by veterinarians included lameness, weight-bearing, pain on palpation, and overall improvement. Parameters assessed by owners included mobility, ability to rise, limping, and overall improvement.

In the first field study (n=109), dogs showed clinical improvement with statistical significance after 14 days of meloxicam treatment for all parameters. In the second field study (n=48), dogs receiving meloxicam showed a clinical improvement after 14 days of therapy for all parameters; however, statistical significance was demonstrated only for the overall investigator evaluation on day 7, and for the owner evaluation on day 14.

## **Safety:**

### Six Week Study

In a six week target animal safety study, meloxicam was administered orally at 1, 3, and 5X the recommended dose with no significant clinical adverse reactions. Animals in all dose groups (control, 1, 3 and 5X the recommended dose) exhibited some gastrointestinal distress (diarrhea and vomiting). No treatment-related changes were observed in hematological, blood chemistry, urinalysis, clotting time, or buccal mucosal bleeding times.

Necropsy results included stomach mucosal petechiae in one control dog, two dogs at the 3X and one dog at the 5X dose. Other macroscopic changes included areas of congestion or depression of the mucosa of the jejunum or ileum in three dogs at the 1X dose and in two dogs at the 5X dose. Similar changes were also seen in two dogs in the control group. There were no macroscopic small intestinal lesions observed in dogs receiving the 3X dose. Renal enlargement was reported during the necropsy of two dogs receiving the 3X and two receiving the 5X dose.

Microscopic examination of the kidneys revealed minimal degeneration or slight necrosis at the tip of the papilla in three dogs at the 5X dose. Microscopic examination of the stomach showed inflammatory mucosal lesions, epithelial regenerative hyperplasia or atrophy, and submucosal gland inflammation in two dogs at the recommended dose, three dogs at the 3X and four dogs at the 5X dose. Small intestinal microscopic changes included minimal focal mucosal erosion affecting the villi, and were sometimes associated with mucosal congestion. These lesions were observed in the ileum of one control dog and in the jejunum of one dog at the recommended dose and two dogs at the 5X dose.

### Six Month Study

In a six month target animal safety study, meloxicam was administered orally at 1, 3, and 5X the recommended dose with no significant clinical adverse reactions. All animals in all dose groups (controls, 1, 3, and 5X the recommended dose) exhibited some gastrointestinal distress (diarrhea and vomiting). Treatment related changes seen in hematology and chemistry included decreased red blood cell counts in seven of 24 dogs (four 3X and three 5X dogs), decreased hematocrit in 18 of 24 dogs (including three control dogs), dose-related neutrophilia in one 1X, two 3X and three 5X dogs, evidence of regenerative anemia in two 3X and one 5X dog. Also noted were increased BUN in two 5X dogs and decreased albumin in one 5X dog.

Endoscopic changes consisted of reddening of the gastric mucosal surface covering less than 25% of the surface area. This was seen in three dogs at the recommended dose, three dogs at the 3X dose and two dogs at the 5X dose. Two control dogs exhibited reddening in conjunction with ulceration of the mucosa covering less than 25% of the surface area.

Gross gastrointestinal necropsy results observed included mild discoloration of the

stomach or duodenum in one dog at the 3X and in one dog at the 5X dose. Multifocal pinpoint red foci were observed in the gastric fundic mucosa in one dog at the recommended dose, and in one dog at the 5X dose.

No macroscopic or microscopic renal changes were observed in any dogs receiving meloxicam in this six month study.

Microscopic gastrointestinal findings were limited to one dog at the recommended dose, and two dogs at the 3X dose. Mild inflammatory mucosal infiltrate was observed in the duodenum of one dog at the recommended dose. Mild congestion of the fundic mucosa and mild myositis of the outer mural musculature of the stomach were observed in two dogs receiving the 3X dose.

### **How Supplied:**

Loxicom oral suspension 0.5 mg/mL: 15 mL and 30 mL bottles with 1 mL and 3 mL measuring syringes.

NDC 55529-042-13 - 0.5 mg/mL - 15 mL

NDC 55529-042-14 - 0.5 mg/mL - 30 mL

**Storage:** Store at controlled room temperature 68-77°F (20-25°C). Excursions permitted between 59°F and 86°F (15°C and 30°C). Brief exposure to temperature up to 104°F (40°C) may be tolerated provided the mean kinetic temperature does not exceed 77°F (25°C); however such exposure should be minimized.

Approved by FDA under ANADA # 200-786

Made in the UK.

Manufactured by:  
Norbrook Laboratories Limited  
Newry, BT35 6QQ, Co. Down, Northern Ireland

Loxicom® is a registered trademark of Norbrook Laboratories Limited

U.S. Patent No. 9,399,013

### **TAKE TIME**



### **OBSERVE LABEL**

### **DIRECTIONS**

Rev. 02/26

624670I02

**Norbrook®**

### **Client Information Sheet**

**Loxicom®**  
**(meloxicam oral suspension)**

## **0.5 mg/mL**

Non-steroidal anti-inflammatory drug for oral use in dogs only.

This summary contains important information about Loxicom<sup>®</sup>. You should read this information before you start giving your dog Loxicom and review it each time the prescription is refilled. This sheet is provided only as a summary and does not take the place of instructions from your veterinarian. Talk to your veterinarian if you do not understand any of this information or if you want to know more about Loxicom.

### **What is Loxicom?**

Loxicom is a prescription non-steroidal anti-inflammatory drug (NSAID) that is used to control pain and inflammation (soreness) due to osteoarthritis in dogs. Osteoarthritis (OA) is a painful condition caused by "wear and tear" of cartilage and other parts of the joints that may result in the following changes or signs in your dog: Limping or lameness, decreased activity or exercise (reluctance to stand, climb stairs, jump or run, or difficulty in performing these activities), stiffness or decreased movement of joints. Loxicom is given to dogs by mouth. **Do not use Loxicom oral suspension in cats. Acute renal failure and death have been associated with the use of meloxicam in cats.**

### **What Kind Of Results Can I Expect When My Dog Is On Loxicom For OA?**

While Loxicom is not a cure for osteoarthritis, it can control the pain and inflammation of OA and improve your dog's mobility.

- Response varies from dog to dog but can be quite dramatic.
- In most dogs, improvement can be seen in a matter of days.
- If Loxicom is discontinued or not given as directed, your dog's pain and inflammation may come back.

### **What Dogs Should Not Take Loxicom?**

Your dog should not be given Loxicom if he/she:

- Has had an allergic reaction to meloxicam, the active ingredient of Loxicom.
- Has had an allergic reaction (such as hives, facial swelling, or red or itchy skin) to aspirin or other NSAIDs.
- Is presently taking aspirin, other NSAIDs, or corticosteroids (unless directed by your veterinarian).

### **Loxicom Should Only Be Given To Dogs**

People should not take Loxicom. Keep Loxicom and all medication out of reach of children. Call your physician immediately if you accidentally take Loxicom.

Keep Loxicom in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

### **How To Give Loxicom To Your Dog**

The actual dose to be given should be prescribed by the veterinarian.

### **Directions for Administration:**

Loxicom<sup>®</sup> oral suspension 0.5 mg/mL is packaged with 2 sizes of dosing syringes. The small syringe is calibrated for use in dogs 1 lb. to 10 lbs. The large syringe is calibrated

for use in dogs 2 lbs. to 30 lbs.

**Only administer Loxicom® oral suspension with the provided syringes. The container should never be used as a dropper bottle for administration of Loxicom® oral suspension.**

### **Dogs 1 to 10 pounds (0.45 kg to 4.5 kg)**

Shake well before use, then remove cap. Particular care should be given with regard to the accuracy of dosing. **To prevent accidental overdosing of small dogs, administer product on food only, never directly into the mouth.** Carefully measure suspension onto food to assure that the correct dose is given before presentation of the food to the dog. The 1 mL syringe provided with the meloxicam concentration of 0.5 mg/mL cannot be used to measure doses for dogs weighing less than 1 lb (0.45kg).

For dogs 1-2 pounds (0.45 kg to 0.90 kg), Loxicom oral suspension 0.5 mg/mL can be placed directly onto the food using the 1 lb graduation mark on the provided 1 mL syringe.


For dogs between 2 - 10 pounds, Loxicom oral suspension can be given using the 1 mL or 3 mL syringe provided in the package (see dosing procedure below). The syringes fit on to the bottle and have a scale in pounds designed to deliver the daily maintenance dose (0.05 mg/lb or 0.1 mg/kg). When using the syringes, the dog's weight should be rounded down to the nearest 1 pound increment. Replace and tighten cap after use.

### **Dogs over 10 pounds (4.5 kg)**


Shake well before use then remove cap. Loxicom oral suspension may be either mixed with food or placed directly into the mouth. Particular care should be given with regard to the accuracy of dosing. Loxicom oral suspension can be given using the 3 mL measuring syringe provided in the package (see dosing procedure below). The syringe fits on to the bottle and has a scale in pounds designed to deliver the daily maintenance dose (0.05 mg/lb or 0.1 mg/kg). When using the syringe, the dog's weight should be rounded down to the nearest 1 pound increment. Replace and tighten cap after use.



*Shake bottle well. Push down and unscrew bottle top. Attach the dosing syringe to the bottle by gently pushing onto the top of the bottle.*



*Turn the bottle/syringe upside down. Pull the plunger out until the medication level corresponds to the dog's body weight in pounds.*



*Turn the bottle right way up and with a twisting movement separate the dosing syringe from the bottle.*



*Push the plunger to empty the contents of the syringe.*

### **What To Tell/Ask Your Veterinarian Before Giving Loxicom**

Talk to your veterinarian about:

- The signs of OA you have observed (for example limping, stiffness).

- The importance of weight control and exercise in the management of OA.
- What tests might be done before Loxicom is prescribed.
- How often your dog may need to be examined by your veterinarian.
- The risks and benefits of using Loxicom.

Tell your veterinarian if your dog has ever had the following medical problems:

- Experienced side effects from Loxicom or other NSAIDs, such as aspirin
- Digestive upset (vomiting and/or diarrhea)
- Liver disease
- Kidney disease

Tell your veterinarian about:

- Any other medical problems or allergies that your dog has now or has had.
- All medicines that you are giving your dog or plan to give your dog, including those you can get without a prescription.

Tell your veterinarian if your dog is:

- Pregnant, nursing or if you plan to breed your dog.

### **What Are The Possible Side Effects That May Occur In My Dog During Loxicom Therapy?**

Loxicom, like other drugs, may cause some side effects. Serious but rare side effects have been reported in dogs taking NSAIDs. Serious side effects can occur with or without warning and in rare situations result in death.

The most common NSAID-related side effects generally involve the stomach and liver or kidney problems. Look for the following side effects that can indicate your dog may be having a problem with Loxicom or may have another medical problem:

- Decrease or increase in appetite
- Vomiting
- Change in bowel movement (such as diarrhea, or black, tarry or bloody stools)
- Change in behavior (such as decreased or increased activity level, incoordination, seizure or aggression)
- Yellowing of gums, skin, or whites of the eyes (jaundice)
- Change in drinking habits (frequency, amount consumed)
- Change in urination habits (frequency, color, or smell)
- Change in skin (redness, scabs, or scratching)

It is important to stop therapy and contact your veterinarian immediately if you think your dog has a medical problem or side effect from Loxicom therapy. If you have additional questions about possible side effects, talk to your veterinarian.

### **Can Loxicom Be Given With Other Medicines?**

Loxicom should not be given with other NSAIDs (for example, aspirin, carprofen, etodolac, deracoxib) or steroids (for example, cortisone, prednisone, dexamethasone, triamcinolone).

Tell your veterinarian about all medicines you have given your dog in the past, and any medicines that you are planning to give with Loxicom. This should include other medicines that you can get without a prescription. Your veterinarian may want to check

that all of your dog's medicines can be given together.

### **What Can I Do In Case My Dog Eats More Than The Prescribed Amount?**

Contact your veterinarian immediately if your dog eats more than the prescribed amount of Loxicom.

### **What Else Should I Know About Loxicom?**

This sheet provides a summary of information about Loxicom. If you have any questions or concerns about Loxicom or osteoarthritis pain, talk to your veterinarian.

As with all prescribed medicines, Loxicom should only be given to the dog for which it was prescribed. **Loxicom oral suspension is for use in dogs only. Do not give Loxicom oral suspension to cats.** It should be given to your dog only for the condition for which it was prescribed. It is important to periodically discuss your dog's response to Loxicom at regular check ups. Your veterinarian will best determine if your dog is responding as expected and if your dog should continue receiving Loxicom.

### **CONTACT INFORMATION**

To report suspected adverse drug experiences, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Norbrook at 1-866-591-5777. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at [www.fda.gov/reportanimalae](http://www.fda.gov/reportanimalae).

Made in the UK.

Manufactured by:  
Norbrook Laboratories Limited  
Newry, BT35 6QQ, Co. Down,  
Northern Ireland

Loxicom<sup>®</sup> is a registered trademark of Norbrook Laboratories Limited

U.S. Patent No. 9,399,013

### **TAKE TIME**



### **OBSERVE LABEL**

### **DIRECTIONS**

Rev. 02/26

624670102

**Norbrook<sup>®</sup>**

**Principal Display Panel - 15 mL Carton Label**

**NDC 55529-042-13**

**Loxicom<sup>®</sup>  
(meloxicam  
oral suspension)**

**0.5 mg/mL**

**Non-steroidal  
anti-inflammatory  
drug for oral use  
in dogs only.**

**Caution: Federal law  
restricts this drug to  
use by or on the order  
of a licensed veterinarian.**

**Approved by FDA  
under ANADA # 200-786**

**15 mL**

***Norbrook***<sup>®</sup>



**Principal Display Panel - 15 mL Bottle Label**

**Loxicom<sup>®</sup>  
(meloxicam  
oral suspension)**

**0.5 mg/mL**

**Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian. Approved by FDA under ANADA # 200-786**

**15 mL**

**Norbrook®**

<p>If signs of illness are seen, treatment should be discontinued immediately and the advice of a veterinarian sought. Refer to the product insert or client information sheet for detailed information.</p>	<p><b>Loxicom®</b> (meloxicam oral suspension) 0.5 mg/mL</p>	<p>NDC 55529-042-13 Shake well. Refer to the package insert for complete product information.</p>	
<p><b>Warnings:</b> Not for use in humans. Keep out of reach of children.</p> <p><b>Indications:</b> Loxicom oral suspension is indicated for the control of pain and inflammation associated with osteoarthritis in dogs.</p> <p>Made in the UK Norbrook Laboratories Limited, Newry, Co. Down, BT35 6QQ, Northern Ireland</p>		<p><b>Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian. Approved by FDA under ANADA # 200-786</b></p> <p>15 mL</p> 	<p><b>Storage:</b> Store at controlled room temperature 68-77°F (20-25°C); excursions permitted between 59°F and 86°F (15°C and 30°C).</p>
		<p>Lot No.: Exp. Date: 323670101 Rev. 02/25</p>	<p><b>OVERPRINT AREA</b> 14 x 20mm</p>

<b>LOXICOM</b>				
meloxicam suspension				
<b>Product Information</b>				
<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:55529-042	
<b>Route of Administration</b>	ORAL			
<b>Active Ingredient/Active Moiety</b>				
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
	meloxicam (UNII: VG2QF83CGL) (meloxicam - UNII:VG2QF83CGL)	meloxicam	0.5 mg in 1 mL	
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:55529-042-13	1 in 1 CARTON		
1		15 mL in 1 BOTTLE, PLASTIC		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	

ANADA	ANADA200786	05/07/2026	
-------	-------------	------------	--

**Labeler -** Norbrook Laboratories Limited (214580029)

**Establishment**

Name	Address	ID/FEI	Business Operations
Norbrook Laboratories Limited		211218325	API MANUFACTURE, MANUFACTURE, ANALYSIS, PACK, LABEL

Revised: 5/2026

Norbrook Laboratories Limited