## MECADOX 11%- carbadox powder Phibro Animal Health

-----

Mecadox 11%

# **DOSAGE AND ADMINISTRATION**

## DOSAGE:

For the treatment of swine dysentery, salmonellosis, colibacillosis, ileitis and other bacterial swine enteritis, add 500 g of Mecadox ® 11%/ton of finished feed (55 ppm).

# **ADMINISTRATION ROUTE**

Oral, mixed into finished feed.

# **PRECAUTIONS:**

To ensure good mixing, the product should be premixed with a small amount of feed prior to mixing with the total ration.

## WARNING:

This product is for veterinary use only.

The administration of this product is restricted to oral use only in the starter phase of swine.

Consumption of products from these animals is prohibited. Later, in the final stage of development, they may be suitable for human consumption.

Withdrawal Period: 42 days.

Its use is prohibited for all adult swine and all ages of other species intended for human consumption.

Operators should wear safety goggles, protective clothing and a respirator when mixing and handling the finished feed.

Keep out of the reach of children and pets.

# **STORAGE CONDITIONS:**

Store the product in a cool, dry place. Do not expose the product to direct and continuous sunlight.

# **DIRECTIONS FOR USE:**

For the treatment of infectious processes of the digestive tract in swine caused by bacteria susceptible to Carbadox, such as:

Brachyspira hyodysenteriae, Lawsonia intracellularis, Salmonella spp. and Escherichia coli.

## Label

۲

## **CONSULT YOUR VETERINARIAN Sale under medical prescription** LOT NO.: DATE OF MANUFACTURE: EXPIRATION DATE:

Mecadox 11 (Mexico) 20KG (MFG USA) 9x14 101-8385-01 10297CR2.qxp 7/15/22 7:23 AM Page 1

# Mecadox®11%

## Registration Number: Q-0259-023 VETERINARY USE ONLY Medicated Premix

COMPOSITION:

Each 1000 g contains:

Carbadox..... Excipients o.s...

## DIRECTIONS FOR USE:

For the treatment of infectious processes of the digestive tract in swine caused by bacteria susceptible to Carbadox, such as: Brachyspira hyodysenteriae, Lawsonia intracellularis, Salmonella spp. and Escherichia coll.

#### DOSAGE:

For the treatment of swine dysentery, salmonellosis, colloacillosis, lieitis and other bacterial swine enteritis, add 500 g of Mecadoxs11%/ton of finished feed (55 ppm).

## ADMINISTRATION ROUTE

Oral, mixed into finished feed.

### WARNING:

This product is for veterinary use only. The administration of this product is restricted to oral use only in the starter phase of swine. Consumption of products from these animals is prohibited. Later, in the final stage of development, they may be suitable for human consumption.

Withdrawal Period: 42 days.

Its use is prohibited for all adult swine and all ages of other species intended for human consumption.

Operators should wear safety goggles, protective clothing and a respirator when mixing and handling the finished feed. Keep out of the reach of children and pets.

### PRECAUTIONS:

To ensure good mixing, the product should be premixed with a small amount of feed prior to mixing with the total ration.

## STORAGE CONDITIONS:

Store the product in a cool, dry place. Do not expose the product to direct and continuous sunlight.

## CONSULT YOUR VETERINARIAN Sale under medical prescription

LOT NO .:

DATE OF MANUFACTURE:

EXPIRATION DATE:

MANUFACTURED BY: International Nutrition Inc. PO Box 27540 7706 I Plaza Omaha, NE 68127 USA IMPORTED AND DISTRIBUTED BY: PB Animal Health de México S, de R.L. de C.V Francisco de Quevedo #117 Torre A Piso 6 Col. Arcos Vallarta ZIP 44130, Guadalajara, Jal. Phone: 33.3818-0529


(i)

Mecadox is a trademark of Phibro Animal Health Corporation



NET CONTENT: 20 Kg

101-8385-01



Labeler - Phibro Animal Health (006989008)

**Registrant -** Phibro Animal Health (006989008)

Revised: 4/2025

849% (7459)

Phibro Animal Health