

**INFANTS DYE FREE IBUPROFEN- ibuprofen suspension/ drops**  
**Walgreen Company**

-----  
**Walgreen Co. Infants Dye-Free Ibuprofen Drug Facts**

**Active ingredient (in each 1.25 mL)**

Ibuprofen 50 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- give with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- mL = milliliter
- measure with the dosing device provided. Do not use with any other device.
- dispense liquid slowly into the child's mouth, toward the inner cheek
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**

### Dosing Chart

Weight (lbs)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

## Other information

- store at 20-25°C (68-77°F)
- do not freeze
- **do not use if printed neckband is broken or missing**

## Inactive ingredients

anhydrous citric acid, glycerin, hypromellose, natural and artificial berry flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

## Questions or comments?

**1-800-719-9260**

## Principal Display Panel

Compare to Motrin® Infants' Drops active ingredient

infants' AGES 6 TO 23 MONTHS

Dye-Free Ibuprofen

CONCENTRATED DROPS

IBUPROFEN ORAL SUSPENSION 50 mg PER 1.25 mL

PAIN RELIEVER / FEVER REDUCER (NSAID)

DYE FREE

LASTS UP TO 8 HOURS

Non-staining

No high fructose corn syrup

BERRY FLAVOR

AGES 6 TO 23 MONTHS

1 FL OZ (30 mL)

Use only with enclosed syringe.

SEE NEW WARNINGS

**Walgreens**  
Compare to Motrin® Infants' Drops active ingredient!<sup>†</sup>  
NDC 0363-2323-10  
**infants'**  
AGES 6 TO 23 MONTHS  
**Dye-Free Ibuprofen**  
CONCENTRATED DROPS  
IBUPROFEN ORAL SUSPENSION  
50 mg PER 1.25 mL  
PAIN RELIEVER / FEVER REDUCER (NSAID)

**DYE FREE**  
LASTS UP TO 8 HOURS

- Non-staining
- No high fructose corn syrup

**AGES 6 TO 23 MONTHS**

**BERRY FLAVOR**

1 FL OZ (30 mL)

Use only with enclosed syringe.

**SEE NEW WARNINGS**

**PHARMACIST RECOMMENDED**  
Walgreens  
Trusted since 1957™  
Health expertise you rely on.™

**Drug Facts**  
**Active ingredient (in each 1.25 mL)** Ibuprofen 50 mg (NSAID)  
**Purposes** Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses temporarily:** ■ reduces fever  
■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Drug Facts (continued)**  
**Warnings**  
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives ■ facial swelling ■ asthma (wheeze/zing)  
■ shock ■ skin reddening ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:  
■ has had stomach ulcers or bleeding problems  
■ takes a blood thinning (anticoagulant) or stomach drug (aspirin, ibuprofen, naproxen, or others)  
■ takes more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**  
■ if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer  
■ right before or after heart surgery

**Ask a doctor before use if**  
■ stomach bleeding warning applies to your child  
■ child has a history of stomach problems, such as heartburn  
■ child has problems or serious side effects from taking pain relievers or fever reducers  
■ child has not been drinking fluids  
■ child has lost a lot of fluid due to vomiting or diarrhea  
■ child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke  
■ child has asthma ■ child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**  
■ under a doctor's care for any serious condition  
■ taking any other drug

**When using this product**  
■ give with food or milk if stomach upset occurs

**Stop use and ask a doctor if**  
■ child experiences any of the following signs of stomach bleeding:  
■ feels faint ■ vomits blood ■ has bloody or black stools  
■ has stomach pain that does not get better  
■ child has symptoms of heart problems or stroke:  
■ chest pain ■ trouble breathing  
■ weakness in one part or side of body  
■ slurred speech ■ leg swelling  
■ the child does not get any relief within first day (24 hours) of treatment  
■ fever or pain gets worse or lasts more than 3 days  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

**Drug Facts (continued)**  
**Directions**  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

■ this product does not contain directions or complete warnings for adult use  
■ do not give more than directed  
■ shake well before using  
■ find right dose on chart below. If possible, use weight to dose; otherwise use age.  
■ mL = milliliter  
■ measure with the dosing device provided. Do not use with any other device.  
■ dispense liquid slowly into the child's mouth, toward the inner cheek  
■ if needed, repeat dose every 6-8 hours  
■ do not use more than 4 times a day

**Dosing Chart**

Weight (lbs)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

**Other information**  
■ store at 20-25°C (68-77°F)  
■ do not freeze  
■ do not use if printed neckband is broken or missing

**Inactive ingredients**  
anhydrous citric acid, glycerin, hydroxyethylcellulose, natural and artificial berry flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

**Questions or comments?**  
1-800-719-9260

<sup>†</sup>Walgreens Pharmacist Survey

<sup>H</sup>This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Motrin® Infants' Drops.

ORG0918-F REV-0719

DISTRIBUTED BY: WALGREEN CO. ITEM 807003 W10106-1019-L  
200 WALGATE RD, DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com  
©2018 Walgreen Co.

3 11917 19545 2

25510 94 C8

## INFANTS' DYE FREE IBUPROFEN

ibuprofen suspension/ drops

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-2323
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	50 mg in 1.25 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
WATER (UNII: 059QF0K00R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

**Product Characteristics**

<b>Color</b>	WHITE (Opaque)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY (mixed)	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-2323-10	1 in 1 CARTON	08/23/2018	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA075217	08/23/2018	

**Labeler** - Walgreen Company (008965063)